Southern Illinois University Carbondale **OpenSIUC**

Final Reports

Cooperative Wildlife Research Laboratory

8-1995

Population Dynamics and Ecology of White-Tailed Deer in Illinois

Alan Woolf Southern Illinois University Carbondale

John L. Roseberry Southern Illinois University Carbondale

Follow this and additional works at: http://opensiuc.lib.siu.edu/cwrl_fr W-87-R-17.

Recommended Citation

Woolf, Alan and Roseberry, John L., "Population Dynamics and Ecology of White-Tailed Deer in Illinois" (1995). Final Reports. Paper 1.

http://opensiuc.lib.siu.edu/cwrl_fr/1

This Article is brought to you for free and open access by the Cooperative Wildlife Research Laboratory at OpenSIUC. It has been accepted for inclusion in Final Reports by an authorized administrator of OpenSIUC. For more information, please contact opensiuc@lib.siu.edu.

FINAL REPORT

W-87-R-17

Population Dynamics and Ecology of White-Tailed Deer in Illinois

Submitted by:

Cooperative Wildlife Research Laboratory, SIUC

Presented to:

Illinois Department of Natural Resources

August 1995

Prepared by: Submitted by:

Alan Woolf, Director Coop. Wildlife Research Lab. Southern Illinois University Carbondale, IL 62901 Alan Woolf, Director Coop. Wildlife Research Lab. Southern Illinois University Carbondale, IL 62901 John L. Roseberry, Senior Scientist Coop. Wildlife Research Lab. Southern Illinois University Carbondale, IL 62901

TABLE OF CONTENTS

	Page
Need	. 1
Objectives	. 2
Executive Summary	. 3
JOB A. Habitat Inventory, Classification, and Analysis	. 5
Objectives	
Methods	. 7
Data Sets	
TIGER/Line(TM) Census Files	
USGS Topographical Maps	. 8
Classification Scheme	. 9
Classification Procedures	
Results and Discussion	15
JOB B. Deer Ecology and Life History in Westcentral and	.
Northern Illinois	53
Objectives	53
JOB C. Population Analysis	54
Objectives	54
Introduction	54 54
Results	55
JOB D. Analyze and Report	57
Objectives	57

- Illinois Deer Harvest Analysis Program (IDHAP)
 Version 1.1 User's Guide and Reference Manual
- Illinois Deer Harvest Analysis and Modeling Program (IDHAMP)
 Version 1.1 User's Guide and Reference Manual
- Illinois Deer Harvest Analysis and Modeling Program (IDHAMP)
 Programmer's Guide Version 1.1

LIST OF TABLES

Table 1	Data on Landsat 5 TM scenes used to classify each
	Illinois county
Table 2	Summary of land use classification accuracy assessment and percent errors of omission (O) and commission (C) for Illinois counties classified from Landsat 5 TM
	satellite data
Table 3	Accuracy assessment and error estimate samples pooled
	for an overall statewide estimate of accuracy 25

LIST OF FIGURES

Figure	1.	Satellite	coverage	e of 1	[llinois	provi	.ded	by	the	<u>,</u>		
		Illinois	Natural B	Histor	ry Survey	<i>y</i>						26

LIST OF APPENDICES

Appendix A.	Counties with greater than 48 ha of orchards were
	digitized using USGS 1:24000-scale topographical
	quadrangle mapssupplied by Morris Library. These
	23 counties represent 73% of land in orchards.
	Cumberland County quad maps were unavailable (51.8
	ha). The orchard vectors were merged into a single
	vector and then converted to a raster for each
	county
Appendix B.	Accuracy assessment and error estimates for Illinois
	counties classified from Landsat 5 TM satellite
	data

FINAL REPORT

STATE OF ILLINOIS

W-87-R-17

Project Period: 1 July 1992 through 30 June 1995

Study: Population dynamics and ecology of white-tailed deer in

Illinois

Prepared by Alan Woolf and John L. Roseberry Cooperative Wildlife Research Laboratory Southern Illinois University at Carbondale

Since modern firearm hunting of deer resumed in Illinois in 1957, county harvest quotas have been based on retrospective interpretation of the previous year's harvest data. Decisions are made in the absence of definitive knowledge of deer habitat distribution, quantity or quality. The productivity of Illinois deer coupled with a generally conservative approach to harvest management has permitted dramatic herd growth in most counties and regions. It is now apparent that more aggressive and innovative harvest strategies will be necessary to effectively manipulate the size and composition of these populations. ensure that these new harvest regimes are both safe and effective, there will need to be more sophisticated and detailed analysis of harvest data as well as the ability to predict and evaluate the effects of proposed and implemented harvest strategies on size and composition of the deer herd and behavior

and performance of hunters. At present, the Illinois deer management program lacks these capabilities.

In addition, more attention must be given to the amount, distribution, and quality of deer habitat on a county and regional basis and how this habitat is spatially oriented in relation to road systems, human habitation, and sensitive agricultural areas. At the local level, habitat variables can often be measured directly. However, special problems arise in the acquisition, storage, analysis, and interpretation of habitat data on a county, regional, or statewide scale. Fortunately, recent advances in remote sensing, geographic information systems (GIS), and habitat modelling offer solutions to these problems. Progressive management of the Illinois deer herd requires that these capabilities be developed and utilized.

Objectives:

- 1. To assess the amount, distribution, and quality of whitetailed deer habitat in Illinois.
- 2. To relate spatial aspects of deer habitat to other important attributes such as hunter access, proximity to human habitation, and agricultural patterns.
- 3. To complete ongoing studies describing current natality rates, fawn recruitment, seasonal movements, and seasonal and annual mortality rates for previously marked deer in westcentral and northern Illinois.

4. To develop interactive, menu-driven, portable computer models and software packages to facilitate analysis of harvest data, predict effects of alternative harvest regimes, and help select appropriate strategies to achieve specific goals and objectives.

EXECUTIVE SUMMARY

This Study is a component of the Grant Agreement entitled

White-tailed Deer Project. The project represents a cooperative

effort between staff of the Illinois Natural History Survey

(INHS), Center for Wildlife Ecology and the Cooperative Wildlife

Research Laboratory at Southern Illinois University at Carbondale

(SIUC). One study objective (Objective 3 above) was the sole

responsibility of INHS staff and the results of that task (Job B,

Deer Ecology and Life History in Westcentral and Northern

Illinois) will be reported separately by the INHS.

Job A (Habitat Inventory, Classification, and Analysis) of this study was designed as a fully collaborative effort. We planned to develop the classification from statewide coverage of Landsat TM scenes selected by INHS staff. Also, we assumed that INHS staff with expertise in classification of Landsat TM scenes would be active participants and would lend technical support to this task. Indeed, we anticipated that a complete classification would be available by the end of Segment 15 so we could focus resources on other tasks planned under Job A. Further, because the land use/land cover classification of the entire state also was an important need of W-106-R-6 (Cooperative Upland Studies),

there was shared responsibility for the task between both projects.

Because of unanticipated staff departures it became clear that the INHS would not participate in developing a statewide classification, and SIUC staff would have to assume sole responsibility. Following consultations with Illinois Department of Natural Resources (IDNR) Program Managers, we made completion of a statewide classification the priority objective of Job A and used all available resources to complete that task. The scope and complexity of producing a land cover classification for Illinois using PC-based software was further complicated by limitations of the Landsat TM data available for the project thru the INHS. This added problem was not resolved until Segment 17 when SIUC's Morris Library acquired Landsat scenes of most portions of Illinois that were better suited for land cover classification than the original scenes we had to work with.

As a result of these unanticipated complications, all planned objectives of Job A were not accomplished. Job A objectives 1 and 2 (the primary objectives agreed to by IDNR Program Managers and project Principal Investigators) were completed and there is now a statewide land cover classification of Illinois available. Job A objective 3 was to be addressed as

a portion of the narrative for Job B reported by the INHS. The lack of data (e.g. the statewide classification) precluded addressing objectives 4 and 5.

All objectives of Job C were completed. In fact, we developed 2 software packages; 1 to archive and analyze deer harvest data, and another that can access and analyze harvest data, but also can model and/or simulate herd performance. The latter software is especially useful for proactive planning and herd management.

In conclusion, accomplishments during this Grant Agreement period have been substantial. Although not all objectives of Job A were completed, prioritization of resources and effort allowed the development of a statewide land cover classification. This was a major accomplishment that will serve the information and management needs of multiple programs within the Illinois Department of Natural Resources.

This project final report describes what was accomplished and methods used to develop the products described. However, the results of this project are, in fact, products not included in the report text. The classified imagery is one major product. It is stored on optical disks at the Cooperative Wildlife Research Laboratory as individual county files. The Laboratory

will use the imagery to support IDNR information needs during subsequent project segments. The other major products are 2 software programs to serve information and management needs of the IDNR Forest Wildlife Program. Two user's manuals describing the software and a programmer's guide are appended to this report.

JOB A. Habitat Inventory, Classification, and Analysis

Objectives: (1) To investigate alternative techniques for classifying white-tailed deer habitat from remote sensing data; (2) to use these techniques and data sources to inventory deer habitat in Illinois; (3) to describe the habitat characteristics of sites selected by dispersing deer and to compare these characteristics with the habitats available within the boundaries of known dispersals from marking sites in northern, westcentral, and eastcentral Illinois; (4) to develop HSI models for the purpose of assessing the relative quality of deer habitat using digital land use classifications from remotely sensed data; and (5) to integrate information relating to spatial distribution of

habitat with other pertinent attributes relating to hunter access, human habitat, and agricultural patterns.

INTRODUCTION

The powerful tools of remote sensing and geographical information system (GIS) software afford resource managers opportunity to develop large data bases that can provide the quantity and quality of information needed for resource management at landscape scales. A land use/land cover classification is a prerequisite foundation upon which resource managers can build GIS data layers that describe location and relative quality of habitat and its spatial relationships to potentially sensitive human developments and agricultural areas.

Our investigations of alternative techniques for classifying white-tailed deer habitat from remote sensing data quickly revealed that only satellite imagery afforded adequate spatial coverage. Landsat 5 TM scenes with 30 m pixel resolution offered an optimum scale (extent and grain) for our proposed classification. The fortuitous availability of statewide Landsat 5 coverage for project use made that the imagery of choice. The number of classes that could be identified in the processed image was considered, and we concluded that a classification consisting

of 6 classes would be appropriate. Generally, the complexity of our classification is similar to that described as Level 1 by Anderson et al. (1976). Finally, we considered the implications of accuracy of the final classification and set a goal of $\geq 90\%$ overall classification accuracy.

METHODS

Data Sets

Landsat 5 TM Data. -- Initially, satellite imagery purchased by the Illinois Natural History Survey was used for this project. This satellite coverage of Illinois consists of 9 full and 2 quarter scenes that are geographically referenced, terrain corrected and mosaicable. The INHS scene dates range from May 1988 to June 1991 (Fig. 1). Data were acquired in TM Fast format on 8 mm Exabyte tapes from INHS. Many of these scenes were difficult to classify which led to a large amount of confusion between several land use classes.

For these confused areas, we used additional satellite imagery owned by the Cooperative Wildlife Research Laboratory (CWRL), and SIUC's Morris Library. These additional scenes range in date from 10 September 1992 to 3 October 1993. Each county

classified, the satellite imagery used, and the scene dates are listed in Table 1.

TIGER/Line(TM) Census Files. -- TIGER (Topographically

Integrated Geographic Encoding and Reference System) is a digital map base used to support Census Bureau programs. The 1990 Census TIGER/Line files contain digital data for features such as streets, rivers and streams, railroads, and political boundaries. The TIGER data set for all Illinois counties was acquired from SIUC's Morris Library.

Aerial Photography. -- National Aerial Photography Program

(NAPP) 1988, black and white, 1:40,000 scale positive prints

available for all of Illinois was the main source for

verification of land use classes. A complete set of photographs

was made available for the project by Morris Library at SIUC.

Color infrared prints from the CWRL map library were used to

verify classifications in some areas. Sets of NAPP black and

white contact print enlargements (3x) that covered bobwhite and

pheasant call count routes in 90 counties were provided by the

IDNR. These photos, with accompanying ground truth information,

were used for accuracy assessments of each county's satellite

classification.

<u>USGS</u> <u>Topographical</u> <u>Maps</u>.--U.S. Geological Survey 7.5-minute quadrangle maps were obtained from both SIUC's Morris Library and the CWRL map library. The most recent copies of these maps were used to identify and digitize orchards in counties that had significant amounts of this land use type. In some cases they were also used to verify other land use types.

Classification Scheme

Six land cover types were identified (crop, woods, grass, water, developed, and orchards) generally similar to the complexity of a classification described as Level 1 by Anderson et al. (1976). Crops included all cultivated acreage in row crops and small grains, as well as miscellaneous cultivated crops. Woods included coniferous forest, deciduous forest, and late old fields. Grass included hay, pasture, fallow fields, Conservation Reserve Program (CRP) fields, field edges, lawns, roadsides, and any other herbaceous cover not considered crop. Water included lakes, rivers, ponds, and other areas with permanent water. Developed areas were defined using TIGER data. Orchards were defined from topographic maps.

Preprocessing

Preprocessing Landsat TM data consisted of importing raw files into Map and Image Processing System (MIPS, MicroImages Inc., Lincoln, NE) software (Miller et al. 1989). Polygons defining broad natural divisions adapted from Schwegman (1973) were digitized over satellite imagery and plotted to binary rasters coregistered to the original scene. These binary rasters were used to extract natural divisions within each scene to limit spectral variability for automatic classification. Cumulus clouds and their shadows were delineated with polygons using the satellite images as a reference. The polygons were plotted to coregistered binary rasters which were used to remove them from the classification to also limit spectral variability.

A principal components analysis was performed on bands 1, 2, and 3 for each natural division and the first principal component (PC1) was saved. A Normalized Difference Vegetation Index (NDVI) also was calculated (Lillesand and Kiefer 1987). Spectral Bands 4, 5, 7, PC1 and NDVI were input in MIPS to an ISOCLASS algorithm which generated approximately 200 ISOCLASS output classes.

TIGER data were processed to provide coregistered maps of developed areas and primary and secondary highways which were combined with the isoclassification of the satellite data.

Developed areas were delineated using neighborhood roads

extracted from TIGER files and plotted to a false color composite image of each scene. A vector of the Illinois state boundary was edited over this image and polygons were drawn around clusters of neighborhood roads. These polygons, which represented developed areas, were cross referenced with hard copy maps of Illinois and defined the "developed" habitat class. Visible quarries and other development areas were also identified in this fashion and put into the developed class.

Primary and secondary highways were extracted from TIGER files for each county within a scene and mosaicked. Primary highways were plotted to a coregistered binary raster and given a width of 2 pixels and digital value of 1. Secondary roads were plotted to the same binary raster, given the same digital value, but with a width of 1 pixel. The polygons representing developed areas were also plotted using this binary raster with a value of 1 on the inside. This binary raster, representing highways and developed areas, was combined with the results of the isoclassification so that developed areas and highways had a digital value of 205 and a distinct color.

Orchards could not be classified using the satellite imagery, but they were identified from USGS 7.5-minute quadrangle maps. The 1992 agricultural census (U.S. Dep. of Commerce 1994)

revealed that 23 counties that had more than 48 ha of orchards accounted for 73% of Illinois orchards. Orchards were manually digitized using the maps for these 23 counties (Appendix A) and combined with their respective county classification.

Classification Procedures

After the initial automatic classifications were run and the ancillary TIGER data added, results (approximately 200 isoclasses) were grouped into meaningful information classes or land use types. We tested 2 methods for lumping classes. The first used 1-mile² (2.59-km²) sample boxes systematically placed over the satellite scene at 10-mile (16.1-km) intervals for ground truth data acquisition. Sample boxes then were delineated on acetate overlays for NAPP black and white positive prints. The land-use classes interpreted within each sample box were drawn onto the acetate overlay.

The land use for each sample box was then digitized to a coregistered raster using the raw satellite image as a reference. These data were used to lump the isoclasses into information classes based on the correlation of the ISOCLASS values to the digital ground truth data within each sample box. This method

proved to be time prohibitive for the large area to be classified and was discontinued.

Alternatively, we examined individual ISOCLASS values for a particular scene and natural division, and used photo verification to determine the most prevalent land use class associated with each ISOCLASS value. Color for each ISOCLASS value was assigned to emulate a false color image. Each ISOCLASS was then flashed and the most prevalent land use associated with that ISOCLASS value was determined. Information about the amount and type of confusion associated with each ISOCLASS value was recorded. After extensive photo verification, we found that many of the resulting ISOCLASS values were confused between 2 or more land use classes.

Each natural division of each scene was reviewed and only ISOCLASSES that showed little or no confusion were retained. A binary raster was created indicating the location of all the confused ISOCLASSES for each scene and natural division. These areas were then re-classified using a maximum likelihood classifier to calculate the statistical probability of a given pixel belonging to a particular land use class (Lillesand and Kiefer 1987). The results of this classifier were combined with the unconfused results from the isoclassification and evaluated.

As an initial accuracy assessment, the percent of land cover types for each extracted county was compared to existing estimates of land use. If these estimates varied from our classification by >10% the county was evaluated using aerial photos.

Even after reclassifying these scenes, acceptable accuracies were not attained for many counties. These counties were then individually re-classified, using only the maximum likelihood classifier. Counties that were not yet processed also were individually classified using the maximum likelihood classifier. Different satellite imagery, owned by the CWRL and SIUC's Morris Library, was then used for counties that still proved difficult to classify accurately. The majority of this imagery were autumn scenes taken in October that were much easier to classify.

After each county was extracted and spectrally classified, a contextual classifier was used to improve the classification.

The contextual classifier was a series of FORTRAN programs that sequentially manipulated the classified output for each county.

The first program detected single isolated pixels of any class and replaced them with the surrounding majority land use class.

The program also removed linear crop features by replacing 1-pixel wide strips of crop, that had grass pixels on opposite

sides with grass. The second program used output from the first to determine size of contiguous crop patches. This program created an output raster with pixel values representing the size of their respective patch. The next program replaced crop patches <13 pixels with the surrounding majority land-use class. The final program replaced any new single pixels created by the previous programs with the surrounding majority land-use class.

Accuracy Assessment

The classification accuracy of the completed counties was assessed using land cover information collected along 0.5-mile (0.8-km) wide and 20-mile (32.25-km) long quail and pheasant call-count routes. The land cover along these routes was identified and mapped on aerial photos by IDNR biologists via field inspections during the summers of 1990 and 1991. Conservation Reserve Program (CRP) enrollment for each county was also recorded along each route.

Representative samples >3 ha of each land cover type were selected on the aerial photographs for each call count route.

Between 50 and 90 samples per county were used depending upon the homogeneity of the landscape and meander of the route. These samples then were located on the classified county image and

their land use types were compared. There were 6 pairs of counties (Gallatin-Hardin, Marshall-Putnam, Jasper-Clay, Stephenson-Winnebago, Carroll-Jo Daviess, and McDonough-Fulton) that had a single route extending across both counties. In these cases, the same set of sample points were used for both county accuracy assessments. An error matrix was constructed for each call count route and used to compute errors of omission and commission, overall accuracy, and the Kappa statistic (Congalton 1991). Accuracy assessment data sampled for each county also were pooled, without the duplicate sample points mentioned above, for statewide estimates.

RESULTS AND DISCUSSION

We classified 99 of 102 Illinois counties (Cook, Du Page, and Lake counties were not classified). The quality (accuracy) of each county classification varied with the quality and acquisition date of the satellite imagery used. Scenes originally acquired from INHS ranged in date from May 26, 1988 to June 10, 1991 and spanned from early spring to summer. The scenes acquired from SIUC's Morris Library ranged only from September 10, 1992 to October 12, 1992 and were much easier to accurately classify.

Overall classification accuracy ranged from 77% for Kendall County to 98% for Macon County and averaged 91.8% over all the counties (Table 2). The Kappa value ranged from 0.58 for Kendall County to 0.98 for Peoria County and averaged 0.88 over all counties. The woods and water classes were the most accurately defined; errors of omission and commission for woods averaged 2.1 and 1.5%, respectively. The errors of omission and commission for water averaged 3.8 and 0.1%, respectively. Average error estimates for crop and grass were notably higher. The errors of omission and commission for crop averaged 3.7 and 17.5% respectively, whereas omission and commission errors for grass averaged 23.6 and 6.4%, respectively. Accuracy assessment data sampled for each county also were pooled for statewide estimates (Table 3). These estimates were very similar to the average county accuracy assessments. Overall statewide accuracy was 92% with a Kappa value of 0.88. Once again, woods and water were the most accurate with errors of omission and commission <3.0%. errors of omission and commission for crop were 3.5 and 16.3%, respectively. Errors of omission and commission for grass were 23.4 and 6.0%, respectively.

Our statewide classification compared well with other land use classifications using Landsat 5 TM data. Sader et al. (1991)

reported a 70% overall classification accuracy (93% for woods) for an area of Costa Rica. Airola and Vogel (1988) classified TM data for New Jersey and reported an overall accuracy of 91.8% (94.1% for woods and 82.4% for agriculture). Moore and Bauer (1990) reported an overall accuracy of 87% for their classification of TM data in northern Minnesota. Because our classification accuracy varied by county and by land use type, we have provided detailed estimates of accuracy for each county we reviewed (Appendix B).

LITERATURE CITED

Airola, T. M., and J. Vogel. 1988. Use of thematic mapper digital data for updating the New Jersey land cover component of the 1987 National Resources Inventory. J. Soil & Water Conserv. 43:425-428.

Anderson, J. R., E. E. Hardy, J. T. Roach and R. E. Witmer.

1976. A land use and land cover classification system for use with remote sensor data. U.S. Geologic Survey Prof.

Paper 964. 26pp.

- Congalton, R. G. 1991. A review of assessing the accuracy of classifications of remotely sensed data. Remote Sensing Environ. 37:35-46.
- Lillesand, T. M., and R. W. Kiefer 1987. Remote sensing and image interpretation. John Wiley & Sons, New York. 721pp.
- Miller, L. D., M. Unverferth, K. Ghormley, and M. P. Skrdla.

 1989. A guide to MIPS. MicroImages, Inc., Lincoln, Neb.

 97pp.
- Moore, M. M., and M. E. Bauer. 1990. Classification of forest vegetation in north-central Minnesota using Landsat multispectral scanner and thematic mapper data. For. Sci. 36:330-342.
- Sader, S. A., G. V. Powell and J. H. Rappole. 1991. Migratory bird habitat monitoring through remote sensing. Int. J. Remote Sensing. 12:363-372.
- Schwegman, J. E. 1973. Comprehensive plan for the Illinois nature preserves system part 2: the natural divisions of Illinois. Ill. Nature Preserves Comm., Springfield. 32pp.
- U.S. Dep. of Commerce 1994. 1992 census of agriculture, volume

 1 geographic area series, part 13, Illinois state and county
 data. U.S. Gov. Printing Office, Washington, DC. 541pp.

Table 1. Data on Landsat 5 TM scenes used to classify each Illinois county.

County	Scene1ª	Scene2	Source	Date1	Date2
Adams	P24R32		INHS	06/24/90	
Alexander	P23R34		MORRIS	10/12/92	
Bond	P23R33		INHS	05/26/88	
Boone	P23R31	P24R31	INHS	06/30/89	06/24/90
Brown	P24R32		INHS	06/24/90	
Bureau	P24R31		INHS	06/24/90	
Calhoun	P24R33		INHS	04/24/91	
Carroll	P24R31		INHS	06/24/90	
Cass	P24R32		INHS	06/24/90	
Champaign	P23R32		INHS	05/26/88	
Christian	P23R33	P23R32	INHS	05/26/88	05/26/88
Clark	P22R33		INHS	06/10/90	
Clay	P23R33	P22R33	INHS	05/26/88	06/10/90
Clinton	P23R33		INHS	05/26/88	
Coles	P22R33		INHS	06/10/90	
Cook	P23R31		MORRIS	09/10/92	
Crawford	P22R33		INHS	06/10/90	
Cumberland	P22R33		INHS	06/10/90	
De Kalb	P23R31		MORRIS	09/10/92	
De Witt	P23R32		INHS	05/26/88	
Douglas	P23R32		INHS	05/26/88	
Du Page	P23R31		MORRIS	09/10/92	
Edgar	P22R32	P22R33	INHS	06/10/90	06/10/90
Edwards	P22R33		INHS	06/10/90	
Effingham	P23R33		INHS	05/26/88	
Fayette	P23R33		INHS	05/26/88	
Ford	P23R32		INHS	05/26/88	
Franklin	P23R34		MORRIS	10/12/92	
Fulton	P24R32		INHS	06/24/90	
Gallatin	P22R34		INHS	06/10/90	
Greene	P24R33		INHS	04/24/91	
Grundy	P23R31		INHS	06/30/89	
Hamilton	P22R34	P22R33	INHS	06/10/90	06/10/90
Hancock	P24R32		INHS	06/24/90	
Hardin	P22R34		INHS	06/10/90	
Henderson	P24R32		INHS	06/24/90	

Henry Iroquois Jackson	P24R31 P23R32 P23R34	P23R31	MORRIS MORRIS	10/03/92 10/12/92 10/12/92	09/10/92
Jasper Jefferson Jersey	P22R33 P23R33 P24R33		INHS INHS INHS	06/10/90 05/26/88 04/24/91	
Jo Daviess Johnson	P24R31 P23R34		INHS MORRIS	06/24/90 10/12/92	

Table 1. Continued.

County	Scene1ª	Scene2	Source	Date1	Date2
Kane	P23R31		MORRIS	09/10/92	
Kankakee	P23R31		MORRIS	09/10/92	
Kendall	P23R31		MORRIS	09/10/92	
Knox	P24R32		INHS	06/24/90	
La Salle	P23R31		INHS	06/30/89	
Lake	P23R31		MORRIS	09/10/92	
Lawrence	P22R33		INHS	06/10/90	
Lee	P24R31		INHS	06/24/90	
Livingston	P23R32		INHS	05/26/88	
Logan	P23R32		INHS	05/26/88	
Macon	P23R32		INHS	05/26/88	
Macoupin	P23R33		INHS	05/26/88	
Madison	P23R33		INHS	05/26/88	
Marion	P23R33		INHS	05/26/88	
Marshall	P24R31		MORRIS	10/03/92	
Mason	P24R32		MORRIS	10/03/92	
Massac	P23R34		MORRIS	10/12/92	
McDonough	P24R32		INHS	06/24/90	
McHenry	P23R31		MORRIS	09/10/92	
McLean	P23R32		INHS	05/26/88	
Menard	P24R32		INHS	06/24/90	
Mercer	P24R31	P24R32	MORRIS	10/03/92	10/03/92
Monroe	P24R33		INHS	04/24/91	
Montgomery	P23R33		INHS	05/26/88	
Morgan	P24R32		INHS	06/24/90	
Moultrie	P23R32		INHS	05/26/88	
Ogle	P24R31		INHS	06/24/90	
Peoria	P24R32		INHS	06/24/90	

Perry	P23R34		MORRIS	10/12/92	
Piatt	P23R32		INHS	05/26/88	
Pike	P24R33	P24R32	INHS	04/24/91	06/24/90
Pope	P23R34		MORRIS	10/12/92	
Pulaski	P23R34		MORRIS	10/12/92	
Putnam	P24R31		MORRIS	10/03/92	
Randolph	P23R34		MORRIS	10/12/92	
Richland	P22R33		INHS	06/10/90	
Rock Island	P24R31		MORRIS	10/03/92	
Saline	P22R34		INHS	06/10/90	
Sangamon	P23R33		INHS	05/26/88	
Schuyler	P24R32		MORRIS	10/03/92	
Scott	P24R32		INHS	06/24/90	
Shelby	P23R33		INHS	05/26/88	
St. Clair	P23R34		MORRIS	10/12/92	
Stark	P24R31		INHS	06/24/90	
Stephenson	P24R31		INHS	06/24/90	

Table 1. Continued.

County	Scene1ª	Scene2	Source	Date1	Date2	
Tazewell	P24R32	P23R32	INHS	06/24/90	05/26/88	
Union	P23R34		MORRIS	10/12/92		
Vermilion	P22R32		INHS	06/10/90		
Wabash	P22R33		INHS	06/10/90		
Warren	P24R32		INHS	06/24/90		
Washington	P23R33		INHS	05/26/88		
Wayne	P22R33		INHS	06/10/90		
White	P22R33	P22R34	INHS	06/10/90	06/10/90	
Whiteside	P24R31		INHS	06/24/90		
Will	P23R31		MORRIS	09/10/92		
Williamson	P23R34		INHS	04/17/91		
Winnebago	P24R31	P23R31	INHS	06/24/90	06/30/89	
Woodford	P23R32		MORRIS	10/12/92		

aP=Path, R=Row.

Table 2. Summary of land use classification accuracy assessment and percent errors of omission (O) and commission (C) for Illinois counties classified from Landsat 5 TM satellite data.

	Crop		Gra	Grass		Woods		er	Accuracya	
County	0	С	0	С	0	С	0	С	Overall(%)	Kappa
Adams	16	28	25	18	0	0	7	0	86	0.81
Alexander	no	accuracy	-							
Bond	no	accuracy	-							
Boone	0	10	4	0	0	0	22	0	97	0.95
Brown	8	19	22	9	0	0	0	10	92	0.88
Bureau	3	21	31	0	0	6	0	0	90	0.85
Calhoun	no	accuracy	y asses	sment						
Carroll	6	6	5	10	6	0	0	0	95	0.93
Cass	3	16	26	5	0	0	0	0	92	0.89
Champaign	0	11	18	0	8	0	0	0	95	0.92
Christian	7	7	15	15	0	0	0	0	93	0.90
Clark	0	33	44	6	4	4	11	0	87	0.81
Clay	4	25	40	13	0	0	14	0	87	0.81
Clinton	8	23	35	10	0	7	0	0	87	0.82
Coles	0	21	38	0	0	0	0	0	91	0.87
Cook	no	accuracy	, asses	sment						
Crawford	14	14	30	20	0	5	0	0	90	0.86
Cumberland	0	7	12	0	0	0	0	0	97	0.96
De Kalb	0	6	17	0	_	_	0	0	96	0.92
De Witt	0	24	24	6	38	0	0	0	87	0.80
Douglas	30	27	56	6	0	0	0	0	85	0.76
Du Page	no	accuracy		sment						
Edgar	3	3	4	9	12	0	0	0	96	0.93

Edwards	4	17	14	5	0	0	8	0	94	0.91
Effingham	4	30	35	12	7	0	12	0	86	0.80
Fayette	0	23	25	0	0	0	0	0	93	0.90
Ford	0	5	10	0	0	0	0	0	97	0.96
Franklin	7	11	12	12	5	0	0	0	93	0.90
Fulton	0	38	45	15	10	5	17	0	83	0.76
Gallatin	0	19	14	0	0	0	0	0	95	0.93
T 1 1 0 0										

Table 2. Continued.

	Crop		Gr	Grass		Woods		iter	Accuracya	
County	0	С	0	С	0	С	0	С	Overall(%)	 Kappa
Greene	no a	ccurac	y asses	sment						
Grundy	7	3	10	20	0	0	0	0	95	0.92
Hamilton	11	0	0	8	0	0	0	0	97	0.96
Hancock	0	6	4	13	0	0	75	0	92	0.88
Hardin	0	19	14	0	0	0	0	0	95	0.93
Henderson	0	38	53	0	0	6	33	0	84	0.76
Henry	0	5	12	0	0	0	0	0	97	0.96
Iroquois	3	6	18	0	0	11	0	0	95	0.91
Jackson	0	18	19	0	0	0	0	0	95	0.93
Jasper	4	25	40	13	0	0	14	0	87	0.81
Jefferson	4	39	43	5	0	0	0	0	87	0.82
Jersey	0	10	18	0	0	0	0	0	96	0.94
Jo Daviess	6	6	5	10	6	0	0	0	95	0.93
Johnson	12	25	9	4	0	0	0	0	95	0.93
Kane	0	12	20	0	0	0	0	0	95	0.93
Kankakee	0	12	25	0	_	_	0	0	93	0.86
Kendall	3	39	86	0	11	11	0	0	77	0.58
Knox	no a	ccurac	y asses	sment						

Lake	no a	ccuracy	asses	sment						
La Salle	0	31	36	0	0	0	0	0	88	0.83
Lawrence	0	18	17	6	5	10	50	0	89	0.85
Lee	0	12	27	0	0	0	0	0	95	0.93
Livingston	0	13	27	0	0	0	0	0	93	0.88
Logan	0	9	12	0	0	0	12	0	96	0.94
Macon	0	4	10	0	0	0	0	0	98	0.97
Macoupin	4	12	15	5	0	0	0	0	94	0.92
Madison	9	9	19	25	7	0	0	0	91	0.87
Marion	4	27	33	6	0	0	0	0	90	0.87
Marshall	0	6	20	0	0	0	0	0	96	0.94
Mason	3	45	74	5	0	6	8	0	80	0.71
Massac	0	17	14	0	0	0	0	0	96	0.94
McDonough	0	38	45	15	10	5	17	0	83	0.76
Table 2. Cor	ntinued	١.								

County	Crop		Grass		Woods		Water		Accuracya	
	0	C	0	С	0	С	0	С	Overall(%)	 Карра
McHenry	no a	accuracy	asses	ssment						
McLean		accuracy								
Menard		accuracy								
Mercer		accuracy								
Monroe	5	33	37	5	0	0	0	0	88	0.84
Montgomery	6	24	31	15	6	0	0	0	89	0.85
Morgan	4	14	24	6	0	0	0	0	93	0.91
Moultrie	no a	accuracy	asses	sment						
Ogle	0	17	17	4	7	13	22	0	90	0.86
Peoria	0	4	9	0	0	0	0	0	98	0.98
Perry	4	12	17	6	0	0	0	0	95	0.93

Piatt	0	7	9	0	0	0	0	0	97	0.96
Pike	4	8	10	5	0	0	0	0	96	0.95
Pope	12	25	9	4	0	0	0	0	95	0.93
Pulaski	4	4	6	6	6	6	0	0	96	0.94
Putnam	0	6	20	0	0	0	0	0	96	0.94
Randolph	4	8	11	11	4	0	0	0	95	0.93
Richland	4	20	31	6	0	0	0	0	92	0.89
Rock Island	3	10	14	5	0	0	0	0	95	0.92
Saline	0	50	52	0	0	0	0	7	85	0.80
Sangamon	2	2	6	18	11	0	0	0	96	0.93
Schuyler	18	7	15	38	0	0	0	0	90	0.87
Scott	8	4	5	10	0	0	0	0	96	0.94
Shelby	0	25	32	10	10	0	0	0	88	0.84
Stark	no a	ccurac	y asses	sment						
St. Clair	0	12	19	0	0	0	0	0	95	0.93
Stephenson	3	31	40	0	0	4	0	0	88	0.82
Tazewell	0	9	15	0	0	0	0	0	96	0.93
Union	16	10	15	23	0	0	0	0	92	0.89
Vermilion	0	6	12	0	0	0	0	0	97	0.95
Wabash	0	19	31	0	0	0	0	0	93	0.90
Warren	0	37	38	0	7	0	0	0	87	0.81
Table 2. Cor	ntinued	•								

County	Crop		Grass		Woods		Water		Accuracya	
	0	С	0	С	0	С	0	C	Overall(%)	 Карра
Washington	4	28	30	5	0	0	8	0	90	0.86
Wayne	14	46	57	10	0	15	0	0	81	0.75
White	12	29	42	17	0	0	0	0	85	0.79
Whiteside	4	19	26	10	5	0	0	0	89	0.84

Will	12	0	0	30	0	0	0	0	94	0.91
Williamson	14	27	24	12	0	0	0	0	89	0.85
Winnebago	3	31	40	0	0	4	0	0	88	0.82
Woodford	2	2	8	0	0	11	0	0	97	0.95

aThe accuracy assessments are as follows:

Overall accuracy = the total number of fields correctly classified, divided by the total number of fields sampled, multiplied by 100.

Table 3. Accuracy assessment and error estimate samples pooled for an overall statewide estimate of accuracy.

Land Use		Cla	ssified M	ар		Err	Accuracy	
Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	(%)
Crop	2134	65	12	0	2211	3.5	16.3	96.5
Grass	341	1144	7	1	1493	23.4	6.0	76.6
Woods	9	17	1331	0	1357	1.9	1.7	98.1
Water	10	7	4	674	695	3.0	0.1	97.0
Sum	2494	1233	1354	675	5756			
Overal	l Accuracy	7: 92%	Kappa	0.88				

^aCalculated according to Congalton 1991.

^bGround truth data from Illinois Department of Natural Resources quail and pheasant call count routes.

Appendix A. Counties with greater than 48 ha of orchards were digitized using USGS 1:24000-scale topographical quadrangle maps supplied by Morris Library. These 23 counties represent 73% of land in orchards. Cumberland County quad maps were unavailable (51.8 ha). The orchard vectors were merged into a single vector and then converted to a raster for each county.

County	Quad	Quad year	Area haª
Union			730.5
	Anna	1978	
	Cobden	1990	
	Makanda	1990	
	Mill Creek	1990	
Jackson			552.8
	Carbondale	1990	
	Cobden	1990	
	Elkville	1978	
	Makanda	1990	
	Murphysboro	1978	
	Pomona	1990	
	Willisville	1968	
Calhoun			448.4
	Brussels	1974	
	Foley	1975	
	Hamburg	1978	
	Kampsville	1980	
St. Clair			241.6
	Cahokia	1974	
	Collinsville	1974	
	Columbia	1979	
	Freeburg	1974	
	French Village	1982	
	Mascoutah	1990	
Jersey			159.5
	Alton	1974	
	Brussels	1974	
	Elsah	1974	
	Grafton	1974	
	Hardin	1978	
	Nutwood	1975	

	Otterville	1983	
Pike			135.6
	Barry	1978	
	Pearl East	1980	
	Pearl West	1980	
	Summer Hill	1981	

County	Quad	Quad year	Area haª	
Adams			132.3	
	Lima	1981		
	Long Island	1981		
	Loraine	1981		
	Quincy East	1971		
	Quincy West	1971		
	Richfield	1981		
	Tioga	1981		
Macoupin	-		97.9	
-	Carlinville West	1979		
	Plainview	1974		
McHenry			90.7	
-	Hebron	1972		
	Fox Lake	1993		
	Marengo North	1970		
Will	5		85.0	
	Joliet	1973		
Jefferson			83.8	
	Harmony	1965		
	Irvington	1974		
	Kell	1978		
	Mount Vernon	1978		
	Walnut Hill	1974		
Marion			81.7	
	Centralia East	1970		
	Harmony	1965		
	Irvington	1974		
	Iuka	1965		

	Kell	1978	
	Salem North	1978	
	Salem South	1978	
Lake			70.4
	Antioch	1993	
	Grayslake	1993	
	Libertyville	1993	
Madison			63.5
	Bethalto	1974	
	Collinsville	1974	
	Elsah	1974	
	Worden	1982	
Randolph			63.5
	Coulterville	1982	
	Kaskaskia	1982	
	Chester	1970	

County	Quad	Quad year	Area haª	
I-To			CO 3	
Wayne	Bluford	1973	60.3	
Rock Island	Blulord	1973	58.7	
NOCK ISLAND	Fort Byron	1991	30.7	
	Milan	1992		
	Silvis	1992		
Marshall	SIIVIS	1991	56.7	
narsnarr	Castleton	1983	30.7	
	La Prairie Center	1983		
Franklin	na rranne center	1905	55.8	
TTUINTIII	Christopher	1968	33.0	
	Harco	1963		
	Johnston City	1976		
	Macedonia	1974		
	Pittsburg	1963		
	Rend Lake Dam	1975		
	Sesser	1975		
	Thompsonville	1976		
	West Frankfort	1978		

Winnebago			55.0
	Ridott	1971	
	South Beloit	1993	
Hancock			50.6
	Augusta	1981	
	Nauvoo	1975	
	Niota	1964	
	Tioga	1981	
Johnson			49.0
	Creal Springs	1976	
	Cypress	1966	
	Goreville	1990	
	Stonefort	1961	
Massac			48.6
	Metropolis	1990	

^aU.S. Department of Commerce. 1994. 1992 Census of agriculture. Vol 1 Part 13. Illinois state and county data.

Appendix B. Accuracy assessment and error estimates for Illinois counties classified from Landsat 5 TM satellite data.

Tanal IIaa		Cl	assified	d Map		Er	ror(%)	Accuracy (%)
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	
Adams Cou	ınty							
Crop	21	4	0	0	25	16.0	28.0	84.
Grass	7	21	0	0	28	25.0	17.9	75.
Woods	0	0	21	0	21	0.0	0.0	100.
Water	0	1	0	13	14	7.1	0.0	92.
Sum	28	26	21	13	88			
(Overall	Accurac	ey: 86%	Kappa	a: 0.81			
Alexandeı	Count	y - no a	accuracy	assessme	ent			
	 >++ n			 ssment				
Bona Cour	icy – II	o accura	cy asse.					
Bond Cour Boone Cou			·					
 Boone Cou			0	0	31	0.0	9.7	100.
Boone Cou	inty				31 24	0.0	9.7	
Boone Cou	31	0	0	0	-			95.
Boone Cou Crop Grass Woods	31 1	0 23	0	0	24	4.2	0.0	95. 100.
Boone Cou Crop Grass Woods Water	31 1 0	0 23 0	0 0 26	0 0 0	24 26	4.2 0.0	0.0	95. 100.
Boone Cou Crop Grass Woods Water Sum	31 1 0 2	0 23 0 0	0 0 0 26 0	0 0 0 7 7	24 26 9	4.2 0.0 22.2	0.0	100. 95. 100. 77.
Boone Cou Crop Grass Woods Water Sum	31 1 0 2 34	0 23 0 0	0 0 26 0	0 0 0 7 7	24 26 9	4.2 0.0 22.2	0.0	95. 100.
Boone Cou Crop Grass Woods Water Sum	31 1 0 2 34	0 23 0 0	0 0 26 0	0 0 0 7 7	24 26 9	4.2 0.0 22.2	0.0	95. 100. 77.
Boone Cou Crop Grass Woods Water Sum Grass Crop	31 1 0 2 34 Overall	0 23 0 0 23 Accurac	0 0 26 0 26 29: 97%	0 0 0 7 7 Kappa	24 26 9 90 a: 0.95	4.2 0.0 22.2	0.0 0.0 0.0	95. 100. 77.
Boone Cou Crop Grass Woods Water Sum Grass Brown Cou	31 1 0 2 34 Overall	0 23 0 0 23 Accurac	0 0 26 0 26 29: 97%	0 0 0 7 7 Kappa	24 26 9 90 a: 0.95	4.2 0.0 22.2	0.0 0.0 0.0	95. 100. 77.
Boone Cou Crop Grass Woods Water Sum Grass Brown Cou Crop Grass Woods	31 1 0 2 34 Overall inty 24 5	0 23 0 0 23 Accurace	0 0 26 0 26 29: 97%	0 0 0 7 7 7 Kappa 	24 26 9 90 a: 0.95 	4.2 0.0 22.2	0.0 0.0 0.0	95. 100. 77. 92. 78. 100.
Boone Cou Crop Grass Woods Water Sum	31 1 0 2 34 Overall inty 24 5	0 23 0 0 23 Accurace	0 0 26 0 26 27: 97%	0 0 0 7 7 Kappa 	24 26 9 90 a: 0.95 	4.2 0.0 22.2	0.0 0.0 0.0	95. 100.

Appendix B. Continued.

Tand Has		Class	sified Ma	ap		E:	rror(%)	\\ aauraau
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
Bureau Co	ounty							
Crop	37	0	1	0	38	2.6	21.1	97.4
Grass	8	18	0	0	26	30.8	0.0	69.2
Woods	0	0	16	0	16	0.0	6.3	100.0
Water	0	0	0	9	9	0.0	0.0	100.0
Sum	45	18	17	9	89			
(Overall	Accurac	cy: 90%	Kappa	a: 0.85			
Calhoun (County	- no acc	curacy as	ssessment	 t			
Carroll (County							
Crop	16	1	0	0	17	5.9	5.9	94.1
Grass	1	18	0	0	19	5.3	10.5	94.
Woods	0	1	16	0	17	5.9	0.0	94.1
Water	0	0	0	6	6	0.0	0.0	100.0
Sum	17	20	16	6	59			
(Overall	Accurac	cy: 95%	Kappa	a: 0.93			
Cass Cour	 nty		. – – – .					
Crop	30	1	0	0	31	3.2	16.1	96.8
Grass	5	14	0	0	19	26.3	5.3	73.
Woods	0	0	17	0	17	0.0	0.0	100.0
Water	0	0	0	11	11	0.0	0.0	100.0
Sum	35	15	17	11	78			
(Overall	Accurac	cy: 92%	Kappa	a: 0.89			
Champaign	n Count	У						
Crop	36	0	0	0	36	0.0	11.1	100.0
Grass	3	14	0	0	17	17.7	0.0	82.4
Woods	1	0	11	0	12	8.3	0.0	91.
Water	0	0	0	10	10	0.0	0.0	100.0
Sum	40	14	11	10	75			

Overall Accuracy: 95% Kappa: 0.92

Appendix B. Continued.

Tand Had		Class	sified Ma	ap		E:	rror(%)	Accuracy (%)
Land Use	Crop	Grass	Woods	Water	Sum	Omission	Commission	
Christia	n Count	.y						
Crop	25	2	0	0	27	7.4	7.4	92.6
Grass	2	11	0	0	13	15.4	15.4	84.6
Woods	0	0	14	0	14	0.0	0.0	100.0
Water	0	0	0	6	6	0.0	0.0	100.0
Sum	27	13	14	6	60			
	Overall	Accurac	cy: 93%	Kappa	a: 0.90			
Clark Co	unty							
Crop	24	0	0	0	24	0.0	33.3	100.0
Grass	7	10	1	0	18	44.4	5.6	55.6
Woods	0	1	23	0	24	4.2	4.2	95.8
Water	1	0	0	8	9	11.1	0.0	88.9
Sum	32	11	24	8	75			
	Overall	Accurac	cy: 87%	Kappa	a: 0.81			
Clay Cou	 inty	. – – –	. – – – .					
Crop	23	1	0	0	24	4.2	25.0	95.8
Grass	6	9	0	0	15	40.0	13.3	60.0
Woods	0	0	15	0	15	0.0	0.0	100.0
Water	0	1	0	6	7	14.3	0.0	85.7
Sum	29	11	15	6	61			
	Overall	Accurac	cy: 87%	Kappa	a: 0.81			
Clinton	County	. – – –						
Crop	24	2	0	0	26	7.7	23.1	92.3
Grass	6	13	1	0	20	35.0	10.0	65.0
Woods	0	0	14	0	14	0.0	7.1	100.0
Water	0	0	0	9	9	0.0	0.0	100.0
Sum	30	15	15	9	69			
	Overall	Accurac	cy: 87%	Kappa	a: 0.82			

Appendix B. Continued.

Tand Has		Clas	sified M	ap		E	rror(%)	Accuracy
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	(%)
Coles Cou	ınty							
Crop	28	0	0	0	28	0.0	21.4	100.0
Grass	6	10	0	0	16	37.5	0.0	62.5
Woods	0	0	13	0	13	0.0	0.0	100.0
Water	0	0	0	11	11	0.0	0.0	100.0
Sum	34	10	13	11	68			
	verall	Accura	cy: 91%	Kappa	a: 0.87			
Cook Cour	nty - n	o accura	acy asse	ssment				
Crawford	County							
Crop	19	2	1	0	22	13.6	13.6	86.4
Grass	3	7	0	0	10	30.0	20.0	70.0
Woods	0	0	21	0	21	0.0	4.8	100.0
Water	0	0	0	8	8	0.0	0.0	100.0
Sum	22	9	22	8	61			
C)verall	Accura	cy: 90%	Kappa	a: 0.86			
Cumberlar	d Coun	ty						
Crop	29	0	0	0	29	0.0	6.9	100.0
Grass	2	14	0	0	16	12.5	0.0	87.5
Woods	0	0	22	0	22	0.0	0.0	100.0
Water	0	0	0	6	6	0.0	0.0	100.0
Sum	31	14	22	6	73			
C	verall	Accura	cy: 97%	Kappa	a: 0.96			
De Kalb C	County							
Crop	32	0	0	0	32	0.0	6.3	100.0
Grass	2	10	0	0	12	16.7	0.0	83.3
Woods	0	0	0	0	0	_	_	-
Water	0	0	0	6	6	0.0	0.0	100.0

Overall Accuracy: 96% Kappa: 0.92

Appendix B. Continued.

		Classi	fied M	ap		E:	rror(%)	7
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
De Witt	County							
Crop	33	0	0	0	33	0.0	24.2	100.0
Grass	4	13	0	0	17	23.5	5.9	76.5
Woods	4	1	8	0	13	38.5	0.0	61.5
Water	0	0	0	7	7	0.0	0.0	100.0
Sum	41	14	8	7	70			
	Overall	Accuracy	∵: 87%	Kappa	a: 0.80			
Douglas	 County							
Crop	32	1	0	0	33	3.0	27.3	97.0
Grass	9	7	0	0	16	56.3	6.3	43.8
Woods	0	0	12	0	12	0.0	0.0	100.0
Water	0	0	0	5	5	0.0	0.0	100.0
Sum	41	8	12	5	66			
	Overall	Accuracy	: 85%	Kappa	a: 0.76			
Du Page	County	- no accu	racy a	ssessment	 :			
Edgar Co	unty							
Crop	29	1	0	0	30	3.3	3.3	96.7
Grass	1	21	0	0	22	4.5	9.1	95.5
Woods	0	1	7	0	8	12.5	0.0	87.5
Water	0	0	0	7	7	0.0	0.0	100.0
Sum	30	23	7	7	67			
	Overall	Accuracy	: 96%	Kappa	a: 0.93			
Edwards	County							
	0.0	1	0	0	24	4.2	16.7	95.8
Crop	23							
Crop Grass	23 3		0	0				
Crop Grass Woods	23 3 0	18			21 22	14.3	4.8	85.7 100.0

Sum 27 19 22 11 79

Overall Accuracy: 94% Kappa: 0.91

Tand Had	2	Clas	sified Ma	ap		E	rror(%)	7 a a u ma a u
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
Effingha	am Count	У						
Crop	22	1	0	0	23	4.3	30.4	95.7
Grass	6	11	0	0	17	35.3	11.8	64.7
Woods	1	0	14	0	15	6.7	0.0	93.3
Water	0	1	0	7	8	12.5	0.0	87.5
Sum	29	13	14	7	63			
	Overall	Accura	cy: 86%	Kappa	a: 0.80			
Fayette	County							
Crop	22	0	0	0	22	0.0	22.7	100.0
Grass	5	15	0	0	20	25.0	0.0	75.0
Woods	0	0	19	0	19	0.0	0.0	100.0
Water	0	0	0	7	7	0.0	0.0	100.0
Sum	27	15	19	7	68			
	Overall	Accura	cy: 93%	Kappa	a: 0.90			
Ford Co	unty							
Crop	40	0	0	0	40	0.0	5.0	100.0
Grass	2	19	0	0	21	9.5	0.0	90.5
Woods	0	0	9	0	9	0.0	0.0	100.0
Water	0	0	0	7	7	0.0	0.0	100.0
Sum	42	19	9	7	77			
	Overall	Accura	cy: 97%	Kappa	a: 0.96			
Frankli	County							
Crop	25	2	0	0	27	7.4	11.1	92.6
Grass	2	15	0	0	17	11.8	11.8	88.2
Woods	1	0	18	0	19	5.3	0.0	94.7
Water	0	0	0	7	7	0.0	0.0	100.0
·va cci	O	O	O	,	,	0.0	0.0	100.0

Sum 28 17 18 7 70

Overall Accuracy: 93% Kappa: 0.90

Appendix B. Continued.

Land Us	0	Classi	ified Ma	ap		E:	rror(%)	Accuracy
Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	(%)
Fulton	County							
Crop	24	0	0	0	24	0.0	37.5	100.0
Grass	9	11	0	0	20	45.0	15.0	55.0
Woods	0	2	17	0	19	10.5	5.3	89.5
Water	0	1	1	10	12	16.7	0.0	83.3
Sum	33	14	18	10	75			
	Overall	Accuracy	y: 83%	Kappa	a: 0.76			
Gallati	n County							
Crop	16	0	0	0	16	0.0	18.8	100.0
Grass	3	19	0	0	22	13.6	0.0	86.4
Woods	0	0	18	0	18	0.0	0.0	100.0
Water	0	0	0	7	7	0.0	0.0	100.0
Sum	19	19	18	7	63			
	Overall	Accuracy	7: 95%	Kappa	a: 0.93			
Greene	County -	no accur	cacy as	sessment				
Grundy	County							
Crop	28	2	0	0	30	6.7	3.3	93.3
Grass	1	9	0	0	10	10.0	20.0	90.0
Woods	0	0	14	0	14	0.0	0.0	100.0
Water	0	0	0	6	6	0.0	0.0	100.0
Sum	29	11	14	6	60			
	Overall	Accuracy	7: 95%	Карра	a: 0.92			
Hamilto	n County	. – – –						
Crop	16	2	0	0	18	11.1	0.0	88.9
Grass	0	25	0	0	25	0.0	8.0	100.0
Woods	0	0	20	0	20	0.0	0.0	100.0
Water	0	0	0	13	13	0.0	0.0	100.0
Sum	16	27	20	13	76			

Overall Accuracy: 97% Kappa: 0.96

Appendix B. Continued.

T 1 TT		Classi	fied Ma	ap		E	rror(%)	Accuracy
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
Hancock	County							
Crop	17	0	0	0	17	0.0	5.9	100.0
Grass	1	22	0	0	23	4.3	13.0	95.7
Woods	0	0	9	0	9	0.0	0.0	100.0
Water	0	3	0	1	4	75.0	0.0	25.0
Sum	18	25	9	1	53			
	Overall	Accuracy	7: 92%	Kappa	a: 0.88			
Hardin (County							
Crop	16	0	0	0	16	0.0	18.8	100.0
Grass	3	19	0	0	22	13.6	0.0	86.4
Woods	0	0	18	0	18	0.0	0.0	100.0
Water	0	0	0	7	7	0.0	0.0	100.0
Sum	19	19	18	7	63			
	Overall	Accuracy	7: 95%	Kappa	a: 0.93			
Henderso	on Count	У						
Crop	21	0	0	0	21	0.0	38.1	100.0
Grass	8	7	0	0	15	53.3	0.0	46.7
Woods	0	0	17	0	17	0.0	5.9	100.0
Water	0	0	1	2	3	33.3	0.0	66.7
Sum	29	7	18	2	56			
	Overall	Accuracy	7: 84%	Kappa	a: 0.76			
 Henry Co								
e.i.ry CC	Juli Cy							
Crop	38	0	0	0	38	0.0	5.3	100.0
Grass	2	15	0	0	17	11.8	0.0	88.2
Woods	0	0	17	0	17	0.0	0.0	100.0
Water	0	0	0	5	5	0.0	0.0	100.0

Sum 40 15 17 5 77

Overall Accuracy: 97% Kappa: 0.96

Tomal II.a	_	Classi	fied M	ap		E	rror(%)	7
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
Iroquois	S County							
Crop	31	0	1	0	32	3.1	6.3	96.9
Grass	2	9	0	0	11	18.2	0.0	81.8
Woods	0	0	9	0	9	0.0	11.1	100.0
Water	0	0	0	4	4	0.0	0.0	100.0
Sum	33	9	10	4	56			
	Overall	Accuracy	: 95%	Kappa	: 0.91			
Jackson	County							
Crop	22	0	0	0	22	0.0	18.2	100.0
Grass	4	17	0	0	21	19.0	0.0	81.0
Woods	0	0	21	0	21	0.0	0.0	100.0
Water	0	0	0	11	11	0.0	0.0	100.0
Sum	26	17	21	11	75			
	Overall	Accuracy	: 95%	Kappa	: 0.93			
Jasper (County							
Crop	23	1	0	0	24	4.2	25.0	95.8
Grass	6	9	0	0	15	40.0	13.3	60.0
Woods	0	0	15	0	15	0.0	0.0	100.0
Water	0	1	0	6	7	14.3	0.0	85.7
Sum	29	11	15	6	61			
	Overall	Accuracy	: 87%	Kappa	: 0.81			
Jefferso	on Count	У						
Crop	22	1	0	0	23	4.3	39.1	95.7
Grass	9	12	0	0	21	42.9	4.8	57.1

Woods Water	0	0	20	0 11	20 11	0.0	0.0	100.0
Sum	31	13	20	11	75			
	Overall	Accuracy:	87%	Kappa:	0.82			

T 1 TT		Classi	fied M	ap		E	rror(%)	3
Land Use Class ^b		Grass	Woods	Water	Sum	Omission	Commission	Accuracy
Jersey C	ounty							
Crop	31	0	0	0	31	0.0	9.7	100.
Grass	3	14	0	0	17	17.6	0.0	82.
Woods	0	0	22	0	22	0.0	0.0	100.
Water	0	0	0	8	8	0.0	0.0	100.
Sum	34	14	22	8	78			
	Overall	Accuracy	: 96%	Kappa:	0.94			
 Jo Davie	ss Coun	 ty						
Crop	16	1	0	0	17	5.9	5.9	94.
Grass	1	18	0	0	19	5.3	10.5	94.
Woods	0	1	16	0	17	5.9	0.0	94.
Water	0	0	0	6	6	0.0	0.0	100.
Sum	17	20	16	6	59			
	Overall	Accuracy	: 95%	Kappa:	0.93			
Johnson	 County							
Crop	7	1	0	0	8	12.5	25.0	87.
Grass	2	21	0	0	23	8.7	4.3	91.
Woods	0	0	25	0	25	0.0	0.0	100.
Water	0	0	0	8	8	0.0	0.0	100.
Sum	9	22	25	8	64			
	Overall	Accuracy	: 95%	Kappa:	0.93			
 Kane Cou	 nty							
Crop	26	0	0	0	26	0.0	11.5	100.

Grass	3	12	0	0	15	20.0	0.0	80.0
Woods	0	0	13	0	13	0.0	0.0	100.0
Water	0	0	0	6	6	0.0	0.0	100.0
Sum	29	12	13	6	60			
	Overall	Accuracy:	95%	Kappa:	0.93			
Sum				-				

		Class	sified M	ap		E	rror(%)	
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy
Kankakee	County	•						-
Crop	34	0	0	0	34	0.0	11.8	100.0
Grass	4	12	0	0	16	25.0	0.0	75.0
Woods	0	0	0	0	0	_	_	-
Water	0	0	0	6	6	0.0	0.0	100.0
Sum	38	12	0	6	56			
	Overall	Accurac	cy: 93%	Kappa	a: 0.86	5		
 Kendall	 County							
Crop	32	0	1	0	33	3.0	39.4	97.0
Grass	12	2	0	0	14	85.7	0.0	14.3
Woods	1	0	8	0	9	11.1	11.1	88.9
Water	0	0	0	4	4	0.0	0.0	100.0
Sum	45	2	9	4	60			
	Overall	Accurac	cy: 77%	Kappa	a: 0.58	3		
Knox Cou	 nty - n	o accura	acy asse	ssment				
 Lake Cou	 nty - n	o accura	 acy asse	ssment				
 La Salle	 County	· •						
Cron	26	0	0	0	26	0 0	30 0	100.0
Crop Grass	26 8	14	0	0	26 22	0.0 36.4	30.8 0.0	63.6
	0	0	13	0	13	0.0	0.0	
Woods	0	0	13	7	13 7	0.0		100.0
Water	U	U	U	/	/	0.0	0.0	100.0
Sum	34	14	13	7	68			

Overall Accuracy: 88% Kappa: 0.83

Appendix B. Continued.

Tand Haa		Class	sified Ma	ap		E	rror(%)	A course
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
Lawrence	County	,						
Crop	22	0	0	0	22	0.0	18.2	100.0
Grass	2	15	1	0	18	16.7	5.6	83.3
Woods	0	1	19	0	20	5.0	10.0	95.0
Water	2	0	1	3	6	50.0	0.0	50.0
Sum	26	16	21	3	66			
	Overall	Accurac	cy: 89%	Карра	a: 0.85			
Lee Coun	ty							
Crop	34	0	0	0	34	0.0	11.8	100.0
Grass	4	11	0	0	15	26.7	0.0	73.3
Woods	0	0	22	0	22	0.0	0.0	100.0
Water	0	0	0	12	12	0.0	0.0	100.0
Sum	38	11	22	12	83			
	Overall	Accurac	cy: 95%	Kappa	a: 0.93			
 Livingst	on Coun	ty						
Crop	31	0	0	0	31	0.0	12.9	100.0
Grass	4	11	0	0	15	26.7	0.0	73.3
Woods	0	0	7	0	7	0.0	0.0	100.0
Water	0	0	0	3	3	0.0	0.0	100.0
Sum	35	11	7	3	56			
	Overall	Accurac	cy: 93%	Kappa	a: 0.88			
Logan Co	 unty							
Crop	35	0	0	0	35	0.0	8.6	100.0
Grass	2	14	0	0	16	12.5	0.0	87.5
Woods	0	0	15	0	15	0.0	0.0	100.0
Water	1	0	0	7	8	12.5	0.0	87.5
Sum	38	14	15	7	74			
	Overall	Accurac	cy: 96%	Kappa	a: 0.94			

Appendix B. Continued.

Tanal IIa		Classi	fied M	ap		Omission Commission 0.0 3.6 1 10.0 0.0 0.0 0.0 1 0.0 0.0 1 0.0 0.0 1 0.0 0.0 1 0.0 0.0 1 0.0 0.0 1 0.0 0.0 1 0.0 0.0 1		
Land Use	e ——— Crop	Grass 1	Woods	Water	Sum	Omission	Commission	Accuracy (%)
Macon Co	ounty							
Crop	28	0	0	0	28	0.0	3.6	100.0
Grass	1	9	0	0	10	10.0	0.0	90.0
Woods	0	0	6	0	6	0.0	0.0	100.0
Water	0	0	0	6	6	0.0	0.0	100.0
Sum	29	9	6	6	50			
	Overall	Accuracy	: 98%	Kappa	a: 0.97			
Macoupi	n County							
Crop	23	1	0	0	24	4.2	12.5	95.8
Grass	3	17	0	0	20	15.0		85.0
Woods	0	0	19	0	19	0.0	0.0	100.0
Water	0	0	0	7	7	0.0	0.0	100.0
Sum	26	18	19	7	70			
	Overall	Accuracy	: 94%	Kappa	a: 0.92			
 Madison	 County							
Crop	29	3	0	0	32	9.4	9.4	90.6
Grass	3	13	0	0	16			81.3
Woods	0	1	14	0	15			93.3
Water	0	0	0	12	12			100.0
Sum	32	17	14	12	75			
	Overall	Accuracy	: 91%	Kappa	a: 0.87			
Marion	 County							
Crop	21	1	0	0	22	4.5	27.3	95.5
Grass	6	12	0	0	18	33.3	5.6	66.7
Woods	0	0	20	0	20	0.0	0.0	100.0
Water	0	0	0	12	12	0.0	0.0	100.0
Sum	27	13	20	12	72			
	Overall	Accuracy	: 90%	Kappa	a: 0.87			

Appendix B. Continued.

Tand Ha	_	Classified Map				E	rror(%)	7 ~ ~
Land Use Class ^b	e ————————————————————————————————————	Grass V	Voods	Water	Sum	Omission	Commission	Accuracy (%)
Marshal:	l County							
Crop	33	0	0	0	33	0.0	6.1	100.0
Grass	2	8	0	0	10	20.0	0.0	80.0
Woods	0	0	8	0	8	0.0	0.0	100.0
Water	0	0	0	5	5	0.0	0.0	100.0
Sum	35	8	8	5	56			
	Overall	Accuracy:	: 96%	Kappa	a: 0.94			
Mason Co	ounty							
Crop	30	1	0	0	31	3.2	45.2	96.8
Grass	14	5	0	0	19	73.7	5.3	26.3
Woods	0	0	16	0	16	0.0	6.3	100.0
Water	0	0	1	12	13	7.7	0.0	92.3
Sum	44	6	17	12	79			
	Overall	Accuracy:	: 80%	Kappa	a: 0.71			
 Massac (County		·					
Crop	24	0	0	0	24	0.0	16.7	100.0
Grass	4	24	0	0	28	14.3	0.0	85.7
Woods	0	0	30	0	30	0.0	0.0	100.0
Water	0	0	0	6	6	0.0	0.0	100.0
Sum	28	24	30	6	88			
	Overall	Accuracy:	: 96%	Kappa	a: 0.94			
McDonou	gh Count	 У						
Crop	24	0	0	0	24	0.0	37.5	100.0
Grass	9	11	0	0	20	45.0	15.0	55.0
Woods	0	2	17	0	19	10.5	5.3	89.5
Water	0	1	1	10	12	16.7	0.0	83.3
Sum	33	14	18	10	75			
	Overall	Accuracy:	: 83%	Kappa	a: 0.76			

Appendix B. Continued.

Tand Has	_	Classified Map				E:	rror(%)	7
Land Use Class ^b		Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
McHenry	County	- no aco	curacy as	ssessment	5			
McLean C	ounty -	no acci	racy as	sessment				
Menard C	ounty -	no acci	racy as	sessment				
Mercer C	ounty -	no acci	racy as	sessment				
Monroe C	ounty							
Crop	20	1	0	0	21	4.8	33.3	95.2
Grass	7	12	0	0	19	36.8	5.3	63.2
Woods	0	0	17	0	17	0.0	0.0	100.0
Water	0	0	0	12	12	0.0	0.0	100.0
Sum	27	13	17	12	69			
	Overall	Accurac	cy: 88%	Карра	a: 0.84			
Montgome	ry Coun	ty						
Crop	16	1	0	0	17	5.9	23.5	94.1
Grass	4	9	0	0	13	30.8	15.4	69.2
Woods	0	1	15	0	16	6.3	0.0	93.7
Water	0	0	0	9	9	0.0	0.0	100.0
Sum	20	11	15	9	55			
	Overall	Accurac	cy: 89%	Карра	a: 0.85)		
 Morgan C	 ounty							
Crop	27	1	0	0	28	3.6	14.3	96.4
Grass	4	13	0	0	17	23.5	5.9	76.5
Woods	0	0	20	0	20	0.0	0.0	100.0
Water	0	0	0	10	10	0.0	0.0	100.0
Sum	31	14	20	10	75			
	Overall	Accurac	cy: 93%	Kappa	a: 0.91			

Appendix B. Continued.

- 1		Class	sified Ma	ap		E		
Land Use	Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
Moultrie	County	- no ac	ccuracy a	assessmer	nt			
 Ogle Cou	 nty							
Crop	23	0	0	0	23	0.0	17.4	100.0
Grass	2	20	2	0	24	16.7	4.2	83.3
Woods	0	1	14	0	15	6.7	13.3	93.3
Water	2	0	0	7	9	22.2	0.0	77.8
Sum	27	21	16	7	71			
	Overall	Accurac	cy: 90%	Kappa	a: 0.86)		
Peoria C	ounty							
Crop	25	0	0	0	25	0.0	4.0	100.0
Grass	1	10	0	0	11	9.1	0.0	90.9
Woods	0	0	13	0	13	0.0	0.0	100.0
Water	0	0	0	11	11	0.0	0.0	100.0
Sum	26	10	13	11	60			
	Overall	Accurac	cy: 98%	Kappa	a: 0.98	}		
Perry Co	unty							
Crop	24	1	0	0	25	4.0	12.0	96.0
Grass	3	15	0	0	18	16.7	5.6	83.3
Woods	0	0	20	0	20	0.0	0.0	100.0
Water	0	0	0	16	16	0.0	0.0	100.0
Sum	27	16	20	16	79			
	Overall	Accurac	cy: 95%	Kappa	a: 0.93	3		
Piatt Co	unty							
Crop	29	0	0	0	29	0.0	6.9	100.0
Grass	2	21	0	0	23	8.7	0.0	91.3
Woods	0	0	10	0	10	0.0	0.0	100.0
Water	0	0	0	8	8	0.0	0.0	100.0

Overall Accuracy: 97% Kappa: 0.96

Appendix B. Continued.

		Classi	fied M	ap		E	_	
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
Pike Cou	ınty							
Crop	24	1	0	0	25	4.0	8.0	96.0
Grass	2	19	0	0	21	9.5	4.8	90.5
Woods	0	0	23	0	23	0.0	0.0	100.0
Water	0	0	0	8	8	0.0	0.0	100.0
Sum	26	20	23	8	77			
	Overall	Accuracy	: 96%	Kappa	a: 0.95			
Pope Cou	nty							
Crop	7	1	0	0	8	12.5	25.0	87.5
Grass	2	21	0	0	23	8.7	4.3	91.3
Woods	0	0	25	0	25	0.0	0.0	100.0
Water	0	0	0	8	8	0.0	0.0	100.0
Sum	9	22	25	8	64			
	Overall	Accuracy	: 95%	Kappa	a: 0.93			
Pulaski	County							
Crop	23	0	1	0	24	4.2	4.2	95.8
Grass	1	17	0	0	18	5.6	5.6	94.4
Woods	0	1	17	0	18	5.6	5.6	94.4
Water	0	0	0	11	11	0.0	0.0	100.0
Sum	24	18	18	11	71			
	Overall	Accuracy	: 96%	Kappa	a: 0.94			
Putnam (County							
Crop	33	0	0	0	33	0.0	6.1	100.0
Grass	2	8	0	0	10	20.0	0.0	80.0
Woods	0	0	8	0	8	0.0	0.0	100.0
Water	0	0	0	5	5	0.0	0.0	100.0
Sum	35	8	8	5	56			

Overall Accuracy: 96% Kappa: 0.94

Appendix B. Continued.

Land Use		Class	sified Ma	ap		E:	Accuracy	
Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	(%)
Randolph	County	,						
Crop	23	1	0	0	24	4.2	8.3	95.8
Grass	2	16	0	0	18	11.1	11.1	88.9
Woods	0	1	21	0	22	4.5	0.0	95.5
Water	0	0	0	11	11	0.0	0.0	100.0
Sum	25	18	21	11	75			
	Overall	Accurac	cy: 95%	Kappa	a: 0.93	3		
Richland	County	,						
Crop	24	1	0	0	25	4.0	20.0	96.0
Grass	5	11	0	0	16	31.2	6.3	68.8
Woods	0	0	22	0	22	0.0	0.0	100.0
Water	0	0	0	10	10	0.0	0.0	100.0
Sum	29	12	22	10	73			
	Overall	Accurac	cy: 92%	Kappa	a: 0.89)		
Rock Isl	and Cou	ınty						
Crop	30	1	0	0	31	3.2	9.7	96.8
Grass	3	18	0	0	21	14.3	4.8	85.7
Woods	0	0	16	0	16	0.0	0.0	100.0
Water	0	0	0	7	7	0.0	0.0	100.0
Sum	33	19	16	7	75			
	Overall	Accurac	cy: 95%	Kappa	a: 0.92	2.		
Saline C	ounty							
Crop	20	0	0	0	20	0.0	50.0	100.0
Grass	10	10	0	1	21	52.4	0.0	47.6
Woods	0	0	19	0	19	0.0	0.0	100.0
Water	0	0	0	15	15	0.0	6.7	100.0
Sum	30	10	19	16	75			
	Overall	Accurac	cy: 85%	Kappa	a: 0.80)		

Appendix B. Continued.

- 1		Class	sified Ma	ap		E	7	
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
Sangamon	County							
Crop	40	1	0	0	41	2.4	2.4	97.6
Grass	1	16	0	0	17	5.9	17.6	94.1
Woods	0	2	16	0	18	11.1	0.0	88.9
Water	0	0	0	13	13	0.0	0.0	100.0
Sum	41	19	16	13	89			
	Overall	Accurac	y: 96%	Kappa	a: 0.93			
Schuyler	County							
Crop	23	5	0	0	28	17.9	7.1	82.1
Grass	2	11	0	0	13	15.4	38.5	84.6
Woods	0	0	20	0	20	0.0	0.0	100.0
Water	0	0	0	11	11	0.0	0.0	100.0
Sum	25	16	20	11	72			
	Overall	Accurac	y: 90%	Kappa	a: 0.87			
Scott Co	 unty							
Crop	23	2	0	0	25	8.0	4.0	92.0
Grass	1	18	0	0	19	5.3	10.5	94.
Woods	0	0	21	0	21	0.0	0.0	100.0
Water	0	0	0	8	8	0.0	0.0	100.0
Sum	24	20	21	8	73			
	Overall	Accurac	y: 96%	Kappa	a: 0.94			
Shelby C	ounty							
Crop	24	0	0	0	24	0.0	25.0	100.0
Grass	6	13	0	0	19	31.6	10.5	68.4
Woods	0	2	18	0	20	10.0	0.0	90.0
Water	0	0	0	6	6	0.0	0.0	100.0
Sum	30	15	18	6	69			
	Overall	Accurac	y: 88%	Kappa	a: 0.84			

Appendix B. Continued.

Land Use		Classi	fied M	ap		E:	rror(%)	A cours ou
Class ^b	Crop	Grass 1	Voods	Water	Sum	Omission	Commission	Accuracy (%)
Stark Co	ounty -	no accura	cy ass	essment				
St. Clai	 ir Count							
Crop	25	0	0	0	25	0.0	12.0	100.0
Grass	3	13	0	0	16	18.8	0.0	81.2
Woods	0	0	13	0	13	0.0	0.0	100.0
Water	0	0	0	6	6	0.0	0.0	100.0
Sum	28	13	13	6	60			
	Overall	Accuracy	: 95%	Kappa	: 0.93			
Stephens	on Coun	 ty						
Crop	31	0	1	0	32	3.1	31.3	96.9
Grass	10	15	0	0	25	40.0	0.0	60.0
Woods	0	0	23	0	23	0.0	4.3	100.0
Water	0	0	0	8	8	0.0	0.0	100.0
Sum	41	15	24	8	88			
	Overall	Accuracy	: 88%	Kappa	: 0.82			
Tazewell	County							
Crop	34	0	0	0	34	0.0	8.8	100.0
Grass	3	17	0	0	20	15.0	0.0	85.0
Woods	0	0	11	0	11	0.0	0.0	100.0
Water	0	0	0	5	5	0.0	0.0	100.0
Sum	37	17	11	5	70			
	Overall	Accuracy	: 96%	Kappa	: 0.93			
Union Co	ounty							
Crop	16	3	0	0	19	15.8	10.5	84.2
Grass	2	11	0	0	13	15.4	23.1	84.6
Woods	0	0	16	0	16	0.0	0.0	100.0
Water	0	0	0	12	12	0.0	0.0	100.0
Sum	18	14	16	12	60			

Overall Accuracy: 92% Kappa: 0.89

Appendix B. Continued.

T 1 TT		Clas	sified M	ap		E	7	
Land Use Class ^b	Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
Vermilio	n Count	У						
Crop	35	0	0	0	35	0.0	5.7	100.0
Grass	2	14	0	0	16	12.5	0.0	87.5
Woods	0	0	8	0	8	0.0	0.0	100.0
Water	0	0	0	6	6	0.0	0.0	100.0
Sum	37	14	8	6	65			
	Overall	Accura	cy: 97%	Kappa	a: 0.95)		
Wabash C	ounty							
Crop	26	0	0	0	26	0.0	19.2	100.0
Grass	5	11	0	0	16	31.3	0.0	68.7
Woods	0	0	16	0	16	0.0	0.0	100.0
Water	0	0	0	10	10	0.0	0.0	100.0
Sum	31	11	16	10	68			
	Overall	Accura	cy: 93%	Kappa	a: 0.90)		
Warren C	ounty							
Crop	19	0	0	0	19	0.0	36.8	100.0
Grass	6	10	0	0	16	37.5	0.0	62.5
Woods	1	0	14	0	15	6.7	0.0	93.3
Water	0	0	0	3	3	0.0	0.0	100.0
Sum	26	10	14	3	53			
	Overall	Accura	cy: 87%	Kappa	a: 0.81			
 Washingt	on Coun	ty						
Crop	24	1	0	0	25	4.0	28.0	96.0
Grass	6	14	0	0	20	30.0	5.0	70.0
Woods	0	0	21	0	21	0.0	0.0	100.0
Water	1	0	0	11	12	8.3	0.0	91.7
Sum	31	15	21	11	78			

Appendix B. Continued.

T 1 TT-	_	Classi	fied M	ap		E	7	
Land Use	e ———— Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy (%)
Wayne Co	ounty							
Crop	19	2	1	0	22	13.6	45.5	86.4
Grass	10	9	2	0	21	57.1	9.5	42.9
Woods	0	0	20	0	20	0.0	15.0	100.0
Water	0	0	0	16	16	0.0	0.0	100.0
Sum	29	11	23	16	79			
	Overall	Accuracy	: 81%	Kappa	a: 0.75			
White Co	 ounty							
Crop	15	2	0	0	17	11.8	29.4	88.2
Grass	5	7	0	0	12	41.7	16.7	58.3
Woods	0	0	13	0	13	0.0	0.0	100.0
Water	0	0	0	5	5	0.0	0.0	100.0
Sum	20	9	13	5	47			
Suill	20	9	13	5	4 /			
	Overall	Accuracy	: 85% ·	Kappa 	a: 0.79			
Whitesi	de Count	У						
Crop	25	1	0	0	26	3.8	19.2	96.2
Grass	5	14	0	0	19	26.3	10.5	73.7
Woods	0	1	18	0	19	5.3	0.0	94.7
Water	0	0	0	1	1	0.0	0.0	100.0
Sum	30	16	18	1	65			
	Overall	Accuracy	: 89%	Kappa	a: 0.84			
Will Co	 unty							
Crop	22	3	0	0	25	12.0	0.0	88.0
Grass	0	10	0	0	10	0.0	30.0	100.0
Woods	0	0	10	0	10	0.0	0.0	100.0
Water	0	0	0	5	5	0.0	0.0	100.0
Sum	22	13	10	5	50			
	Overall	Accuracy	: 94%	Kappa	a: 0.91			

Appendix B. Continued.

		Classi	fied Ma	ар		E:	rror(%)	3
Land Use Class ^b	e ——— Crop	Grass	Woods	Water	Sum	Omission	Commission	Accuracy
William		L						
WIIII	son Coun	Ly						
Crop	19	3	0	0	22	13.6	27.3	86.4
Grass	6	19	0	0	25	24.0	12.0	76.0
Woods	0	0	23	0	23	0.0	0.0	100.0
Water	0	0	0	10	10	0.0	0.0	100.0
Sum	25	22	23	10	80			
	Overall	Accuracy	·: 89%	Kappa:	: 0.85			
Winnebag	go Count	у У						
Crop	31	0	1	0	32	3.1	31.3	96.9
Grass	10	15	0	0	25	40.0	0.0	60.0
Woods	0	0	23	0	23	0.0	4.3	100.0
Water	0	0	0	8	8	0.0	0.0	100.0
Sum	41	15	24	8	88			
	Overall	Accuracy	7: 88%	Kappa:	: 0.82			
Woodford	d County							
Crop	41	0	1	0	42	2.4	2.4	97.6
Grass	1	11	0	0	12	8.3	0.0	91.7
Woods	0	0	9	0	9	0.0	11.1	100.0
Water	0	0	0	7	7	0.0	0.0	100.0
Sum	42	11	10	7	70			
	Overall	Accuracy	7: 97%	Kappa:	: 0.95			

^aCalculated according to Congalton 1991.

 $^{^{\}rm b}\text{Ground}$ truth data from Illinois Department of Natural Resources quail and pheasant call count routes.

JOB B. <u>Deer Ecology and Life History in Westcentral and Northern</u> Illinois

Objectives: (1) To determine age specific natality and seasonal and annual survival rates for deer in westcentral and northern Illinois; (2) to determine seasonal movement patterns and habitat selection of deer in Westcentral and Northern Illinois; (3) to integrate the natality and survival data collected from this study within new population models for the Illinois deer herd.

This job and its objectives are assigned to the Illinois
Natural History Survey and are reported in a separate document.

JOB C. Population Analysis

Objectives: (1) To develop interactive, menu-driven, portable computer models and software packages to analyze population data, model herd performance, and predict outcome of alternative harvest strategies on herd size, herd composition, and hunter behavior and success; and (2) to assist the Illinois Department of Natural Resources in integrating this system into their deer management program.

INTRODUCTION

Since the inception of modern firearm deer hunting in 1957,
Illinois has required mandatory check-in of all deer harvested.
This practice has resulted in perhaps the most extensive and
detailed deer harvest data set available to any state agency.
The primary purpose of Job C was to assemble this and
supplementary information into an accessible database and provide
the tools for its analysis.

METHODS

Initial software design was accomplished in close consultation with Division of Wildlife Forest Wildlife Program

managers and biologists. Final design and function of the computer programs were based primarily on management needs and data availability. Software was written in Microsoft® FORTRAN version 5.0 and originally compiled on a Gateway 2000 486/66 microcomputer. The program is designed to run on PC-compatible microcomputers using DOS 3.0 or later and i386 or larger processors. The executable file requires 286k of storage and the current data files requires 4.8mb of storage.

RESULTS

Objectives of Job C were met with the development and distribution of the Illinois Deer Harvest Analysis and Modeling Program (IDHAMP). This menu-driven PC-based computer program is designed to store, retrieve, and analyze historical white-tailed deer harvest data from Illinois and to model and simulate alternative future harvest strategies. The program can retrieve certain basic shotgun harvest data from 1957 to the present. Harvest data can also be retrieved separately for archery, muzzleloader, and handgun seasons. Data can be retrieved and examined by individual county, special area, management region, or statewide. Data also can be retrieved by day of season or permit type. Selected data output includes percent hunter

success, sex/age composition of the harvest, fawn:doe ratios, population sex ratios, indices of male mortality rates and female harvest rates, potential prevalence of trophy males (≥3.5 years old), and hunter and harvest densities per square mile of total area and woods. Estimates of population size and trends based on population reconstruction, kill/effort, various sex/age harvest indices, and population modeling are also available in graphic or tabular form. Alternative harvest strategies also can be simulated and evaluated using the program's modeling capabilities. A modified version of IDHAMP that does not include modeling capabilities also was produced and distributed.

Two user's manuals describing all program functions and how to use them are appended to this report. Also appended is a programmer's guide which includes a description of all program subroutines, a list of all variable names and their definitions, and the complete source code for IDHAMP.

JOB D. Analyze and Report

Objectives: (1) To analyze results and prepare products from Jobs A and B; and (2) report and discuss findings and present products in a timely manner.

Objectives for this job were met by development of the products to meet the objectives of Jobs A and C, and timely reporting of progress by means of quarterly progress reports and annual performance reports. Also, meetings were held with Division of Wildlife Resources Forest Wildlife Program staff to implement software application. Finally, Roseberry presented the following paper at the 1995 SE Deer Study Group meeting:

Roseberry, J. L., P. Shelton, J. Kube, and A. Woolf. 1995. Computer assisted management of white-tailed deer in Illinois. 18th Ann. Meeting of the Southeastern Deer Study Group, San Antonio, TX. February 26-28, 1995.