

2009

Establishment of the Regeneration System for *Vicia faba* L.

Shimaa Bahgat

Omer A. Shabban

Osama El-Shihy

David A. Lightfoot

Southern Illinois University Carbondale, ga4082@siu.edu

Hany A. El- Shemy

Follow this and additional works at: http://opensiuc.lib.siu.edu/psas_articles

Published in *Current Issues in Molecular Biology*, (2009) 11 (Suppl. 1): i47-54

Recommended Citation

Bahgat, Shimaa, Shabban, Omer A., El-Shihy, Osama, Lightfoot, David A. and El- Shemy, Hany A. "Establishment of the Regeneration System for *Vicia faba* L." (Jan 2009).

This Article is brought to you for free and open access by the Department of Plant, Soil, and Agricultural Systems at OpenSIUC. It has been accepted for inclusion in Articles by an authorized administrator of OpenSIUC. For more information, please contact opensiuc@lib.siu.edu.

Establishment of the Regeneration System for *Vicia faba* L.

Shimaa Bahgat¹, Omer A. Shabban¹, Osama El-Shihy², David. A. Lightfoot³ and Hany A. El-Shemy^{1,3*}

¹Faculty of Agriculture Research Park (FARP) and Biochemistry Department, Faculty of Agriculture, University of Cairo, 12613 Giza, Egypt

²Plant Physiology Department, University of Cairo, Faculty of Agriculture, 12613 Giza, Egypt

³Center of Excellence: The Illinois Soybean Center, Department of Plant, Soil and Agriculture Systems, SIUC, Carbondale, IL 62901, USA

Received 4 August 2008

Revised 13 September 2008

Accepted 11 October 2008

Abstract

A reliable regeneration system for faba bean has been difficult to establish and therefore, the genetic improvement of *Vicia faba* L. was delayed. The paper describes a method of somatic embryo induction in callus of *V. faba*. Two Egyptian faba bean cultivars 'Giza 2' and '24 Hyto' were used. Callus was induced from epicotyls and shoot tips cultured on MS or Gamborg medium supplemented with 3% sucrose and 0.025% (w/v) for each of ascorbic and citric acid, 0.8% agar and different concentrations of 10 mg/l BAP, 0.5 mg/l of each NAA and 2,4-dichlorophenoxyacetic acid (M1) and 1 mg/l BAP and 0.5 mg/l NAA (M2). The media with BAP, NAA and 2,4-D were optimal for embryogenic callus induction. Somatic embryos developed after transfer of the callus to 1/2 B5 medium with no plant growth regulators. There were various stages of somatic embryo development present including globular, heart-shaped, torpedo, and cotyledonary stages. Embryos developed into plantlets and plants were regenerated. RAPD analyses were performed to investigate the genetic stability of the regenerated plants obtained from different treatments and different explants. The cultivar Giza 2 exhibited more genetic stability than cultivar 24 Hyto. In conclusion, a regeneration system was established suitable for both gene transformation and the isolation of somaclonal mutants. The regeneration system will be used in order to improve the nutritional value of faba bean.

Introduction

Legumes are the third largest family of dicotyledons, *Leguminosae*. Examples include soybean (*Glycine max*), pea (*Pisum sativum*), faba bean (*Vicia faba*), mung bean

(*Vigna radiata*), chickpea (*Cicer arietinum*) and common bean (*Phaseolus vulgaris*). Faba bean (*Vicia faba* L.) is one of the most important grain Legumes in the world. Because Faba bean is an excellent source of protein, it is an important pulse crop used for human consumption and animal feeding, faba bean also plays a great role in the biological fixation of aerial nitrogen (Duke, 1981; Jelenić *et al.*, 2000). This species alone occupies nearly 3.2x10⁶ ha worldwide in 1991 (FAO statistics, 1992) with a world production close to 4.5 million tons in 2004 (Gutierrez, 2006). Even though faba bean has been cultivated in many countries, 49% of the total production comes from China followed by Egypt (10, 8%), Ethiopia (8.5%), and Australia by 5.8% in 2000 (FAO, statistics 2001). Unfortunately, its susceptibility to environmental conditions, biotic and abiotic stress and instability of the yield made this crop less attractive. Also, difficulties in pollination control and the limited genetic pool has led to slow down progress in varietal improvement (Bond, 1987; Selva *et al.*, 1989). Today to improve faba against biotic and abiotic stress and to improve its nutritional value with sulphur amino acids, traditional breeding programs could be supplemented by gene transfer technology which requires an efficient and suitable regeneration system, which occurs via organogenesis or embryogenesis, either directly from the excised tissue or indirectly after formation of callus depending on internal and external plant factors (Kuchuk, 2001). In tissue culture faba bean appears to be recalcitrant towards *in vitro* regeneration (Khalafalla and Hattori, 2000), due to difficulties in regeneration from callus, and the releasing of phenolic compounds, so studies on *in vitro* culture of faba bean are therefore difficult to carry out (Böttinger *et al.*, 2001). Regeneration system of faba bean through indirect somatic embryogenesis was reported by Griga *et al.* (1987) they were used the cotyledons on media supplemented with 2,4D. Somatic embryos at globular stage to early torpedo stage were released, but their development was terminated at the late-torpedo stage. In tissue culture, the regenerated plants from cultured cells are known to exhibit genetic and epigenetic changes collectively called somaclonal variations (Razdan, 1994). In this study we described a new method for regeneration of faba bean through somatic embryogenesis, with a development to complete plants. RAPD-PCR molecular markers were performed to identify the genetic variability between the control plants and the regenerated plants of each cultivar.

Materials and methods

Plant material

Two cultivars of faba bean Giza 2 and 24 Hyto (cv. Giza 2 was obtained from Agricultural Research Center, Egypt and cv. 24 Hyto was obtained from Agronomy Department,

*For correspondence: helshemy@hotmail.com

Faculty of Agriculture, Cairo University, Egypt) were used for this study. Segments of shoot tips and epicotyls were the sources of *in vitro* experimental materials.

Preparation of explants

Seeds of faba bean were sterilized in 70% (v/v) ethanol for 30 second, followed by 20% (v/v) commercial bleach for 15min and washed 3–4 times with sterile distilled water. Sterilized seeds were germinated on half-strength MS medium (Murashige and Skoog, 1962) supplemented with 15g/l sucrose and solidified with 8g/l agar. The seeds were germinated in darkness at 25°C. After a week shoot tips and epicotyls were cut with scalpel blade into segments of 5–7mm before cultivation on callus induction media.

Culture media

Two different treatments (M1& M2) were used in this study. The MS medium and Gamborg medium (Gamborg *et al.* 1968) were used for plant tissue culture. All of callus and regeneration media were supplemented with 3% (w/v) sucrose and 0.025% (w/v) for each of ascorbic and citric acid (Sigma, USA). The pH of all cultures was adjusted to 5.7 then solidified with 0.8% (w/v) agar and autoclaved at 121°C for 25 min. The cultures were incubated at 25°C ± 1 under a 16/8h light/dark photoperiod regime provided by white fluorescent light at 716 Lux.

M1 treatment

Epicotyls and shoot tips of the two cultivars were placed in callus induction medium CIM1 which contained MS medium with 10 mg/l BAP, 0.5 mg/l of each NAA and 2, 4D (Sigma, USA), and sub-cultured every 3–4 weeks, after two months calluses were transferred to a regeneration medium RM1 containing Gamborg medium or MS medium supplemented with B5 vitamins and 2 mg/l BAP. When globular structures appeared on the surface of callus with a white-greenish brilliant color, or a single embryo with a long structure had formed, they were separated from the callus and transferred to RM1 medium supplemented with 20 mg/l of glutamine (Sigma, St Louis, MO, USA). After 4–6 weeks, Nemours somatic embryos were formed (at globular stages) on an embryogenic callus, they were transferred to hormone free medium 1/2 B5 to encourage the development of the somatic embryos into plants. They were sub-cultured every 4 weeks. The hormone-free medium caused the formation of hundreds of somatic embryos. Each was separated and allowed to develop into a plant. Normally they formed shoots and roots in the same medium. Such normal plantlets were transferred to the soil, covered at first for adaptation to the glasshouse. Covers were gradually removed to allow hardening. Plants were allowed to flower and self fertilize.

M2 treatment

Epicotyls and shoot tips of the faba bean cvs. were placed in the callus induction medium CIM2 which contained MS medium with 1 mg/l BAP and 0.5 mg/l NAA. Within 6–7 weeks calluses were formed and embryo-like structures started to be formed. The embryo-like structures were excised and transferred to the regeneration medium RM2, containing MS medium supplemented with B5 vitamins,

1.2 mg/l BAP, and 0.5 mg/l NAA. Somatic embryos were formed at different stages. They were excised when they appeared to be at the beginning of the cotyledonary stage. Embryos developed to complete plants in 1/2 B5 medium without plant growth regulators. After the elongation of shoots and roots, plantlets were transferred to the soil to prepare them for adaptation to glass-house conditions.

RAPD analysis

Total DNA was isolated from each control and regenerated plant using the CTAB protocol (Doyle and Doyle, 1987). PCR reactions were performed in total volume of 25µl and 6 random Primers (from the OPA set) were used as described in Table 1 (all primers were obtained from Operon Technologies Inc., Alameda, CA, USA). DNA amplification was carried out in a T-gradient thermal cycler (Biometra, Göttingen, FRG), PCR reactions were hot –started at 95°C for 4 min and subjected to 40 cycles as follows; 1 min at 94°C; 2 min at 35°C; and 2 min at 72°C. The last extension phase was prolonged to 7 min at 72°C. PCR products were separated by electrophoresis in 1.5 % agarose gel and visualized by Ultraviolet illumination after staining with ethidium bromide, and photographed using gel-documentation system (BIO-RAD, Hercules CA).

Results

Callus tissues were initiated from epicotyls segments and shoots tip of two Egyptian faba bean cvs. Giza 2 and 24 Hyto using two different treatments M1 and M2. All the induction media were supplemented with various concentrations of auxins and cytokinins, segments of shoot tips and epicotyls induced a mixture of greenish, brownish and yellowish compact callus (Fig. 1A-F).

The M1 treatments consisted of CIM1 (callus induction medium) and RM1 (regeneration medium). In CIM1 shoot tips and epicotyls of each cultivar were cultured for two months during which time they were sub-cultured every 3–4 weeks. They had produced a compact yellowish callus; after two months the calluses were transferred to regeneration medium RM1. After 3–7 weeks, structures that looked like an embryo with long structure were obtained, or sometimes a globular smooth structures with white-greenish transparent color, were formed on the surface of callus-derived from shoot tips. Cultivar Giza 2 formed these structures after 3 weeks while cv. 24 Hyto formed the same structures after 7 weeks of culturing in RM1 medium. The embryo-like structures were separated and cultured in the same RM1 medium, but supplemented with 20 mg/l glutamine. Within a week these embryo

Table 1. Nucleotide sequences of six primers used in RAPD analysis.

Sequence (5'-3')	Primer
AATCGGGCTG	OPA04
AGGGGTCTTG	OPA05
GAAACGGGTG	OPA07
GGGTAACGCC	OPA09
GTGATCGCAG	OPA10
CAGCACCCAC	OPA13




Fig. 1 Examples of embryogenic callus from *V. faba* cvs. Giza 2 and 24 Hyto. Panels A and B show segments of shoot tips and epicotyls. Panels C and D show segments of shoot tips and epicotyls on callus induction media after 10–15 days. Panels E and F show shoot tips and epicotyls respectively after two months on callus induction medium. Panel E shows callus derived from shoot tips of cv. 24 Hyto on CIM1 medium. Panel F shows callus derived from an epicotyl of cv. 24 Hyto in CIM2 medium which turned brown due to phenolic compounds.

structures grew and developed into globular stage (Fig. 2F, G), 3 weeks later, a lot of somatic embryos with white-greenish color were formed in clusters with embryogenic callus (Fig. 2H). For further development these embryos were transferred to hormone free medium (1/2 B5 medium). It was also noted that, the hormone free medium enhanced the production of new generations of somatic embryos. Each embryo was capable of producing a lot of embryos over a period of 14 months. Therefore there was asynchronous development of embryos to complete plants over that time. The somatic embryos which derived from shoot tips of cv. Giza 2 readily developed to fertile plants, while the cultivar 24 Hyto did not, being arrested at the torpedo stage. Although the embryos of cv. Giza 2 developed slowly, this cultivar produced more embryos than that of cv. 24 Hyto (Table 2).

Epicotyls from cv. Giza 2 which cultured on M1 also formed embryogenic callus with somatic embryos in the

Table 2. Effect of cultivar generally on fresh weight of callus, Callus (%) and regeneration through somatic embryogenesis (%).

Cultivar	Fresh weight of callus	Callus (%)	Calluses producing somatic embryos (%)
Giza 2	2.14 ^b	69.38 ^b	4.14 ^a
24 Hyto	3.69 ^a	72.91 ^a	3.31 ^b

Means followed by the same letter are not significantly different at $p \leq 0.05$ using Duncan's multiple range test.

globular stage. However, these globular structures did not develop further and eventually became contaminated. The experiment was repeated many times without any success. Epicotyls from cv. 24 Hyto formed callus but not embryos.

The other treatment (M2) induced callus from epicotyls of each cv. but failed to induce callus from shoot tips where, shoots formed directly (Fig. 3A). These shoots were transferred to 1/2 B5 medium supplemented with 3 mg/l IBA for rooting. Epicotyls from cv. Giza 2 induced small calluses with a greenish- yellow color, but all failed to produce any somatic embryos. Only a few adventitious shoots were obtained from buds on the callus surface (Fig. 3B).

Calluses derived from epicotyls of 24 Hyto formed creamy- brownish masses (Fig. 1F), that increased in weight rapidly (Table 5). These calluses were embryogenic after 7 weeks of culturing on CIM2 medium (Fig. 2A). To develop the embryos, calluses were transferred to RM2 medium (Fig. 2B). Calluses developed and produced somatic embryos in clusters at different growth stages including globular and heart stages and the early torpedo stages (Fig. 2C). The embryos were separated at the beginning of the cotyledonary stage (Fig. 2D), and each transferred to hormone free medium (1/2 B5) for their development. The hormone free medium increased the numbers of embryos formed by stimulating many generations of somatic embryos with different stages forming on the first embryo explants. The adventitious embryos developed in the same medium to become young plantlets (Fig. 2E). These plantlets then transferred to the green house for adaptation.

The number of total shoots which developed from somatic embryos varied between each cultivar. In cultivar Giza 2, the calluses that derived from shoot tips were reproducible and produced a lot of somatic embryos. Hence within 8 months after the formation of the first embryo, 400 shoots per callus were developed from somatic embryos. However, in cultivar 24 Hyto, within 8 months, only 127 shoots had developed from somatic embryos derived from the epicotyl calluses.

Therefore cultivar differences and explant differences play important roles determining both callus formation and regeneration percentages; the type of structures that develop callus and the regeneration frequency of those structures (Tables 2 and 3).

A significant response towards callus, and regeneration percentage were observed, due to the hormonal compositions of each treatment (Table 4). Although the M2 medium had the highest value of the

Table 3. Effect of explants generally on, fresh weight of callus, callus (%) and regeneration through somatic embryogenesis (%) for both cultivars (Giza 2 and 24 Hyto).

Type of explant	Fresh weight of callus	Callus (%)	Calluses producing somatic embryos (%)
Shoot tip	1.64 ^b	49.58 ^b	6.76 ^a
Epicotyl	4.19 ^a	92.71 ^a	0.70 ^b

Means followed by the same letter are not significantly different at $p \leq 0.05$ using Duncan's multiple range test.




Fig. 2. *In vitro* regeneration through indirect somatic embryogenesis from *V.faba* cvs. Giza 2 and 24Hyto. **A–E** Somatic embryos derived from calluses of epicotyls of cv. 24 Hyto. **A** Embryogenic cells formed on callus derived from epicotyls in CIM2 medium. **B** Differentiation of the embryogenic cells in RM2 medium. **C** Different stages of somatic embryos were formed in RM2 medium including globular, heart and torpedo stages. **D** Maturation of somatic embryos into the cotyledonary stage in hormone free medium with asynchronous development. **E** The development of somatic embryos to young plantlets. **F–N** Different stages of somatic embryos of cv. Giza 2 which formed on RM1 medium from calluses derived from shoot tips. **F–G** The stages of embryos formation including formation of globular stage from a single embryo, this process took place in RM1 medium supplemented with 20mg/l glutamine. **H** White-green structures identified as the early stages of somatic embryos were obtained on the previous medium including formation of embryogenic callus, these embryos were transferred subsequently to hormone free medium for their development. **I–K** Different stages of somatic embryos were formed in clusters in hormone free medium. **L** A cluster of somatic embryos developed before separating. **M** Development of embryos to young plantlets with shoots and roots in the hormone free medium. **N** Rooted plants that were transferred to the soil for acclimatization.




Fig. 3. **A** Direct organogenesis was formed from the shoot tips of cv. Giza 2 on CIM2 medium. **B** Adventitious shoots were formed from callus of epicotyls of cv. Giza 2 on M2 treatment.

callus fresh weight, the M1 treatment was the most effective in each of regeneration and callus percentages, so we detected that there was not any relationship between the weight of callus and the frequency of the regeneration.

Also the interactions between the three elements of this study (cultivars, explants and treatments) in all of the callus and regeneration percentages, showed that the best explant which led to the highest percentage of regeneration was the shoot tip of cv. Giza 2 using M1 treatment (Tables 5–8).

RAPD analysis

RAPD analyses were performed to investigate the genetic stability of the regenerated plants of each cultivar obtained from different treatments and different explants. Therefore six random primers were used (Table 1) to amplify DNA from many genomic locations. The 100 bp DNA molecular marker (with range of 100, 200, 300, 400, 500, 600, 700, 800, 900 and 1,000 bp) was used in this experiment. Among the RAPD profiles (Fig. 4) the cultivar Giza 2 showed identity between the control and the earliest regenerated plant in all profiles, except for new fragments detected at 1000 bp (OPA-04), 550 bp (OPA-05), and 500 bp of (OPA-07). Cultivar 24 Hyto showed identity in DNA banding patterns between the control and the earliest regenerated plant only with primer OPA-13. The other 5 primers derived DNA banding patterns showed new fragments at 300 bp (OPA-05) and 400 bp (OPA-09), whereas some fragments were absent at 750 bp (OPA-04), 450 bp (OPA-07) and 800 bp (OPA-10). Therefore, the cultivar Giza 2 exhibited more genetic stability than cultivar 24 Hyto.

Table 4. Effect of treatments generally on fresh weigh of callus, callus (%) and regeneration through somatic embryogenesis(%) for both cultivars (Giza 2 and 24 Hyto).

Treatment	Fresh weight of callus	Callus (%)	Calluses producing somatic embryos (%)
M1	2.66 ^a	97.52 ^a	6.89 ^a
M2	3.17 ^a	44.78 ^b	0.56 ^b

Means followed by the same letter are not significantly different at $p \leq 0.05$ using Duncan's multiple range test.

Table 5. Effect of interaction between treatments and cultivars on fresh weight of callus, callus (%), and regeneration through somatic embryogenesis (%).

Treatment	Cultivar	Fresh weight of callus	Callus (%)	Calluses producing somatic embryos (%)
M1	Giza 2	2.99 ^a	96.20 ^b	8.28 ^a
	24 Hyto	2.33 ^a	98.83 ^a	5.50 ^b
M2	Giza 2	1.29 ^a	42.57 ^d	0.00 ^d
	24 Hyto	5.05 ^a	46.98 ^c	1.12 ^c

Means followed by the same letter are not significantly different at $p \leq 0.05$ using Duncan's multiple range test.

Table 6. Effect of interaction between treatments and explants on fresh weight of callus, callus (%), and regeneration through somatic embryogenesis (%) for both cultivars (Giza 2 and 24 Hyto).

Treatment	Explant	Fresh weight of callus	Callus (%)	Calluses producing somatic embryos (%)
M1	Shoot tip	3.28 ^b	99.17 ^a	13.51 ^a
	Epicotyl	2.04 ^c	95.87 ^b	0.27 ^c
M2	Shoot tip	0.00 ^d	0.00 ^d	0.00 ^c
	Epicotyl	6.35 ^a	89.55 ^c	1.12 ^b

Means followed by the same letter are not significantly different at $p \leq 0.05$ using Duncan's multiple range test.

Table 7. Effect of interaction between cultivars and explants on fresh weight of callus, callus (%), and regeneration through somatic embryogenesis (%).

Cultivar	Explant	Fresh weight of callus	Callus (%)	Calluses producing somatic embryos (%)
Giza 2	Shoot tip	1.71 ^c	49.33 ^c	8.01 ^a
	Epicotyl	2.57 ^b	89.43 ^b	0.27 ^d
24 Hyto	Shoot tip	1.57 ^c	49.83 ^c	5.50 ^b
	Epicotyl	5.82 ^a	95.98 ^a	1.12 ^c

Means followed by the same letter are not significantly different at $p \leq 0.05$ using Duncan's multiple range test.

Discussion

This study describes a new reproducible system for plant regeneration via somatic embryogenesis using callus cultures derived from shoot tips and epicotyls of two Egyptian cultivars of *V. faba*. Many investigators have suggested that the difficulties of indirect regeneration from faba bean were due to the accumulation of phenolic compounds, which led to cell death (Böttinger *et al.*, 2001). Here it was shown to be very important to avoid the accumulation of the phenolic compounds within the first two weeks of culturing the explants on the callus induction media. However some calluses that turned to brown or black colors later (Fig. 1 F), presumably due to the phenolic compounds, still showed a good ability to divide and produce several generations of somatic embryos. (Fig. 2 A, B, I, G). The addition of ascorbic and citric acids to the media helped to decrease or prevent the production or oxidation of the phenolic compound, therefore enhancing the production of somatic embryos. The addition of citric acid to media was reported by Anthony *et al.*, 2004, to

Table 8. Effect of interactions among treatments, cultivars and explants on fresh weight of callus, callus (%), and regeneration through somatic embryogenesis (%).

Treatment	Cultivar	Explant	Fresh weight of callus	Callus (%)	Calluses producing somatic embryos (%)
M1	Giza 2	Shoot tip	3.42 ^b	98.67 ^a	16.02 ^a
		Epicotyl	2.55 ^b	93.73 ^b	0.53 ^d
	24 Hyto	Shoot tip	3.13 ^b	99.67 ^a	11.00 ^b
		Epicotyl	1.53 ^c	98.00 ^a	0.00 ^d
M2	Giza 2	Shoot tip	0.00 ^d	0.00 ^d	0.00 ^d
		Epicotyl	2.59 ^b	85.13 ^c	0.00 ^d
	24 Hyto	Shoot tip	0.00 ^d	0.00 ^d	0.00 ^d
		Epicotyl	10.11 ^a	93.97 ^b	2.25 ^c

Means followed by the same letter are not significantly different at $p \leq 0.05$ using Duncan's multiple range test.

prevent the browning and to enhance the formation of somatic embryos in *Conostephium pendulum*. Also the addition of ascorbic acid was beneficial for long-term common bean (*Phaseolus vulgaris*) callus maintenance (Mohamed *et al.*, 1993).

In this study, callus induction media used enrichment with cytokinin (as BAP) that was higher in concentration than auxins in each treatment. Cytokinin was important for the formation of callus and initiation of somatic embryogenesis. Recently, attempts to regenerate plants via somatic embryogenesis focused on pre-cultivation of the initial plants on cytokinin- enriched medium (Veltcheva *et al.*, 2005). High cytokinin to auxin ratios may modify the physiological state of the initial explants and initiate a somatic embryogenesis program.

The hormone free medium enhanced the development of somatic embryos through different stages (Fig. 5) and led to regeneration of complete plants (Pickardet *et al.*, 1989; Albrecht and Kohlenbach, 1989). Also the hormone

free medium enhanced the formation of secondary embryos.

Here, within 8 months after the formation of the first embryo, 400 shoots per callus were developed from the somatic embryos derived from shoot tips of the cultivar Giza 2 on M1 treatment. However, these calluses maintained embryogenic potential over a period of 14 months. The number of shoots was increased due to the effect of the hormonal composition of the M1 treatment. Further, the type of explant used (shoot tip) with its endogenous hormones may have played an important role in this regeneration system. In the M2 treatment within 8 months only 127 shoots per callus were developed from the somatic embryos that derived from calluses of epicotyl of cultivar 24 Hyto. The difference between each cultivar in the number of shoots was varied due to the type of explants used and the hormonal composition of each treatment. Cultivar genetics seems to have an important effect on the regeneration rate.




Fig. 4. RAPD profiles of faba bean cvs. were amplified with six RAPD primers. M: DNA molecular marker 100bp. Lane 1: Control plant of cv. Giza 2 Lane 2: Regenerated plant of cv. Giza 2 via somatic embryogenesis . Lane 3: Control plant of cv. 24 Hyto. Lane 4: Regenerated plant of cv. 24 Hyto via somatic embryogenesis.




Fig. 5. Different stages of somatic embryogenesis (globular stage, heart stage, torpedo stage, cotyledonary stages, and young plantlets) of cv. Giza 2 were formed in the hormone free medium.

During the development of somatic embryos some of the developed plantlets formed roots rapidly in the same hormone free medium. However, sometimes the number of roots was up to 3 per plantlet (Fig. 6). A delay in the formation of roots was observed in some plantlets, in some cases the normal shoots were developed but root formation was suppressed.

Difficulties in the regeneration from the callus tissues of faba bean led to a limited number of publications on cell and tissue culture of this species. A regeneration system of faba bean through indirect somatic embryogenesis was reported by Griga *et al.* (1987). An indirect somatic embryogenesis system for *Vicia narbonensis* was reported by Pickardet *et al.* (1989) and also reported by Albrecht and Kohlenbach (1989). Another kind of regeneration is the organogenesis that was described in faba bean by Taha and Francis, (1990), Tegeder *et al.* (1995) and Böttinger *et al.* (2001).



Fig. 6. Plants developed from somatic embryogenesis of cv. Giza 2 with number of roots ranged from 1–3 roots.

PCR-based RAPD (Random amplified polymorphic DNA) markers have been applied to many types of genetic studies because of their simplicity and ease of use (Gutierrez *et al.*, 2006). In this study, RAPD analysis was performed to identify potential somaclonal variations among the regenerated plants from somatic embryogenesis. To investigate the effects of the media and subcultures, on genetic stability, the regenerated plants were obtained after long-term of subcultures (after about 13 months of the first culture, from the later generations).

More new bands were detected in the regenerated plants of cultivar Giza 2 while more bands were missing in regenerated plants of cultivar 24Hyto. Both the deletions underlying the loss of DNA fragments and the modifications underlying the appearance of new fragments among the regenerated plants may be used to help to improve faba bean genetically. Furthermore, RAPD analysis could be used to trace the genetic changes in the regenerated plants at the genome level. RAPD polymorphism in callus-derived plants has been reported in wheat by Brown *et al.*, 1993, and turmeric (Salvi *et al.*, 2001). However, Isabel *et al.* (1993) detected no polymorphism in regenerated plants of *Picea Mariana*. The source of the explants, mode of regeneration (somatic embryogenesis/organogenesis/axillary bud multiplication), genotypes used and the plant growth regulators applied are known to play a major role in determining the presence or absence of variations (Razdan, 1994; Damasco *et al.*, 1996). Several investigators (Link *et al.*, 1995; Torres *et al.*, 1993; Vaz Patto *et al.*, 1999; Gutierrez *et al.*, 2006; and Gutierrez *et al.*, 2007) used RAPD molecular markers to study the genetic variability and relationships among accessions, lines and cultivars of faba bean. Therefore, a link between natural and induced genetic variation might be made by further analyses.

Acknowledgments

We thank Dr. Said Abd-ELTawab (Cotton Research institute, Agricultural Research Center, Egypt) for his assistance with statistical analysis.

References

- Albrecht, C., and Kohlenbach, H.W. (1989) Induction of somatic embryogenesis in leaf derived callus of *Vicia narbonensis* L. *Plant Cell Rep.* **8**, 267–269.
- Anthony, J.M., Senaratna, T., Dixon, K.W., and Sivasithamparam, K. (2004) The role of antioxidants for initiation of somatic embryos with *Conostephium pendulum* (Ericaceae) *Plant Cell Tiss. Org. Cult.* **78**, 247–252.
- Bond, D.A. (1987) Recent developments in breeding of field beans (*Vicia faba* L.) *Plant Breed.* **99**, 1–26.
- Böttinger, P., Steinmetz, A., Schieder, O., and Pickardt, T. (2001) *Agrobacterium*-mediated transformation of *Vicia faba*. *Mol. Breed.* **8**, 243–254.
- Brown P.T.H, Lange, F.D., Kranz, E., and Lörz, H. (1993) Analysis of single protoplasts and regenerated plants by PCR and RAPD technology. *Mol. Gen. Genet.* **237**, 311–317.
- Damasco, O.P., Graham, G.C., Henry, R.J., Adkins, S.W., Smith, M.K., and Godwin, I.D. (1996) Random amplified polymorphic DNA (RAPD) detection of dwarf off-types in micropropagated Cavendish (*Musa SPP AAA*) bananas. *Plant Cell Rep.* **16**, 118–123.
- Doyle, J.J., and Doyle, J.L. (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* **19**, 11–15.
- Duke, J.A. (1981) *Handbook of Legumes of World Economic Importance.* (New York: Plenum Press), pp. 199–265.
- Gamborg, O.L., Miller, R.A., and Ojima, K. (1968) Nutrient requirements of suspension cultures of soybean root cells. *Exp. Cell Res.* **50**, 151–158.
- Griga, M., Kubalaková, M., and Tejklová, E. (1987) Somatic embryogenesis in *Vicia faba* L. *Plant Cell Tiss. Org. Cult.* **9**, 167–171.
- Gutierrez, N., Avila, C.M., Duc, G., Marget, P., Suso, M.J., Moreno, M.T., and Torres, A.M. (2006) CAPs markers to assist selection for low vicine and convicine contents in faba bean (*Vicia faba* L.) *Theor. Appl. Genet.* **114**, 59–66.
- Gutierrez, N., Avalia, C.M., Rodriguez-Suarez, C., Moreno, M.T., and Torres, A.M. (2007) Development of SCAR markers linked to a gene controlling absence of tannins in faba bean. *Mol. Breed.* **19**, 305–314.
- Isabel, N., Tremblay, L., Michaud, M., Tremblay, F.M., and Bousquet, J. (1993) RAPDs as an aid to evaluate the genetic integrity of somatic embryogenesis-derived populations of *Picea mariana* (Mill.) B.S.P. *Theor. Appl. Genet.* **86**, 81–87.
- Jelenić, S., Mitrikeski, P.T., Papeš, D., and Jelaska, S. (2000) *Agrobacterium*-mediated transformation of broad bean *Vicia faba* L. *Food Technol. Biotechnol.* **38**, 167–172.
- Khalafalla, M., and Hattori, K. (2000) Ethylene inhibitors enhance *in vitro* root formation on faba bean shoots regenerated on medium containing thidiazuron. *Plant Growth Regul.* **32**, 59–63.
- Kuchuk, N. (2001) Biotechnology. In: *Carbohydrates in Grain Legume Seeds*, C.L. Hedly, ed. (Wallingford, UK: CABI Publishing), pp. 145–207.
- Link, W., Dixkens, C., Singh, M., Schwall, M., and Melchinger, A.E. (1995) Genetic diversity in European and Mediterranean faba bean germ plasm revealed by RAPD markers. *Theor. Appl. Genet.* **90**, 27–32.
- Mohamed, M.F., Coyne, D.P., and Read, P.E. (1993) Shoot organogenesis in callus induced from pedicel explants of common bean (*Phaseolus vulgaris* L.) *J. Amer. Soc. Hort. Sci.* **118**, 158–162.
- Murashige, T., and Skoog, F. (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* **15**, 473–497.
- Pickardt, T., Huancaruna Perales, E., and Schieder, O. (1989) Plant regeneration via somatic embryogenesis in *Vicia narbonensis*. *Protoplasma* **149**, 5–10.
- Razdan, M.K. (1994) *An Introduction to Plant Tissue Culture* (New Delhi: Oxford & IBH Publishing CO. PVT. LTD), pp. 225–243.
- Salvi, N.D., George, L., and Eapen, S. (2001) Plant regeneration from leaf base callus of turmeric and random amplified polymorphic DNA analysis of regenerated plants. *Plant Cell. Tiss. Org. Cult.* **66**, 113–119.
- Selva, E., Stouffes, B., and Briquet, M. (1989) *In vitro* propagation of *Vicia faba* L. by micro-cutting and multiple shoot induction. *Plant Cell Tiss. Org. Cult.* **18**, 167–179.
- Taha, R.M., and Francis, D. (1990) The relationship between polyploidy and organogenetic potential in embryo and root-derived tissue cultures of *Vicia faba* L. *Plant Cell Tiss. Org. Cult.* **22**, 229–236.
- Tegeger, M., Gebhardt, D., Schieder, O., and Pickardt, T. (1995) Thidiazuron-induced plant regeneration from protoplasts of *Vicia faba* cv. Mythos. *Plant Cell Rep.* **15**, 164–169.
- Torres, A.M., Weeden, N.F., and Martín, A. (1993) Linkage among isozym, RFLP and RAPD markers in *Vicia faba*. *Theor. Appl. Genet.* **85**, 937–945.
- Vaz Patto, M.C., Torres, A.M., Koblizkova, A., Macas, J., and Cubero, J.I. (1999) Development of a genetic composite map of *Vicia faba* using F₂ populations derived from trisomic plants. *Theor. Appl. Genet.* **98**, 736–743.
- Veltcheva, M., Svetleva, D., Petkova, SP., and Perl, A. (2005) *In vitro* regeneration and genetic transformation of common bean (*Phaseolus vulgaris* L.)— Problems and progress. *Scientia Horticulturae.* **107**, 2–10.

Caliciviruses

Molecular and Cellular Virology

Edited by: G.S. Hansman, J. Jiang, K.Y. Green
c. 250 pp., April 2010

ISBN: 978-1-904455-63-9 \$310 / £159

The most important research findings.
Timely and comprehensive reviews.
Discussion of past and current research.

Epstein-Barr Virus

Latency and Transformation

Edited by: Erle S. Robertson
c. 220 pp., April 2010

ISBN: 978-1-904455-62-2 \$310 / £159

Expert virologists comprehensively review this important subject from a genetic, biochemical, immunological, and cell biological perspective. Essential reading.

Anaerobic Parasitic Protozoa

Genomics and Molecular Biology

Edited by: C.G. Clark, P.J. Johnson, R.D. Adam
c. 210 pp., March 2010

ISBN: 978-1-904455-61-5 \$310 / £159

Internationally acclaimed researchers critically review the most important aspects of research on anaerobic parasitic protozoa.

Lentiviruses and Macrophages

Molecular and Cellular Interactions

Edited by: Moira Desport
c. 410 pp., March 2010

ISBN: 978-1-904455-60-8 \$310 / £159

Top lentivirus and macrophage specialists comprehensively review cutting-edge topics in the molecular and cellular biology of the lentivirus-macrophage interaction.

Microbial Population Genetics

Edited by: Jianping Xu
c. 230 pp., March 2010

ISBN: 978-1-904455-59-2 \$310 / £159

Details the major current advances in microbial population genetics and genomics.

Borrelia

Molecular Biology, Host Interaction and Pathogenesis

Edited by: D. Scott Samuels and Justin D. Radolf
c. 630 pp., March 2010

ISBN: 978-1-904455-58-5 \$310 / £159

Written by renowned scientists in the field who have made seminal contributions to the field, this book is a comprehensive guide to the pathogenic *Borrelia*.

Influenza

Molecular Virology

Edited by: Qinghua Wang and Yizhi Jane Tao
c. 200 pp., February 2010

ISBN: 978-1-904455-57-8 \$310 / £159

NS1, hemagglutinin, nucleoprotein, glycoproteins, M2 channel, virulence, polymerase, microarrays, vaccine design.

RNA Interference and Viruses

Current Innovations and Future Trends

Edited by: Miguel Angel Martínez
c. 280 pp., February 2010

ISBN: 978-1-904455-56-1 \$310 / £159

Expert RNAi specialists from around the world have teamed up to produce a timely and thought-provoking review of the area.

Retroviruses

Molecular Biology, Genomics and Pathogenesis

Edited by: Reinhard Kurth and Norbert Bannert
c. 520 pp., January 2010

ISBN: 978-1-904455-55-4 \$310 / £159

Genomics, molecular biology and pathogenesis, comprehensively covering all the recent advances.

Metagenomics

Theory, Methods and Applications

Edited by: Diana Marco
x + 212 pp., January 2010

ISBN: 978-1-904455-54-7 \$310 / £159

Essential reading for all researchers performing metagenomics studies. Highly recommended.

Aspergillus

Molecular Biology and Genomics

Edited by: M. Machida and K. Gomi
x + 238 pp., January 2010

ISBN: 978-1-904455-53-0 \$310 / £159

Systematics, bioinformatics, systems biology, regulation, genetics, genomics, metabolism, ecology, development.

Environmental Molecular Microbiology

Edited by: Wen-Tso Liu and Janet K. Jansson
viii + 232 pp., January 2010

ISBN: 978-1-904455-52-3 \$310 / £159

Current technology and applications. Microbial diversity, phylogeny, communities, 16S rRNA, metagenomics, metaproteomics, microarrays, fingerprinting, soil, water, plants, humans, biofilms.

Neisseria

Molecular Mechanisms of Pathogenesis

Edited by: Caroline Genco and Lee Wetzler
x + 270 pp., January 2010

ISBN: 978-1-904455-51-6 \$310 / £150

Genomics, biofilms, adhesion, invasion, immunity, complement, apoptosis, vaccine, epidemiology, antibiotic resistance.

Frontiers in Dengue Virus Research

Edited by: K.A. Hanley and S.C. Weaver
viii + 304 pp., January 2010

ISBN: 978-1-904455-50-9 \$310 / £150

Evolution, epidemiology, translation, replication, pathogenesis, host, animal models, mosquito interactions, transmission, vaccine, drugs, immunotherapy.

ABC Transporters in Microorganisms

Edited by: Alicia Ponte-Sucre
xii + 260 pp., August 2009

ISBN: 978-1-904455-49-3 \$310 / £150

Pili and Flagella

Current Research and Future Trends

Edited by: Ken Jarrell

x + 238 pp., August 2009

ISBN: 978-1-904455-48-6 \$310 / £150

Lab-on-a-Chip Technology

Edited by: K. E. Herold and A. Rasooly

Vol 1: Fabrication and Microfluidics

ISBN: 978-1-904455-46-2 \$310 / £150

Vol 2: Biomolecular Separation and Analysis

ISBN: 978-1-904455-47-9 \$310 / £150

Bacterial Polysaccharides

Current Research and Future Trends

Edited by: Matthias Ulbrich

xii + 358 pp., June 2009

ISBN: 978-1-904455-45-5 \$310 / £150

Microbial Toxins:

Current Research and Future Trends

Edited by: Thomas Proft

viii + 192 pp., May 2009

ISBN: 978-1-904455-44-8 \$310 / £150

Acanthamoeba:

Biology and Pathogenesis

Author: Naveed Khan

viii + 290 pp., February 2009

ISBN: 978-1-904455-43-1 \$310 / £150

Bacterial Secreted Proteins: Secretory Mechanisms and Role in Pathogenesis

Edited by: Karl Wooldridge

xii + 512 pp., April 2009

ISBN: 978-1-904455-42-4 \$310 / £150