

10-2017

A Pilot Study on the Effects of Curcumin on Parasites, Inflammation, and Opportunistic Bacteria in Riding Horses

Samantha Wuest

Southern Illinois University Carbondale, ssparks2@siu.edu

Rebecca Lynn Atkinson

Southern Illinois University Carbondale, ratkinsn@siu.edu

Stephanie D. Bland

Southern Illinois University Carbondale, sbland@siu.edu

Darcie Hastings

Southern Illinois University Carbondale, darueh2@siu.edu

Follow this and additional works at: http://opensiuc.lib.siu.edu/asfn_articles

In press June 29, 2017.

Recommended Citation

Wuest, Samantha, Atkinson, Rebecca L., Bland, Stephanie D. and Hastings, Darcie. "A Pilot Study on the Effects of Curcumin on Parasites, Inflammation, and Opportunistic Bacteria in Riding Horses." *Journal of Equine Veterinary Science* 57 (Oct 2017): 46-50. doi:10.1016/j.jevs.2017.06.010.

This Article is brought to you for free and open access by the Department of Animal Science, Food and Nutrition at OpenSIUC. It has been accepted for inclusion in Peer-reviewed Articles by an authorized administrator of OpenSIUC. For more information, please contact opensiuc@lib.siu.edu.

A Pilot Study on the Effects of Curcumin on Parasites, Inflammation, and Opportunistic Bacteria
in Riding Horses

S. Wuest[‡], R. L. Atkinson^{‡1}, S. D. Bland[‡] and D. Hastings

[‡]Department of Animal Science, Food & Nutrition, Southern Illinois University,
Carbondale, IL 62901

¹ Corresponding author: ratkinsn@siu.edu; 1205 Lincoln DR, Carbondale, IL 62901

1 **ABSTRACT:**

2 Twelve riding horses were utilized to examine the effects of curcumin on intestinal parasites,
3 inflammation, and the fecal shedding of *Streptococcus bovis/equinus* complex (SBEC),
4 *Clostridium difficile* and *Clostridium perfringens*. Known for having anti-inflammatory,
5 antimicrobial, and antiparasitic properties it was hypothesized that curcumin would decrease
6 parasite shedding, inflammation, and opportunistic bacteria found in the GIT of riding horses.
7 Horses were randomly assigned to one of the following treatments (n = 6/treatment): 1) no
8 curcumin, control (CON); or 2) 15 g of 95% pure curcumin, (CUR). Curcumin was dosed per
9 day for 30 d. Fecal samples were evaluated for shedding of ova and concentrations of selected
10 bacteria. Blood samples taken pre and post riding intervals and evaluated for erythrocyte
11 sedimentation rate (ESR) for inflammation. All data were analyzed for repeated measures.
12 Treatment had no effect ($P \geq 0.58$) on total fecal egg count, strongyles, or ascarids. Treatment
13 had no effect on ESR ($P \leq 0.42$); however, ESR decreased ($P = 0.0006$) on d 14 in CUR horses.
14 Treatment had no effect ($P \geq 0.34$) on concentrations of SBEC, *C. difficile*, or *C. perfringens*.
15 Curcumin was not an effective compound against intestinal parasites or fecal microbial strains
16 examined when administered for 30 days; but could potentially decrease inflammation.
17 Curcumin has been observed to have many beneficial effects in other species, however, more
18 research is needed to evaluate those benefits in horses.

19 **Keywords:** Curcumin; Equine; Inflammation; Opportunistic bacteria; Parasites

20 **1. INTRODUCTION**

21 Horse intestinal parasites pose an economic and health risk that are of concern to both
22 breeders and horse owners [1]. The body of a horse is host to millions of microscopic organisms

23 who utilize the horse's oxygen, nutrients, and body heat for survival. Parasites, such as
24 *Strongylidae* (strongyles) can cause emaciation and anemia; while ascarids are known to cause a
25 blockage in the intestines, which if not taken care of properly can lead to death [2,3]. Moreover,
26 parasites, which can be found in the intestines of horses of all breeds, both sexes, and all age
27 classes [4] can cause inflammation within the gastrointestinal tract (GIT).

28 In addition to parasite-induced inflammation, inflammation can also occur due to the
29 athletic lifestyle required of domesticated horses. Repetitive stress applied to the joints from
30 speed work, jumping, and extreme hindquarter thrust, results in inflammatory changes to the
31 bone structure, joint anatomy, and synovial fluid as well as predisposed factors. Although some
32 horses are diagnosed with lameness in their younger years, many develop the progressive
33 problem over time, whether it is mild or severe [5]. Therefore, inflammation in a horse can be
34 due to several factors including, illness, injury, GIT parasites, and even an altered hindgut
35 microbiome.

36 The gut microbiota is one of the densest, most dynamic, and complex microorganism
37 populations located in the body [6]. Gut microbiota act against pathogens, aid in digestion and
38 absorption, and stimulate the immune system [7,8]. If the microbiome is altered, this could result
39 in gastrointestinal diseases, such as enterocolitis, diarrhea, ulcers, anorexia, colic, and even death
40 [9,10]. *Streptococcus bovis/equinus* complex (SBEC), *Clostridium difficile*, and *Clostridium*
41 *perfringens* are bacteria found in the hindgut microbiome that are considered opportunistic due
42 to GIT issues when the immune system is compromised. SBEC is a group of human and animal
43 derived streptococci that are commensal, opportunistic pathogens, or food fermentation
44 associates [11]. *C. difficile* is commonly associated with the onset of colic in horses, but has also
45 been isolated from foals with diarrhea. *C. perfringens* causes enterocolitis in neonatal foals; in

46 addition, this species produces endotoxins that can cause diarrhea and severe damage to the
47 mucosa [9]. When compared to other mammals, little research has been conducted on the
48 microbiota in the GIT of horses [9].

49 Curcumin is the active ingredient in turmeric [*Curcuma longa*] that is not only known for
50 having anti-inflammatory properties, but also possessing antimicrobial, wound healing, and
51 antiparasitic properties [12,13]. In addition to curcumin having many biological activities, it is
52 relatively safe and well-tolerated [14]. Testing curcumin has shown effective antiparasitic
53 properties, it was an effective compound against *R. cesticillus* in birds [12], strongyles in cattle
54 [15], and fecal egg shedding in goats [16]. The indication of the safety and efficacy of curcumin
55 provided a solid basis for evaluating its antiparasitic and antimicrobial properties in riding
56 horses. We hypothesized that curcumin would decrease parasite shedding, inflammation, and
57 opportunistic bacteria found in the GIT of riding horses. The main objectives were to evaluate
58 fecal shedding of intestinal parasite ova and selected opportunistic bacteria as well as erythrocyte
59 sedimentation rate when dosing curcumin at 15 g per day for 30 days to riding horses.

60 **2. Materials and Methods**

61 Twelve horses, ten Southern Illinois University of Carbondale (SIUC) owned riding
62 horses and two privately owned riding horses were used for this study. All horses were between
63 the ages of five and twenty years old, and did not have any concurrent illnesses and/or ailments;
64 they also did not receive any medications or dewormer for 30 days prior to the commencement
65 of this research trial. The predominate breed utilized was Quarter Horse (nine), one mustang, one
66 warmblood, and one draft horse. Care and handling of animals used in this study was approved
67 by Southern Illinois University Animal Care and Use Committee (Protocol 15-041).

68 Horses were randomly assigned to one of the following treatments: 1) no curcumin
69 (CON) or 2) 15 g of 95% pure curcumin (CUR). The average age of the CON horses was 12.5
70 years old \pm 7 years, while the average age of the CUR horses was 13.5 \pm 7 years. There were 3
71 gelding and 3 mares on CON and 6 mares on CUR treatments. The CON horses received 3 - 4
72 alfalfa cubes, moistened with water, and mashed. The CUR horses received the same alfalfa cube
73 mash as CON horses but with 15 g of curcumin (Noble Elephant Supplement, San Dimas, CA)
74 mixed into the mash. The dosage of curcumin was based off the recommended dose of one
75 tablespoon, which equates to 15 g per horse [17]. Horses were gathered and samples were
76 collected at 1100 daily. Once samples were collected horses were fed either CON or CUR
77 treatment. All horses were housed on pasture and grazed *ad libitum* when not ridden. All horses
78 were ridden for an average of three hours daily for four days a week.

79 ***2.1 Analysis of Intestinal Parasites***

80 A fresh fecal sample was collected from each horse at 1100, prior to initiation of the
81 study (d 0), and then daily for 30 days. Fecal samples were collected, placed in a Whirl-Pak[®]
82 sample bag (Nasco, Fort Atkinson, WI) then placed in the refrigerator. Once all samples were
83 collected (approximately one hour later), 3 g of feces was weighed and the remainder of sample
84 was immediately frozen at -20°C for later analysis of opportunistic bacteria. Fecal parasite load
85 was determined using the modified Wisconsin sugar flotation method and the centrifugation
86 technique with a specific gravity between 1.20 to 1.33; strongyles, ascarids, and total eggs were
87 counted and recorded [18,19].

88 ***2.2 Erythrocyte Sedimentation Rate***

89 Blood samples were collected via jugular venipuncture using a 5 mL syringe with a 20-
90 gauge, 1.5-inch needle and transferred to a 5 mL vacutainer K2 Ethylenediaminetetraacetic acid
91 tube (Fischer Scientific, Franklin Lakes, NJ) for erythrocyte sedimentation rate (ESR)
92 assessment (Globe Scientific, Paramus, NJ). The westergren ESR test was performed on days 0,
93 3, 7, 10, 14, 17, 21, 24, and 28 to examine inflammation pre- and post-riding. Samples collect on
94 days 0, 7, 14, 21, and 28 were collected prior to riding. Samples collected on days 3, 10, 17, and
95 24 were collected after four days of riding in which horses were ridden for an average of three
96 hours per day.

97 ***2.3 Fecal Analysis of Opportunistic Bacteria***

98 *Growth of Bacteria*

99 Pure cultures of selected opportunistic bacteria were grown and used as standards for
100 qPCR. Trypticase soy broth (30 g/L) and yeast extract (3 g/L) medium was made for SBEC,
101 according to Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ)
102 (Germany) media recipes. *Clostridium* medium (17 g/L digest casein, 3 g/L digest soy, 5 g/L
103 NaCl, 2.5 g/L K₂Pho₄ and dextrose) was made for both *C. difficile* and *C. perfringens*, according
104 to Difco™ (Becton, Dickson and Company, Sparks, MD). Ten mL of broth was pipetted into
105 glass Hungate tubes and deoxygenated with nitrogen. Rubber stoppers and metal caps were
106 crimped on the tubes and then were autoclaved at 121°C, 15 psi, for 15 min. Hungate tubes were
107 inoculated with pelleted strains of bacteria, *C. difficile*, *C. perfringens*, and SBEC. Dense
108 bacterial samples were transferred to a new Hungate tube every three days for ten days to ensure
109 pure cultures (Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ)
110 (Germany)). These pure cultures contained the forward and reverse primers in Table 1.

111 DNA Extraction

112 DNA was extracted from the daily fecal samples collected using PowerFecal Mo Bio
113 DNA Extraction Kits (Mo Bio Laboratories, Carlsbad, CA). The pure cultures that were used as
114 standards for qPCR (Bio-Rad MyiQ Optical System Software 2.0) were extracted using
115 PowerFecal Mo Bio DNA Extraction Kits (Mo Bio Laboratories, Carlsbad, CA) and then
116 purified using UltraClean 15 DNA Purification Kits (Mo Bio Laboratories, Carlsbad, CA). All
117 DNA extractions were assessed for concentration using a Nano Drop ND-1000
118 Spectrophotometer (Wilmington, DW).

119 Real Time qPCR

120 Real-time qPCR reactions were performed in triplicate using a MyiQ thermocycler
121 (BioRad, Hercules, CA) in a total volume of 25 μ L. The reaction mixture was composed of
122 Maxima SYBR Green/ROX qPCR (Thermo Scientific, Waltham, MA), the corresponding
123 forward and reverse primers (Eurofins MWG Operon, Louisville, KY) (Table 1), extracted DNA,
124 and sterile water. The thermal cycling protocols for *C. difficile* [20], *C. perfringens* [21], and
125 SBEC [22] were utilized at an average starting concentration of DNA at 52 ng/ μ L, 20 ng/ μ L, and
126 17 ng/ μ L, respectively.

127 **2.4 Statistical Analysis**

128 All data were analyzed using the MIXED procedure of SAS (SAS 9.4 Inst., Inc., Cary,
129 NC) for repeated measures. The model included treatment, day, and treatment \times day interactions.
130 An autoregressive covariance structure (AR1 of the MIXED procedure of SAS) was determined
131 to be most appropriate based on Akaike's Information Criterion. There were no treatment \times day
132 interactions; therefore, only treatment means are reported. Comparisons of main effects were

133 determined using least square means and Fisher's protected LSD ($P \leq 0.05$) and a trend set at (P
134 ≤ 0.10).

135 **3. Results and Discussion**

136 ***3.1 Analysis of Intestinal Parasites***

137 Dosing curcumin at the recommended rate of 15 g per horse had no effect ($P \geq 0.58$) on
138 the amount of shedding of total fecal egg count, strongyles, or ascarids; however, there was a day
139 effect. Total egg counts for both CON and CUR horses decreased ($P \leq 0.05$) from d 1 to 10 then
140 increased from d 11 to 19 followed by a decrease from d 20 to 29 with a substantial increase on d
141 30. This pattern of decreasing and increasing ova was due to both strongyles ($P = 0.05$) and
142 ascarids ($P = 0.0007$) decreasing and increasing for both CON and CUR horses. The increase
143 and decrease of shed ova, approximately every ten days, would suggest the pattern of the
144 strongyles lifecycle was observed [23]; these intestinal parasites can develop from an egg to
145 larval stage 3 in a period of around one week in the optimal temperatures (8 – 38°C). The study
146 was conducted at the onset of spring when it is expected for fecal egg counts to be higher than in
147 the winter months [23, 24]. In contrast, when dosing cattle turmeric at 100 mg/ml a decrease in
148 strongyles was observed [15]. The same dose was also determined to be 100% effective against
149 adult gastrointestinal tract nematodes (roundworms) [15]. Similarly, a decrease in fecal egg
150 shedding was observed when goats were dosed 0.2% or 0.6% curcumin of the daily ration for 60
151 days when compared to d 0 [16]. In the current study, it is possible that 30 days was not a long
152 enough dosing period or that a larger dose is needed to observe the antiparasitic effects of
153 curcumin. It should also be noted that treatments in the current study were randomly assigned
154 prior to d 0 fecal egg counts and unfortunately most of the high shedding horses were randomly
155 assigned to CUR, which could have influenced the results. The results would suggest that dosing

156 curcumin for 30 days is not effective for intestinal parasite control in riding horses during the
157 spring months of the Midwest region and that further research is needed to examine effects of
158 curcumin with a longer dosing period or at an increased dose. In the current study, curcumin was
159 utilized based on previous parasite work, mentioned above, and also do to its lower cost to
160 supplement. However, the bioavailability of curcumin is noted to be minimal due to being
161 hydrophobic, low intrinsic activity, poor absorption, and high rate of metabolism and elimination
162 from the body [25]. It is possible that the results would have been different had liposomal
163 curcumin been utilized. In rats, oral administration of liposome-encapsulated curcumin showed
164 increased bioavailability of curcumin [26, 27].

165 ***3.2 Erythrocyte Sedimentation Rate Analysis***

166 There were no treatment x time interactions and treatment had no effect on ESR ($P \leq$
167 0.42); however, a day effect ($P \leq 0.001$) was observed (Table 2). ESR within the CUR horses,
168 significantly decreased ($P = 0.0006$) on d 14 and again on d 21 ($P = 0.02$) compared to d 0
169 (Figure 1). However, there was no difference ($P = 0.64$) between d 0 and d 28. There was no day
170 effect ($P = 0.29$) for ESR in CON horses. This would suggest that it would take at least 14 days
171 before curcumin could potentially decrease inflammation in riding horses. These findings are
172 similar to Farinacci et al. (2009) [28], who observed curcumin down regulating COX-2, TNF- α ,
173 and IL-6 when mares and foals were administered 4mg/kg of CURCUVET®; and on d 15, COX-
174 2 was significantly down-regulated. However, in the current study inflammation on day 28 was
175 not significantly different compared to day 0 which would further suggest that the bioavailability
176 of curcumin is minimal, as previously discussed. Moreover, factors such as an increase in
177 ambient temperature, experience of the rider and saddle fit could have contributed to a similar
178 ESR on day 28 compared to day 0.

179 3.3 Fecal Analysis of Opportunistic Bacteria

180 There was no treatment ($P = 0.34$) or day effect ($P = 0.53$) on concentration of *C.*
181 *perfringens* (Table 2). Similarly, there was no treatment effect for *C. difficile*, ($P = 0.51$) or
182 SBEC ($P = 0.69$). However, *C. difficile*, for both CON and CUR horses, had a significant day
183 effect ($P = 0.0001$) with all horses having higher concentrations on d 0 and d 1 compared to all
184 other days. Furthermore, concentrations of SBEC were affected by day ($P = 0.05$) with CON and
185 CUR horses having similar concentrations at d 0 and d 1. However, the CON horses had an
186 increase in SBEC concentration on d 9, 25, 27, and 30 when compared to d 0, while the CUR
187 horses had increased concentrations on d 9, 17, and 27 compared to day 0. The data from this
188 study would indicate that curcumin can be dosed at 15 g per day for 30 days with no adverse side
189 effects on opportunistic bacteria concentrations, however, more research is needed to evaluate
190 the antimicrobial properties of curcumin when dosed to horses over a longer duration and at
191 varying dosages.

192 5. Conclusions

193 The antiparasitic and antimicrobial properties of curcumin were not observed when 15 g
194 of curcumin was orally dosed to twelve riding horses for 30 days. The inability for curcumin to
195 decrease the parasite shedding load would suggest that curcumin will need to be dosed for longer
196 periods of time or at higher dosages, if utilizing for intestinal parasite control. However, it is
197 possible that curcumin can decrease inflammation after 14 days of administration. More research
198 is needed to further evaluate the benefits of supplementing curcumin to horses, especially for
199 intestinal parasitic control.

200 Funding: This work was partially supported by the Illinois Equine Industry Research and
201 Promotion Board.

202 **References**

- 203 [1] Nielson, M. K., J. Monrad, and S. N. Olsen. 2006. Prescription-only anthelmintics - A
204 questionnaire survey of strategies for surveillance and control of equine strongyles in
205 Denmark. *Vet. Parasitol.* 135:47-55.
- 206 [2] Evans, J. W., A. Borton, H. Hintz, L. Dale Van Vleck. 1977. *The Horse*. 2nd Ed. W. H.
207 Freeman and Company. New York. P. 622-630.
- 208 [3] Clayton, H. M. 1986. Ascarids: Recent Advances. *Vet. Clin. North Am. Equine Pract.* 2(2):
209 313-328.
- 210 [4] Van Doorn, D. C. K., M. Eysker, F. N. J. Kooyman, J. A. Wagenaar, and H. W. Ploeger.
211 2012. Searching for ivermectin resistance in Dutch horses. *Vet. Parasitol.* 185:355-358.
- 212 [5] Bland, S. 2015. Canine osteoarthritis and treatments: A review. *Vet. Sci. Dev.* 5:2-25.
- 213 [6] Costa, M. C., L. G. Arroyo, E. Allen-Vercoe, H. R. Stampfli, P. T. Kim, A. Sturgeon, and J.
214 S. Weese. 2012. Comparison of fecal microbiota of healthy horses and horses with colitis
215 by high throughput sequencing of the V3-V5 region of the 16s rRNA gene. *PLoS ONE*.
216 7(7):1-11.
- 217 [7] Marteau, P., P. Pochart, J. Dore, C. Bera-Maillet, A. Bernalier, and G. Cortheir. 2001.
218 Comparative study of bacterial groups within the human cecal and fecal microbiota.
219 *Appl. Environ. Microbiol.* 67(10).
- 220 [8] Costa, M. C. and S. Weese. 2012. The equine intestinal microbiome. *Anim. Health Res. Rev.*
221 13(1):121-128.
- 222 [9] Dicks, L. M. T., M. Botha, E. Dicks, and M. Botes. 2014. The equine gastro-intestinal tract:
223 An overview of the microbiota, disease and treatment. *Livest. Sci.* 160:69-81.

- 224 [10] Bland, S. 2016. Therapeutic and safety evaluation of curcumin's antimicrobial and anti-
225 inflammatory properties of canine and equine. PhD Diss. Southern Illinois Univ.,
226 Carbondale.
- 227 [11] Jans, C., L. Meile, C. Lacroix, and M. Stevens. 2015. Genomics, evolution, and molecular
228 epidemiology of the *Streptococcus bovis*/*Streptococcus equinus* complex (SBSEC).
229 Infect. Genet. Evol. 33:419-436.
- 230 [12] El-Bahy, N. and E. Bazh. 2015. Anthelmintic activity of ginger, curcumin, and praziquantel
231 against *Raillietina cesticillus* (in vitro and in vivo). J. Parasitol. Res. 114:2427-2434.
- 232 [13] Ghosh, S., S. Banerjee, and P. Sil. 2015. The beneficial role of curcumin on inflammation,
233 diabetes, and neurodegenerative disease: A recent update. Food Chem. Toxicol. 83:111-
234 124.
- 235 [14] Collison, S. 2014. Curcumin and inflammatory diseases. Today's Dietician. 16(9):56-63.
- 236 [15] Amin, M. R., M. Mostofa, M. E. Hoque, and M. A. Sayed. 2009. In vitro anthelmintic
237 efficacy of some indigenous medicinal plants against gastrointestinal nematodes of cattle.
238 J. Bangladesh Agril. Univ. 7(1):57-61.
- 239 [16] Afrin, F., M. M. R. Chowdhury, S. S. Saha, Md. Aatur Rahman, Md. Ali Asgar, and Md.
240 Kamrul Islam. 2016. Efficacy of curcumin against gastro-intestinal parasites in goat. Eur.
241 J. Pharm. Med. Research. 3(8):158-165.
- 242 [17] Getty, J. M. 2016. Consider Curcumin for Joint Inflammation.
243 <http://gettyequinenutrition.biz/Library/ConsiderCurcuminforJointInflammation.htm>.
244 (Accessed 13 June 13 2017).
- 245 [18] Dryden, M.W., P. A. Payne, R. Ridley, and V. Smith. 2005. Comparison of common fecal
246 flotation techniques for the recovery of parasite eggs and oocysts. Vet. Ther. 6(1):15-28.

- 247 [19] University of Pennsylvania School of Veterinary Medicine. 2006.
248 <http://cal.vet.upenn.edu/projects/parasit06/website/modWis.htm>. (Accessed 10 Feb.
249 2016).
- 250 [20] Avbersek, J., M. Cotman, and M. Ocepek. 2011. Detection of *Clostridium difficile* in
251 animals: Comparison of real-time PCR assays with the culture method. *J. Med.*
252 *Microbiol.* 60:1119-1125.
- 253 [21] Karpowicz, E., A. Novinscak, F. Bärlocher, and M. Filion. 2009. qPCR quantification and
254 Genetic characterization of *Clostridium perfringens* populations in biosolids composted
255 for 2 years. *J. Appli. Microbiol.* 108:571-581.
- 256 [22] Hastie, P. M., K. Mitchell, and J. M. D. Murray. 2008. Semi-quantitative analysis of
257 *ruminococcus flavefaciens*, *fibrobacter succinogenes* and *stretococcus bovis* in the equine
258 large intestine using real-time polymerase chain reaction. *British J. Nutr.* 100:561-568.
- 259 [23] Johnstone, C. 2000. Parasites and parasitic diseases of domestic animals.
260 http://cal.vet.upenn.edu/projects/merial/Strongls/strong_8a1.htm. (Accessed 8 Oct 2016).
- 261 [24] Thomas, H. S. 2014. Pasture management for parasite control.
262 <http://www.thehorse.com/articles/33480/pasture-management-for-parasite-control>.
263 (Accessed 29 Sept 2016).
- 264 [25] Anard, P., A. B. Kunnumakkara, R. A. Newman, and B. B. Aggarwal. 2009. Bioavailability
265 of curcumin: Problem and promises. *Mol. Pharm.* 4(6):807-818.
- 266 [26] Takahashi, M., S. Uechi, K. Takara, Y. Asikin, and K. Wada. 2009. Evaluation of an oral
267 carrier system in rats: bioavailability and antioxidant properties of liposome-encapsulated
268 curcumin. *J. Agric. Food. Chem.* 57:9141–9146.

- 269 [27] Prasad, S., A. K. Tyagi, and B. B. Aggarwal. 2014. Recent developments in delivery,
270 bioavailability, absorption and metabolism of curcumin: The golden pigment from golden
271 spice. *Cancer Res. Treat.* 46(1):2-18.
- 272 [28] Farinacci, M., B. Gaspardo, M. Colitti, and B. Stefanon. 2009. Dietary administration of
273 Curcumin modifies transcriptional profile of genes involved in inflammatory cascade in
274 horse leukocytes. *Ital. J. Anim. Sci.* 8(Suppl. 2):84-86.
- 275

276 Table 1. Forward and Reverse Primers used for real time PCR
277

Strains	Forward Primers (5'-3')	Reverse Primers (5'-3')
SBEC ¹	GCCTACATGAAGTCGGAATCG	TACAAGGCCCGGGAACGTA
<i>C. difficile</i> ²	CAAGTTGAGCGATTTACTTCGGTAA	CTAATCAGACGCGGGTCCAT
<i>C. perfringens</i> ³	AAATGTAACAGCAGGGGCA	TGAAATTGCAGCAACTCTAGC

278 ¹*Streptococcus bovis/equinus* complex [22]

279 ²*Clostridium difficile* [20]

280 ³*Clostridium perfringens* [21]

281

282 Table 2. Effects of 15 g of 500 mg/g of 95% curcumin dosed once daily to riding horses on fecal
 283 parasite load (number of eggs/3 grams of feces), erythrocyte sedimentation rate, and fecal
 284 opportunistic bacteria.

285
 286


Parameters	Treatments ¹			P-Value	
	CON	CUR	SEM	TRT	DAY
Total Egg	127	156	37.8	0.58	0.05
Strongyles	123	153	37.7	0.58	0.05
Ascarids	3.22	3.11	0.24	0.74	0.0007
ESR ² , mm/hr	89.6	94.5	4.21	0.42	0.001
SBEC ³ , ng/ μ L	1.93	1.66	0.46	0.69	0.04
<i>C. difficile</i> , ng/ μ L	4108	8080	4136	0.51	0.0001
<i>C. perfringens</i> , ng/ μ L	0.01	0.0001	0.01	0.34	0.53

287 ¹ Treatments: CON= control; CUR= curcumin dosed at 15g/day

288 ² Erythrocyte Sedimentation Rate

289 ³ *Streptococcus bovis/equinus* complex

290 Figure 1. Effect of curcumin on Erythrocyte Sedimentation Rate over time (days).



291