

*Ethnobotanical Leaflets 12: 44-55, 2008.*

## **Ethnomedicinal Survey of Botanicals Used in Treating Sexually Transmitted Diseases in Ekiti State, Nigeria**

**J. Kayode<sup>1,3</sup> and G.M. Kayode<sup>2</sup>**

<sup>1</sup>Department of Plant Science, University of Ado-Ekiti, Ado-Ekiti, Nigeria.

<sup>2</sup>Department of Adult Education, University of Ado-Ekiti, Ado-Ekiti, Nigeria.

<sup>3</sup>Author to whom correspondence should be addressed. E-mail: josmodkay@yahoo.com

**Issued 29 January 2008**

### **ABSTRACT**

This paper focuses on the botanicals used in the treatment of sexually transmitted diseases in Ekiti State, Nigeria. The authors identified 49 plants belonging to 30 angiosperm families that were being used by the natives of Ekiti State for the cure of sexually transmitted diseases. Most of the wild species were rare or not very abundantly distributed in nature. The proportion of cultivated to wild species, however, was relatively low. Species that were under cultivation by the natives were being grown largely for reasons other than medicinal value. The methods of extraction were mostly predatory and annihilative. Considerable proportions of the identified botanicals were obtained from the forest. The need for the conservation of most of the species cannot be over emphasized. Strategies towards the attainment of this goal were proposed.

### **INTRODUCTION**

There has been an increase in worldwide realization of the use of medicinal plants in various traditional health systems of developing countries. For example, recent estimates by the World Health Organization (WHO) revealed that about 80% of the population in Africa relies on traditional medicine of which the botanicals constituted greater components. It is estimated that about 30,000 botanical species are now recorded for their medicinal properties.

These botanicals had over the years been subjected to wide and unsustainable use (Kayode 2002). They are now diminishing at an alarming rate. Although studies on the ethnomedicinal utilization of botanicals abound in Nigeria, these studies were conducted on scattered basis usually based on the ethnic composition of the country. Presently, a gross

dearth of documentation abounds on the ethnomedicinal utilization of botanicals among the Ekiti, a distinct Yoruba tribe that constitutes over 98% of the 1.6 million inhabitants (EKSG 1997) of Ekiti state. At present sexually transmitted diseases (STD) are perhaps the most devastating diseases in the state. The diseases include Gonorrhoea, Trichomoniasis, Chlamydial infection, Syphilis and, more recently, the Acquired Immune Deficiency Syndrome (AIDS).

The aim of this study therefore is to identify botanicals used traditionally in the management of sexually transmitted diseases in Ekiti State, and propose sustainable strategies for the conservation of these species.

## **MATERIALS AND METHODS**

The study was conducted in Ekiti State, which is situated in the southwestern part of Nigeria. Ekiti has a land area of about 7000km<sup>2</sup>, and is situated between latitude 7<sup>0</sup>25' and 8<sup>0</sup>20' North and longitude 5<sup>0</sup>00' and 6<sup>0</sup>00' East. The state has a population of about 1.6 million (EKSG 1997), 75% of whom are farmers who live in rural areas. There are two climatic seasons in the state, a dry season from November to February and a rainy season from March to October. The annual rainfall is about 1150mm (Kayode and Faluyi 1994). According to Smith and Montgomery (1962), the soil is overlying metamorphic rocks of basement complex, which shows greater variations in size and mineral composition.

## **METHODS**

A combination of social surveys and direct field observation (Kayode 2002) was used in the study. The entire state was divided into three zones based on the existing political delineation. These zones are Ekiti Central, Ekiti South and Ekiti North. In each zone, three major markets were selected, the major criterion for selection being the level of patronage by residents from both rural and urban centers in the zone. In each of these markets, vendors of medicinal plant species were identified and interviewed with the aid of a semi-structured matrix. The interviews focused on plant species used in curing sexually transmitted diseases.

The botanical species were identified by the vendors; the part(s) of the species used and methods of application during utilization were identified and recorded. Voucher specimens of the species were obtained and taken to the herbarium of the Department of Plant Science, University of Ado-Ekiti, for scientific identification and preservation.

The abundance of the species identified was determined in the study area. For this purpose, five rural communities, which were far from urban influence, were selected in each zone. In each of these communities, the abundance of each of the identified species was determined within 5 kilometers radius from the center of each community using the abundance

scale defined by Kayode (1999) as follows: Rare when the number of the individual species found available within the defined area was less than 5; Occasional when between 5 and 10 individuals were found; Frequent when between 11 and 30 individuals were found; Abundant when between 31 and 100 individuals were found; And, very abundant when more than 100 individuals were found. Also in each community, ten elderly respondents were randomly selected and interviewed on their knowledge of the utilization of the identified botanical species.

Also in each zone, five key informants who were knowledgeable in the use of botanical species were identified and interviewed. These included herbalists and community development officers. Secondary information on the active principles present in the identified species was obtained from the literature, especially Oliver (1960), Gbile (1986) and Gill (1992).

## RESULTS AND DISCUSSION

The following 49 plant species belonging to 30 families were identified as being used for curing sexually transmitted diseases in the study area:

### Alliaceae

#### *Allium cepa*

Local Name: Alubasa

Parts used: Leaves, bulb

Major source: Market

Abundance at source: Very abundant

Active Principle: Riboflavin, n-propyl disulphide

#### *Allium ascalonicum*

Local Name: Alubasa

Parts used: Whole plant

Major source: Household farms

Abundance at source: Rare

Active Principle: Riboflavin

### Amaranthaceae

#### *Amaranthus spinosus*

Local Name: Tete elegun

Parts used: Leaves, stem

Major source: Farms

Abundance at source: Abundant

Active Principle: Tannins, saponin, hydrocyanic acid

***Cyathula prostrata***

Local Name: Shawere pepe  
Parts used: Leaves, stems  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Tannins, saponin

**Annonaceae*****Haxelobus monopetalus***

Local Name: Lapawe  
Parts used: Roots, Stems, Leaves  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Saponin, inulin, essential oil

**Apocynaceae*****Landolphia owariensis***

Local Name: Ibo-akitipa  
Parts used: Leaves, roots, stem bark, seeds  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Saponin, tannins

**Asclepiadaceae*****Secamone afzelii***

Local Name: Alu  
Parts used: Stems, Leaves  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Alkaloids

**Bignoniaceae*****Kigelia africana***

Local Name: Pandoro  
Parts used: Leaves, roots, stem bark, fruit  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Saponin, tannins, inulins, B-amyrin (Msonths 1986)

***Sterospermum kunthianum***

Local Name: Akoko-igbo  
Parts used: Leaves, roots, stem bark, fruits  
Major source: Forest

Abundance at source: Rare  
Active Principle: Tannins, saponin

## Burseraceae

### *Canarium schweifuthii*

Local Name: Origbo  
Parts used: Stem bark  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Saponin, tannins, resin, amyryl, limonene, phellandrina (Gill 1992).

## Cactaceae

### *Opuntia dillenii*

Local Name: Oro  
Parts used: Stem, roots  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Tannins, saponin

## Caesalpiaceae

### *Azelia africana*

Local Name: Apa  
Parts used: Root  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Alkaloid, Tannins

### *Cassia podocarpa*

Local Name: Asunrin  
Parts used: Leaves  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Anthraquinones

### *Macrolobium macrophyllum*

Local Name: Aba  
Parts used: Stem bark  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Tannins, saponin

### *Mezoneuran benthamianum*

Local Name: Ajuju  
Parts used: Leaves, stem, roots  
Major source: Forest

Abundance at source: Rare  
Active Principle: Saponins, mucilage

## Caricaceae

### *Carica papaya*

Local Name: Ibepe  
Parts used: Leaves, fruits, roots  
Major source: Household farms  
Abundance at source: Abundant  
Active Principle: Carpaine, saponin, tannins, nicotinic acid, tocopherol, papain

## Colchicaceae

### *Gloriosa superba*

Local Name: Ewe-aje  
Parts used: Leaves  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Superbin, colchicin, gloriosine, gloriosol, phytosterils, stigmasterin

## Combretaceae

### *Terminalia catapa*

Local Name: Odan  
Parts used: Stem bark  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Tannins

### *Terminalia glaucescens*

Local Name: Odan  
Parts used: Stem bark, roots  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Alkaloids, tannins

## Connaraceae

### *Cnestis ferruginea*

Local Name: Omu-aje  
Parts used: Leaves, roots, fruits, seeds  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Glycosidea

## Dilleniaceae

### *Tetracera alnifolia*

Local Name: Opon  
Parts used: Leaves, roots  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Glycoside – syringin, tannis

## **Euphorbiaceae**

### ***Alchornea cordifolia***

Local Name: Ipa  
Parts used: Leaves, stem bark, fruits, roots  
Major source: Forest  
Abundance at source: Frequent  
Active Principle: Inulin, tannins, alchornin, alkaloid

### ***Alchornea laxiflora***

Local Name: Pepe  
Parts used: Stem  
Major source: Forest  
Abundance at source: Frequent  
Active Principle: Alkaloid

### ***Manihot esculenta***

Local Name: Ege  
Parts used: Leaves, tubes  
Major source: Household farm  
Abundance at source: Very abundant  
Active Principle: Alkaloid, saponins, tannins

### ***Phyllanthus niruri***

Local Name: Asasa  
Parts used: Leaves, stem, roots  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Saponins, phyllanthin, hypophlenthin

## **Lamiaceae**

### ***Ocimum basilicum***

Local Name: Efinrin-wewe  
Parts used: Leaves, stem, roots  
Major source: Household farms  
Abundance at source: Frequent  
Active Principle: Essential oils, methylcinnamate, thymol, terpenes

## **Malvaceae**

### ***Abuilon mauritianum***

Local Name: Furu  
 Parts used: Leaves, roots  
 Major source: Forest  
 Abundance at source: Rare  
 Active Principle: Tannins, saponin

### ***Hibiscus esculentus***

Local Name: Ila  
 Parts used: Fruits, seeds  
 Major source: Household farms  
 Abundance at source: Very abundant  
 Active Principle: Essential oils-farnesol

### ***Sida cordifolia***

Local Name: Iseketu pupa  
 Parts used: Leaves, roots  
 Major source: Forest  
 Abundance at source: Abundant  
 Active Principle: Alkaloid-ephedrine

## **Meliaceae**

### ***Trichilia prieuriana***

Local Name: Awe  
 Parts used: Roots  
 Major source: Forest  
 Abundance at source: Rare  
 Active Principle: Tannins, saponin

## **Moraceae**

### ***Ficus asperifolia***

Local Name: Eripin  
 Parts used: Leaves, stem bark, roots  
 Major source: Forest  
 Abundance at source: Occasional  
 Active Principle: Tannins

### ***Ficus capensis***

Local Name: Opoto  
 Parts used: Roots  
 Major source: Forest  
 Abundance at source: Occasional  
 Active Principle: Tannins

## **Mimosaceae**

### ***Tetrapluera tetreptera***



Local Name: Aridan  
Parts used: Stem bark  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Saponins-Aridanu, essential oils, scopoletin

## **Papaveraceae**

### ***Argemone mexicana***

Local Name: Egunarigbo  
Parts used: Roots  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Alkaloids-berberine, protopine

## **Passifloraceae**

### ***Adenia lobata***

Local Name: Dodo  
Parts used: Leaves, stem  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Flaviroid

## **Papilionaceae**

### ***Desmodium adecendens***

Local Name: Epa-ile  
Parts used: Leaves  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Tannis

### ***Erythrina senegalensis***

Local Name: Ologun-sese  
Parts used: Leaves, stem bark, seeds  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Alkaloid – hypaphorine

## **Polygalaceae**

### ***Securidaca longepedunculata***

Local Name: Ofodo  
Parts used: Leaves  
Major source: Forest  
Abundance at source: Rare  
Active Principle: Saponin-glycosides, tannins, valerianate methylsalicylate

## Rutaceae

### *Citrus aurantifolia*

Local Name: Osan-wewe  
 Parts used: Stem and root barks  
 Major source: Household farms  
 Abundance at source: Abundant  
 Active Principle: Essential oils

### *Fagara macrocarpa*

Local Name: Ata igbo  
 Parts used: Stem and root barks  
 Major source: Forest  
 Abundance at source: Rare  
 Active Principle: Alkaloids-xanthofegarine, erythrofagarin, fagaramide, f-methoxy-dihydroneitidine

### *Fagara zanthoxyloides*

Local Name: Ata  
 Parts used: Root and stem barks  
 Major source: Forest  
 Abundance at source: Rare

Active Principle: Alkaloi, p-hydroxybenzoic acid, 2-hydroxymethyl benzoic acid, vanillic acid, inulin,  
 Saponin

## Sapindaceae

### *Bligha sapida*

Local Name: Ishin  
 Parts used: Stem bark  
 Major source: Household farms  
 Abundance at source: Abundant  
 Active Principle: Saponin, hypoglycin, tannins, steroidal alkaloid

### *Cardiospermum halicacabium*

Local Name: Shaworo  
 Parts used: Roots  
 Major source: Forest  
 Abundance at source: Rare  
 Active Principle: Saponins

## Scrophulariaceae

### *Scoparia dulcis*

Local Name: Aya  
 Parts used: Roots

Major source: Forest  
Abundance at source: Rare  
Active Principle: Alkaloids, inulin, saponins, tannins

## **Solanaceae**

### ***Capsicum frutescens***

Local Name: Ata wewe  
Parts used: Fruits  
Major source: Household farms  
Abundance at source: Very abundant  
Active Principle: Capsaicin, oil, ascorbic acid

### ***Solanum nigrum***

Local Name: Odu  
Parts used: Leaf  
Major source: Household farms  
Abundance at source: Abundant  
Active Principle: Alkaloid-solanine, solamarine, scopolin , scopoletin, aesculin,, isoscopoletin, demisine, solarmagine, solasodabiro, tomatine, solauricine, solangustine

### ***Solanum vervascifolium***

Local Name: Ikan  
Parts used: Leaves, fruits, roots  
Major source: Farms  
Abundance at source: Abundant  
Active Principle: Alkaloid-solanine, saponins

## **Tiliaceae**

### ***Glyphaea brevis***

Local Name: Atori  
Parts used: Leaves  
Major source: Forest  
Abundance at source: Occasional  
Active Principle: Tannins, saponin

## **Verbenaceae**

### ***Gmelina arborea***

Local Name: Melaina  
Parts used: Leaves  
Major source: Government Reserve Forest  
Abundance at source: Very abundant  
Active Principle: Tannins

Most of these species were rare in abundance and the proportion of the cultivated species was relatively low. Species cultivated were meant for other purposes other than their medicinal value. The methods of extraction were mostly predatory and annihilative. Such methods, as previously observed by Homman (1994), Kayode and Ogunleye (2008), entailed the destruction of source(s) in such a rate that the regeneration is slower than the rate of extraction. Thus, predatory and annihilation usually results in increasing scarcity of species. Although some of the species were extracted by non-predatory and gathering methods, yet collections were observed to be by pulling or cutting of the branches thus making such collection destructive. Field observations revealed that collections were done indiscriminately without any consideration for size and age, thus resulting in species depletion. Also, the lower-altitude harvesting by a larger number of households in the study area due to the less vegetation cover per inhabitants may be detrimental to the survival of these species.

Considerable proportions of the identified botanicals were obtained from the forest. Thus the increasing conversion of valuable natural environments in the study area to monoculture plantations of exotic timber and agriculture will likely lead to a continued erosion of botanical diversity in the study area. Thus, some of the presently rare species require urgent domestication while in-situ and ex-situ conservation methods should be embarked upon. These, according to Shinwari and Khan (2000) will require the protection of plant species in their natural habitats followed by ex-situ devices by growing the rare species and subsequently re-introducing them into their natural environment. The domestication of most of the botanicals identified is now desirable, further research activities are still required to develop deep understanding of the life cycles, pollination, and dispersal mechanisms in most of the botanicals. The populace should be enlightened on the dangers in the loss of biological diversity. Kayode (2006) had also advocated the need to accommodate the indigenous farmers in both planning and execution of conservation activities. This strategy is still relevant in the study area.

## REFERENCES

EKSG 1997. First Anniversary Celebration of Ekiti State. Government Press, Ado-Ekiti, Nigeria. 22pp.

Gbile, Z. O. 1986. Ethnobotany, taxonomy and conservation of medicinal plants. Pp. 13-29. In: Sofowora, A. (Ed.). The state of medicinal plants research in Nigeria. University of Ibadan Press, Ibadan, Nigeria.

- Gill, L. S. 1992. Ethnomedicinal uses of plants in Nigeria. Uniben Press, Benin-City, Nigeria. 276pp.
- Homman, A. K. O. 1994. Plant extractivism in the Amazon: Limitations and possibilities. In: Extractivism on Regional Development. Ctusener-God, M. and Sachs, I. (Eds.). MAB Digest 18, UNESCO, Paris. pp. 34-57.
- Kayode, J and Faluyi, M. A 1994. Studies on self and cross compatibility studies on soybeans (*Glycine max*) in a tropical environment. Nigerian Journal of Botany 7:55- 61.
- Kayode, J. (2006). Conservation of indigenous medicinal botanicals in Ekiti State, Nigeria. Journal of Zhejiang University SCIENCE-B 7 (9): 713 -718.
- Kayode, J. and Ogunleye, T. (2008). Checklist and Status of Plant Species Used as Spices in Kaduna State of Nigeria. Research Journal of Botany 3 (1), 35-40
- Oliver, B. (1960). Medicinal plants in Nigeria. University of Ibadan Press, Ibadan, Nigeria. 139pp.
- Shinwari, M. I. and Khan, M. A. 2000. Folk use of medicinal herbs of Magalla Hills National Park, Islamabad. Journal of Ethnopharmacology, 69: 45-56.
- Smith, A. J. and Montgomery R.T. 1962. *Soil and land use in central western Nigeria*. The Government Press, Ibadan, Nigeria.