

2-2007

Hartman-Grobman Theorems along Hyperbolic Stationary Trajectories

Edson A. Coayla-Teran
Universidade Federal da Bahia

Salah-Eldin A. Mohammed
Southern Illinois University Carbondale, salah@sfde.math.siu.edu

Paulo Régis C. Ruffino
Universidade Estadual de Campinas

Follow this and additional works at: http://opensiuc.lib.siu.edu/math_articles

 Part of the [Dynamical Systems Commons](#), and the [Ordinary Differential Equations and Applied Dynamics Commons](#)

This is a pre-copy-editing, author-produced PDF of an article accepted for publication in *Discrete and Continuous Dynamical Systems - Series A* following peer review. The definitive publisher-authenticated version "Hartman-Grobman theorems along hyperbolic stationary trajectories," *Discrete and Continuous Dynamical Systems - Series A*, 17(2), 281-292 is available online at:
<http://aimsciences.org/journals/dcdsA/online.jsp>

Recommended Citation

Coayla-Teran, Edson A., Mohammed, Salah-Eldin A. and Ruffino, Paulo Régis C. "Hartman-Grobman Theorems along Hyperbolic Stationary Trajectories." (Feb 2007).

This Article is brought to you for free and open access by the Department of Mathematics at OpenSIUC. It has been accepted for inclusion in Articles and Preprints by an authorized administrator of OpenSIUC. For more information, please contact opensiuc@lib.siu.edu.

Hartman-Grobman theorems along hyperbolic stationary trajectories

Edson A. Coayla-Teran

*Instituto de Matemática - UFBA. Av. Ademar de Barros s/n,
40170-110 Salvador-BA, Brazil. e-mail: eact@im.ufba.br.*

Salah-Eldin A. Mohammed¹

*Department of Mathematics, Southern Illinois University at Carbondale,
Carbondale, Illinois 62901, e-mail: salah@sfde.math.siu.edu*

and

Paulo Régis C. Ruffino²

*Departamento de Matemática, Universidade Estadual de Campinas,
13.081-970 - Campinas - SP, Brazil. e-mail: ruffino@ime.unicamp.br*

Abstract

We extend the Hartman-Grobman theorems on discrete random dynamical systems (RDS), proved in [7], in two directions: For continuous RDS and for hyperbolic stationary trajectories. In this last case there exists a conjugacy between traveling neighbourhoods of trajectories and neighbourhoods of the origin in the corresponding tangent bundle. We present applications to deterministic dynamical systems.

Key words: Random dynamical systems, Hartman-Grobman theorems, hyperbolic stationary trajectories.

MSC2000 subject classification: 93E99, 37D40, 34A34, (37H15).

¹Project supported by FAPESP grant no. 02/01095-0.

²Research partially supported by CNPq grant n° 301112/2003 – 7.

1 Introduction

The celebrated Hartman-Grobman theorem (HGT, for short) plays a fundamental role in the theory of dynamical systems. Essentially, among other features, it allows one to make topological classification of the dynamics in a neighbourhood of hyperbolic fixed points. This classification is based on the existence of a conjugacy of the local dynamics with the linearized system at a hyperbolic fixed point. For the original papers we mention Hartman [10] and [11], and Grobman [9].

One of the first results concerning HGT in random dynamical systems (RDS, for short) is due to Wanner [19] for discrete systems. His argument was based on random difference equation, such that the construction was made ω -wise. His proof is completed by showing that the choice of random homeomorphisms is, in fact, measurable. In our intrinsic approach, Coayla-Teran and Ruffino [7], we have looked for the conjugacy in an appropriate Banach space of random homeomorphisms. The arguments in there correspond to an appropriate extension of the deterministic arguments, with the state space enlarged by the probability space. Hence the norms and other constants were composed with $L^1(\Omega)$ norm.

Here, we extend the Hartman-Grobman theorems on discrete RDS in two directions: For continuous (hyperbolic) RDS and for hyperbolic stationary trajectories. In this last case we mean that there exists a conjugacy between traveling neighbourhoods of the trajectory and neighbourhoods of the origin in the corresponding tangent bundle, see figure 1 in Section 4.

The proofs of the main results in this article are applications of the local discrete random HGT, analogous to deterministic case, where the proof of the continuous version uses the discrete version, see Palis and de Melo [14]).

The article is organized as follows: In Section 2 we present preliminaries results; mainly the HGT for hyperbolic random fixed point, Theorem 2.1. This result was originally proved in [7] and as we said before, this is a key result to the proof of the other theorems in this article. In Section 3 we present the first extension: we lift the fixed point hypothesis and consider hyperbolic stationary orbits on discrete systems. Section 4 starts extending the HGT to continuous RDS, still in this section we show the continuous version for stationary trajectories. Finally, in Section 5 we present direct applications of the random versions of HGT to chaotic deterministic dynamical systems.

2 Definitions and preliminary results

In this section we present the main technical tools we shall use in the proofs of next sections. It can be skipped if the reader is already familiarized with random dynamical systems (RDS) in terms of cocycle theory, as in L. Arnold [1].

2.1 Random norm and stationary trajectory

To set up the notation let (θ_t) be a group of ergodic transformations on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, where $t \in \mathbb{T}$, with $\mathbb{T} = \mathbb{Z}$ or \mathbb{R} . A continuous (perfect) cocycle $\varphi(t, \omega)$ on \mathbb{R}^n over θ is a map over the space of local diffeomorphisms $\text{Dif}_{loc}(\mathbb{R}^n)$ denoted by $\varphi : \Omega \times \mathbb{T} \rightarrow \text{Dif}_{loc}(\mathbb{R}^n)$ such that for all $\omega \in \Omega$:

- (i) $\varphi(0, \omega) = Id$;
- (ii) $\varphi(t, \omega)$ is continuous on t ;
- (iii) it has the cocycle property:

$$\varphi(t + s, \omega) = \varphi(t, \theta_s(\omega)) \circ \varphi(s, \omega).$$

We deal with perfect cocycles once for every crude cocycle there exists a perfect cocycle such that they are indistinguishable, see L. Arnold and M. Scheutzow [2] or L. Arnold [1]. This cocycle generates the following random system on \mathbb{R}^n :

$$x_t = \varphi(t, \omega)x_0. \tag{1}$$

The concept of a cocycle over a measurable transformation on a probability space generalizes many interesting (random or not) dynamical systems, including those which are generated by random equations and stochastic differential equations, see Arnold [1].

Given a (locally) invertible cocycle, if

$$\sup_{-1 \leq t \leq 1} \log^+ \|D\varphi_t\| \quad \text{and} \quad \sup_{-1 \leq t \leq 1} \log^+ \|D\varphi_t^{-1}\|$$

are in $L^1(\Omega, \mathbb{P})$, then the multiplicative (Osseledec) ergodic theorem (MET) establishes a random linear algebra for RDS, see e.g. Arnold [1], Ruelle [15], Katok and Hasselblatt [12], among others. Moreover, this random linear algebra allows one to introduce a measurable random norm $\|x\|_\omega^2 = \langle x, x \rangle_\omega$, also known as Lyapunov norm (Katok and Hasselblatt [12]). For the definition

of this norm and a survey of its properties we would suggest an interested reader to see [1, Thm. 3.7.4] and references therein. We only recall that the random norm is such that the exponential behaviour of the linearized cocycle $D\varphi_t$ imitates the exponential behaviour of a deterministic linear systems with respect to the Euclidean norm, up to an exponential (“small”) correction term. More precisely: if Λ is the maximum of the modulus of the Lyapunov spectrum and $a > 0$ is smaller than the minimum of the modulus of these exponents, then,

$$e^{-(\Lambda+a)|t|} \|x\|_\omega \leq \|\Phi(t, \omega)x\|_{\theta_t\omega} \leq e^{(\Lambda+a)|t|} \|x\|_\omega \quad \text{for } t \in \mathbb{T},$$

For each $\varepsilon > 0$ there exists a random variable $B_\varepsilon(\cdot) : \Omega \rightarrow [1, +\infty)$ which provides an equivalence between the random norm and the Euclidean norm, i.e. for all $x \in \mathbb{R}^d$:

$$\frac{1}{B_\varepsilon(\omega)} \|x\| \leq \|x\|_\omega \leq B_\varepsilon(\omega) \|x\| \quad (2)$$

Moreover, we have that $B_\varepsilon(\theta_t\omega) \leq B_\varepsilon(\omega)e^{\varepsilon|t|}$, see [1] or [5].

We define the main spaces which we are going to work with, see also [7].

- a) $\text{Homeo}(\Omega, \mathbb{R}^m)$ denotes the space of random homeomorphisms given by measurables $h(\cdot, \cdot) : \Omega \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ such that for each $\omega \in \Omega$, $h(\omega, \cdot) : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a homeomorphism.
- b) $C_b(\Omega, \mathbb{R}^m)$ denotes the space of random bounded continuous maps $u(\omega, \cdot)$ such that:

$$\|u\|_{C_b(\Omega, \mathbb{R}^m)} = \mathbb{E} \left[\sup_{x \in \mathbb{R}^m} \|u(\omega, x)\|_\omega \right] < +\infty.$$

- c) $C_{0,b}(\Omega, \mathbb{R}^m) \subset C_b(\Omega, \mathbb{R}^m)$ denotes the subspace of random bounded continuous maps which fix the origin, i.e. $u \in C_{0,b}(\Omega, \mathbb{R}^m)$ if $u(\omega, 0) = 0$ a.s.. We shall denote the norm in $C_b(\Omega, \mathbb{R}^m)$ restricted to this subspace by $\|\cdot\|_{C_{0,b}(\Omega, \mathbb{R}^m)}$.

The spaces $C_{0,b}(\Omega, \mathbb{R}^m)$ and $C_b(\Omega, \mathbb{R}^m)$ with the norm defined above are Banach spaces, see [7, Prop. 2.2].

Let $\varphi(\omega, t, \cdot)$ be a cocycle over a family of ergodic transformations $\theta_t : \Omega \rightarrow \Omega$ on a complete probability space $(\Omega, \mathcal{F}, \mathbf{P})$, with $t \in \mathbf{T}$, where $\mathbf{T} = \mathbb{Z}$

or \mathbb{R} . We say that an \mathcal{F} -measurable random variable $Y : \Omega \rightarrow \mathbf{R}^d$ is a *stationary trajectory* of $\varphi(t, \omega, \cdot)$ if

$$\varphi(t, \omega, Y(\omega)) = Y(\theta_t \omega)$$

for all $t \in \mathbf{T}$ and every $\omega \in \Omega$. We say that a stationary trajectory is hyperbolic if the Lyapunov spectrum along its trajectory does not contain zero. For examples, see Section 3, where we show that support of invariant measures are stationary trajectories.

For discrete systems, frequently one finds the notation $\varphi(\omega, \cdot)$ and $\theta^k(\omega)$, with $k \in \mathbf{Z}$, more natural than $\varphi(1, \omega, \cdot)$ and $\theta_k(\omega)$, respectively. Hence, $\varphi(k, \omega, \cdot) = \varphi(\theta^{k-1}(\omega), \cdot) \circ \dots \circ \varphi(\omega, \cdot)$.

2.2 Fixed point random Hartman-Grobman Theorem

Analogous to most of proofs of the continuous HGT, the arguments of our proofs of the random versions of this theorem along hyperbolic stationary trajectory are based on the discrete local version for a hyperbolic fixed point of random diffeomorphisms.

We shall denote by $C_0^1(\Omega, \mathbf{R}^d)$ the space of random C^1 -local diffeomorphisms which fix the origin. The discrete cocycle generated by the pair (f, θ) is defined by:

$$\varphi(n, \omega)(x) = f(\theta^{n-1}) \circ \dots \circ f(\theta(\omega)) \circ f(\omega)(x).$$

Its linearization is still a cocycle, i.e., if we denote by $A(\omega) = D_0 f(\omega)$ the derivative of f at the origin, then,

$$D\varphi(n, \omega)(x) = A(\theta^{n-1}) \circ \dots \circ A(\theta(\omega)) \circ A(\omega)(x).$$

For reader's convenience we rewrite this preliminary Hartman-Grobman result for random diffeomorphisms with hyperbolic fixed point:

Theorem 2.1 (HGT: fixed point, local discrete case) *With the notations above, let $f \in C_0^1(\Omega, \mathbf{R}^d)$ be a random local diffeomorphism whose linearized dynamical system generated by (A, θ) is hyperbolic. Then, for \mathbb{P} -a.s., there exists a positive random variable $v(\omega)$ and a local homeomorphism $h \in \text{Homeo}(\Omega, B(0, v(\omega)); h(B(0, v(\omega))))$ such that:*

$$f(\omega, x) = h^{-1}(\theta\omega)A(\omega)h(\omega)(x),$$

for all x in the domain of the composition. The random conjugacy h is unique of the form $I + u$ with $u \in C_{0,b}(\Omega, \mathbb{R}^m)$.

Proof:

The proof is quite technical, see [7]. To figure out the idea of the localization, we recall that there is a requirement that the non-linear component $\Psi = f - A$ is Lipschitz in a neighbourhood of the origin with respect to the random norm with an appropriated Lipschitz constant equals to L . The relation between this constant and the random radius $0 \leq v(\omega) \leq 1$ is such that for all x in the ball $B(0, 2v(\omega))$ we have

$$\|D_x \Psi(\omega, \cdot)\| \leq \frac{L}{6 e B(\omega)^2}.$$

Locally this is obviously satisfied since $D_x \Psi$ is continuous with respect to x and $D_0 \Psi \equiv 0$. □

3 Discrete case with stationary orbits

This section presents a generalization of Theorem 2.1 to hyperbolic stationary orbits, in particular, to hyperbolic invariant probability measure; it corresponds to the first extension of HGT for fixed point.

We shall decompose a cocycle around a stationary trajectory writing:

$$\varphi(\omega, \cdot) = D_{Y(\omega)} \varphi(\omega, \cdot) + \Psi(\omega, \cdot),$$

where $D_{Y(\omega)} \varphi$ is the derivative of the cocycle φ at the stationary point $Y(\omega)$ and Ψ is its non-linear part.

Theorem 3.1 (HGT, discrete stationary orbit) *Let $Y(\omega)$ be a hyperbolic stationary orbit for a time-discrete C^1 -cocycle $\varphi(k, \omega, \cdot)$. Then, there exists a random homeomorphism $h(\omega) : U(\omega) \rightarrow W(\omega)$ where $U(\omega)$ is a random neighbourhood of $Y(\omega)$ and $W(\omega)$ is a neighbourhood of the origin in the tangent space $T_{Y(\omega)} \mathbf{R}^d$, such that:*

$$\varphi(\omega, x) = h^{-1}(\theta(\omega)) \circ (D_{Y(\omega)} \varphi(\omega, \cdot)) \circ h(\omega),$$

for all x in the domain of the composition.

Proof:

The argument consists of using an adequate centralization of the cocycle around the stationary orbit in the following sense: define another cocycle $\hat{\varphi}$ by

$$\hat{\varphi}(\omega, x) := \varphi(\omega, x + Y(\omega)) - Y(\theta(\omega)).$$

One checks that $\hat{\varphi}$ is a new cocycle such that the origin is a hyperbolic fixed point. The linearized system is now given by $D_0\hat{\varphi}(t, \omega) = D_{Y(\omega)}\varphi(t, \omega, \cdot) \circ I$, where the identity $I : T_0\mathbf{R}^d \rightarrow T_{Y(\omega)}\mathbf{R}^d$ is the derivative of the translation $x \mapsto x + Y(\omega)$.

By Theorem 2.1, there exists a local random topological conjugacy $\hat{h}(\omega) : \hat{U}(\omega) \rightarrow \hat{V}(\omega)$ where $\hat{U}(\omega)$ and $\hat{V}(\omega)$ are open neighbourhood of the origin in \mathbf{R}^d , such that

$$\hat{h}(\theta\omega) \circ \hat{\varphi}(\omega, \cdot) = D_0\hat{\varphi}(\omega, \cdot) \circ \hat{h}(\omega).$$

Finally, just define:

$$h(\omega)(x) = \hat{h}(\omega)(x - Y(\omega)).$$

The formula of the statement holds with $U(\omega) = Y(\omega) + \hat{U}(\omega)$. □

Let μ be an ergodic invariant probability measure on \mathbb{R}^d for a given random dynamical systems generated by (φ, θ) as in the hypothesis of Theorem 3.1 (in particular for a Markovian i.i.d. systems). One method of finding stationary orbits is constructing an equivalent cocycle on an enlarged underlying probability space. Let $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ be the following probability space:

$$\tilde{\Omega} = \Omega \times \mathbb{R}^d, \quad \tilde{\mathcal{F}} = \mathcal{F} \otimes \mathcal{B}(\mathbb{R}^d), \quad \tilde{\mathbb{P}} = \mathbb{P} \otimes \mu.$$

The family of ergodic transformation in $\tilde{\Omega}$ will be defined by:

$$\tilde{\theta}(\tilde{\omega}) = (\theta(\omega), \varphi(\omega, x))$$

with $\tilde{\omega} = (\omega, x)$. One easily sees that $\tilde{\theta}$ is $\tilde{\mathbb{P}}$ -preserving and ergodic, Carverhill [6]. We define the new equivalent cocycle $\tilde{\varphi}(x, \tilde{\omega})$ by:

$$\tilde{\varphi}(x, \tilde{\omega}) = \varphi(x, \omega)$$

with $\omega = \pi_1(\tilde{\omega})$. Note that $\tilde{Y} : \tilde{\Omega} \rightarrow \mathbb{R}^d$ given by $\tilde{Y}(\tilde{\omega}) = \pi_2(\tilde{\omega})$ is a stationary orbit for the cocycle generated by $(\tilde{\varphi}, \tilde{\theta})$ over the probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$.

Corollary 3.2 (HGT on invariant probability measures) *Let μ be an invariant ergodic probability measure on \mathbf{R}^d for a Markov process generated by a C^1 -cocycle φ . If the system is hyperbolic μ -a.s., then, for μ -a.e. $x \in \mathbf{R}^d$, there exists a random homeomorphism $h(x, \omega) : U(x, \omega) \rightarrow W(x, \omega)$ where $U(x, \omega)$ is a random local neighbourhood of x and $W(x, \omega)$ is a neighbourhood of the origin in $T_x\mathbf{R}^d$, such that:*

$$\varphi(\omega, y) = h^{-1}(\varphi(\omega, x), \theta(\omega)) \circ (D_x\varphi(\omega, \cdot)) \circ h(x, \omega)(y),$$

for all y in the domain of the composition.

Proof:

As stated before, $\tilde{Y}(\tilde{\omega})$ is a stationary trajectory for the cocycle $\tilde{\varphi}(\tilde{\omega}, \cdot)$. By Theorem 3.1, for each $\tilde{\omega} = (x, \omega)$ there exists a neighbourhood $V_{(x, \omega)}$ and a random homeomorphism $\tilde{h}(\tilde{\omega}) : V_{(x, \omega)} \rightarrow W_{(x, \omega)} \subset T_x \mathbf{R}^d$ such that

$$\tilde{h}(\tilde{\theta}(\tilde{\omega})) \circ \tilde{\varphi}(\tilde{\omega}, y) = D_x \tilde{\varphi}(\omega, \cdot) \circ \tilde{h}(\tilde{\omega})(y).$$

Now we establish the natural definition $H(x, \omega) := \tilde{H}(\tilde{\omega})$. □

Example: Random Hénon Map.

Consider the chaotic system generated by a random Hénon map $H : (\omega, (x, y)) \mapsto (\alpha(\omega) + \beta(\omega)y - x^2, x)$, where $\alpha, \beta : \Omega \rightarrow \mathbb{R}$ are real random variables on the probability space $(\Omega, \mathcal{F}, \mathbf{P})$. Consider the system $H^{(n)}(\omega, x)$ generated by the RDS (H, θ) , with $\theta : \Omega \rightarrow \Omega$ an ergodic transformation, i.e.

$$H^{(n)}(\omega, (x, y)) = H(\theta^{n-1}\omega, \cdot) \circ \dots \circ H(\theta\omega, \cdot) \circ H(\omega, (x, y)).$$

Assume that ρ is an (ergodic) invariant probability measure on \mathbb{R}^2 for this system. By the corollary above, there exists a random homeomorphism $h(\omega, (x, y)) : U((x, y), \omega) \rightarrow W(\omega)$ defined $(\mathbf{P} \otimes \rho)$ -a.s. such that:

$$\begin{pmatrix} \alpha + \beta y - x^2 \\ x \end{pmatrix} = h^{-1}(\theta\omega, H(\omega, (x, y))) \circ \begin{pmatrix} -2x & \beta(\omega) \\ 1 & 0 \end{pmatrix} \circ h(\omega, (x, y)).$$

4 Continuous Case

To establish the main result, namely, a Hartman-Grobman theorem for hyperbolic stationary trajectories we shall first establish the result for a continuous C^1 -cocycle φ which has the origin as a hyperbolic fixed point.

We shall assume that the cocycle φ is a continuous process in the group of diffeomorphisms, in fact most of the interesting cocycles in finite dimensional state space, independently of how they are generated (stochastic differential equation, random equation, etc) satisfies this assumption, see Arnold [1] and references therein.

Initially, consider discrete systems generated by time-one map $\varphi(1, \omega, \cdot)$. If the origin is a hyperbolic fixed point for the continuous system, it is also a hyperbolic fixed point for this discretization. Let $V(\omega) = B(0, v(\omega))$ be the neighbourhood of conjugacy for this discrete cocycle, as stated in Theorem 2.1. With this notation we have:

Lemma 4.1 *Let $\varphi(t, \omega, \cdot)$ be a C^1 -cocycle on \mathbf{R}^d such that the origin is a hyperbolic fixed point. There exists a neighbourhood $U(\omega) = B(0, u(\omega))$ such that:*

$$\varphi(t, \omega, x) \in V(\theta_t \omega),$$

for all $x \in U(\omega)$ and $t \in [0, 2]$. Obviously $u(\omega) \leq v(\omega)$.

Proof:

Firstly, note that, by hyperbolicity, the derivative $D_0\varphi(t, \omega, \cdot)$ is a non-singular matrix, hence, there exists a ball centered at zero with radius $R(t, \omega) > 0$ where $\varphi(t, \omega, \cdot)$ is invertible. Let $R(\omega)$ be the (measurable) random radius defined by

$$R(\omega) = \inf\{R(t, \omega); t \in [0, 2]\}.$$

By continuity on t the variable $R(\omega)$ is strictly positive.

We claim that $\varphi(1, \theta_s \omega, x)$ is continuous on $s \in [0, 2]$ for all x in $B(0, R(\omega))$. In fact, in this case:

$$\varphi(1, \theta_s \omega, x) = \phi(1 + s, \omega, \cdot) \circ \phi(s, \omega, x)^{-1},$$

and the expression on the right hand side is continuous on s .

This continuity of $\varphi(1, \theta_s \omega, x)$ on s implies that the radius of conjugacy in the discrete (time-one) case $v(\theta_s \omega)$, as defined in the proof of Theorem 2.1, is also continuous on $s \in [0, 2]$. Hence,

$$r := \inf\{v(\theta_s \omega), s \in [0, 2]\} > 0,$$

for all $\omega \in \Omega$. By continuity of $\varphi(t, \omega, x)$ on t , for each ω there exists a strictly positive radius $u(\omega)$ such that $\|\varphi(t, \omega, x)\| < r$ for all x such that $\|x\| < u(\omega)$. Moreover, one can choose $u(\omega)$ such that it is measurable. \square

As we said before, considering the lemma above, the proof of the local HGT goes with the same ideas of the global version, [7, Thm 5.1], which is in fact a random adaptation of the ideas in Palis and de Melo [14] or S. Sternberg [18, Lemma 4].

We shall write, as before

$$\varphi(t, \omega, \cdot) = \Phi(t, \omega, \cdot) + \Psi(t, \omega, \cdot),$$

where Φ is the linearization at the origin $\Phi(t, \omega, \cdot) = D_0\varphi(t, \omega, \cdot)$.

Theorem 4.1 (HGT for continuous cocycles, fixed point) *Consider $\varphi(t, \omega, x)$, a C^1 -cocycle on \mathbf{R}^d . If the origin is a hyperbolic fixed point, then there exists a random conjugacy between φ and its linearization $\Phi = D\varphi$.*

More precisely: there exists a random neighbourhood $U(\omega) = B(0, u(\omega))$ and a measurable random homeomorphism $H(\omega) : U(\omega) \rightarrow H(U(\omega))$ such that

$$\varphi(t, \omega, \cdot) = H^{-1}(\theta_t \omega) \Phi(t, \omega) H(\omega)(\cdot).$$

for all x in $U(\omega)$ and $t \in [-1, 1]$.

Proof:

As stated before, let h be the random conjugacy of Theorem 3.1 for the discrete system generated by $\varphi(1, \omega, x)$. Let $u(\omega)$ be the strictly positive random radius defined in the lemma above. For $x \in U(\omega)$, define:

$$H(\omega, x) = \int_0^1 \Phi(-s, \theta_s \omega) h(\theta_s \omega) \varphi(s, \omega, x) ds. \quad (3)$$

Lemma 4.1 guarantees that the composition makes sense and that the integrand is continuous on s , hence $H(\omega)$ is well-defined. Note also that, by hyperbolicity, $\Phi(s, \omega)$ is an invertible cocycle and $\Phi(-s, \theta_s \omega) = \Phi(s, \omega)^{-1}$.

Let t be in the interval $[-1, 1]$, then

$$\Phi(t, \omega) H(\omega)(x) = \int_0^1 \Phi(t-s, \theta_s \omega) h(\theta_s \omega) \varphi(s-t, \theta_t \omega, \cdot) ds \circ \varphi(t, \omega, x).$$

Changing the variable $r = s - t$, we have:

$$\begin{aligned} \Phi(t, \omega) H(\omega)(x) &= \int_{-t}^{1-t} \Phi(-r, \theta_{r+t} \omega) h(\theta_{r+t} \omega) \varphi(r, \theta_t \omega, \cdot) dr \circ \varphi(t, \omega, x) \\ &= \left[\int_{-t}^0 \Phi(-r, \theta_{r+t} \omega) h(\theta_{r+t} \omega) \varphi(r, \theta_t \omega, \cdot) dr \right. \\ &\quad \left. + \int_0^{1-t} \Phi(-r, \theta_{r+t} \omega) h(\theta_{r+t} \omega) \varphi(r, \theta_t \omega, \cdot) dr \right] \circ \varphi(t, \omega, x). \end{aligned}$$

The first integral inside the bracket equals:

$$\int_{-t}^0 \Phi(-r-1, \theta_{r+t+1} \omega) [\Phi(1, \theta_{t+r} \omega) h(\theta_{r+t} \omega) \varphi(-1, \theta_{r+t+1} \omega, \cdot)] \varphi(r+1, \theta_t \omega, \cdot) dr. \quad (4)$$

But Theorem 2.1 states that

$$\Phi(1, \theta_{t+r} \omega) h(\theta_{r+t} \omega) \varphi(-1, \theta_{r+t+1} \omega, \cdot) = h(\theta(\theta_{r+t} \omega)).$$

Note that the domain makes sense because the formula above is applied to a point

$$\varphi(r+1, \theta_t \omega, \cdot) \circ \varphi(t, \omega, x) = \varphi(t+r+1, \omega, x) \in B(0, u(\theta_{t+r+1} \omega)),$$

with $t + r + 1 < 2$.

Hence, expression (4) equals:

$$\int_{-t}^0 \Phi(-r - 1, \theta_{r+t+1}\omega) h(\theta(\theta_{r+t}\omega)) \varphi(r + 1, \theta_t\omega, \cdot) dr,$$

which, changing variables again with $s = r + 1$, becomes:

$$\int_{1-t}^1 \Phi(-s, \theta_{s+t}\omega) h(\theta_{s+t}\omega) \varphi(s, \theta_t\omega, \cdot) ds.$$

Hence,

$$\begin{aligned} \Phi(t, \omega) H(\omega)(x) &= \left[\int_{1-t}^1 \Phi(-s, \theta_{s+t}\omega) h(\theta_{s+t}\omega) \varphi(s, \theta_t\omega, \cdot) ds \right. \\ &\quad \left. + \int_0^{1-t} \Phi(-r, \theta_{r+t}\omega) h(\theta_{r+t}\omega) \varphi(r, \theta_t\omega, \cdot) dr \right] \circ \varphi(t, \omega, x) \\ &= \left[\int_0^1 \Phi(-s, \theta_{s+t}\omega) h(\theta_{s+t}\omega) \varphi(s, \theta_t\omega, \cdot) ds \right] \circ \varphi(t, \omega, x) \\ &= H(\theta_t\omega) \varphi(t, \omega, x). \end{aligned}$$

We still have to prove that $H(\omega)$ is indeed a homeomorphism. We shall conclude this fact by showing that $H = I + u$ with u in the space $C_{0,b}(\Omega, \mathbb{R}^m)$. Therefore, by the uniqueness stated in Theorem 2.1, $H(\omega) = h(\omega)$ a.s.. For readers convenience, we leave this part of the prove to be done in the next lemma.

□

Lemma 4.2 *The random conjugacy H has the form $H = I + u$ with u in the space $C_{0,b}(\Omega, \mathbb{R}^m)$.*

Proof: We write

$$H = I + \int_0^1 \Phi(-s, \theta_s\omega) (\Psi(s, \omega) + u(\theta_s\omega) \varphi(s, \omega)) ds$$

and note that the component

$$\int_0^1 \Phi(-s, \theta_s\omega) u(\theta_s\omega) \varphi(s, \omega) ds$$

belongs to $C_{0,b}(\Omega, \mathbb{R}^m)$, since $u \in C_{0,b}(\Omega, \mathbb{R}^m)$ and Φ is linear. Moreover, given an $\varepsilon > 0$, there exists a random variable $\delta(\omega) \leq u(\omega)$ such that, if $x \in B(0, \delta(\omega))$, then for $s \in [0, 1]$:

$$\|\Psi(s, \omega, x)\|_{\theta_s \omega} \leq \varepsilon.$$

Hence the other component

$$\int_0^1 \Phi(-s, \theta_s \omega)(\Psi(s, \omega))$$

also belongs to $C_{0,b}(\Omega, \mathbb{R}^m)$. □

Theorem 4.2 (HGT for continuous stationary trajectories) *Let $Y(\omega)$ be a hyperbolic stationary trajectory for a cocycle $\varphi(t, \omega, \cdot)$. Then, there exists a random homeomorphism $H(\omega) : U(\omega) \rightarrow W(\omega)$ where $U(\omega)$ is a neighbourhood of $Y(\omega)$ and $W(\omega)$ is a neighbourhood of origin in the tangent space $T_{Y(\omega)}\mathbf{R}^d$ such that:*

$$\varphi(t, \omega, \cdot) = H^{-1}(\theta_t \omega) \circ D_{Y(\omega)}\varphi(t, \omega, \cdot) \circ H(\omega),$$

for all $t \in [-1, 1]$.

Proof:

Again, using the same centralization argument as in Theorem 3.1, define a new cocycle $\hat{\varphi}$ by:

$$\hat{\varphi}(t, \omega, x) := \varphi(t, \omega, x + Y(\omega)) - Y(\theta_t \omega).$$

The linearized flow is now given by $D_0\hat{\varphi}(t, \omega) = D_{Y(\omega)}\varphi_t \circ I$, where the identity $I : T_0\mathbf{R}^d \rightarrow T_{Y(\omega)}\mathbf{R}^d$ is the derivative of the translation $x \mapsto x + Y(\omega)$.

By Theorem 4.1 there exists a random conjugacy $\hat{H}(\omega) : U(\omega) \rightarrow V(\omega)$ where $U(\omega)$ and $V(\omega)$ are open neighbourhood of the origin in \mathbf{R}^d , such that

$$\hat{\varphi}(t, \omega, \cdot) = \hat{H}^{-1}(\theta_t \omega) \circ D_0\hat{\varphi}(t, \omega, \cdot) \circ \hat{H}(\omega).$$

Finally, as in the discrete case, just define:

$$H(\omega)(x) = \hat{H}(\omega)(x - Y(\omega)).$$

□

Note that, if φ is a solution flow of an Itô stochastic differential equation, the cocycle $\hat{\varphi}$ is non-adapted (hence is not a solution of any Itô stochastic

equation). Due to this fact, there is a good advantage on having a HGT for general cocycle instead of the particular case of solutions of stochastic differential equations.

Analogously to the discrete case, if the cocycle generates a Markov processes (typically, consider those cocycles which are either deterministic flows or solution flows of stochastic differential equations), if there exists μ an ergodic invariant probability measure on \mathbb{R}^d for $\varphi(t, \omega, \cdot)$, we can construct an stationary trajectory by constructing an equivalent cocycle on an enlarged underlying probability space. Again, consider the probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, as defined before Corollary 3.2. The family of ergodic transformation is now given by:

$$\tilde{\theta}_t(\tilde{\omega}) = (\theta_t(\omega), \varphi_t(\omega, x))$$

It preserves the ergodic probability measure $\tilde{\mathbb{P}}$. Define the new equivalent cocycle $\tilde{\varphi}_t(\tilde{\omega}, x)$ by:

$$\tilde{\varphi}_t(\tilde{\omega}, x) = \varphi_t(\omega, x)$$

with $\omega = \pi_1(\tilde{\omega})$. Note that $\tilde{Y} : \tilde{\Omega} \rightarrow \mathbb{R}^d$ given by $\tilde{Y}(\tilde{\omega}) = \pi_2(\tilde{\omega})$ is a stationary trajectory for the cocycle generated by $(\tilde{\varphi}_t, \tilde{\theta})$ over the probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$.

Corollary 4.3 (HGT on invariant probability measures) *Let μ be an invariant ergodic probability measure on \mathbf{R}^d for a Markov process associated with the cocycle φ (say, solution of an stochastic differential equation). Assume hyperbolicity μ -a.s., then, for μ -a.e. $x \in \mathbf{R}^d$, there exists a random homeomorphism $H(x, \omega) : U(x, \omega) \rightarrow W(x, \omega)$ where $U(x, \omega)$ is a random neighbourhood of x and $W(x, \omega)$ is a neighbourhood of the origin in the tangent space $T_x \mathbf{R}^d$ such that:*

$$\varphi(t, \omega, y) = H^{-1}(x_t(\omega), \theta_t \omega) \circ D_{x_t} \varphi(t, \omega, \cdot) \circ H(x, \omega)(y),$$

for all y in the domain of the composition, where $x_t = \varphi(t, \omega, x)$ and $t \in [0, 1]$.

Proof:

The proof is essentially the same as in the discrete case. Take $\tilde{Y}(\omega)$ the stationary trajectory for the cocycle $\tilde{\varphi}(t, \tilde{\omega}, \cdot)$. By Theorem 4.2, for each $\tilde{\omega} = (x, \omega)$ there exists a neighbourhood $V_{(x, \omega)}$ and a random homeomorphism $\tilde{H}(\tilde{\omega}) : V_{(x, \omega)} \rightarrow W_{(x, \omega)} \subset T_x \mathbf{R}^d$ such that

$$\tilde{H}(\tilde{\theta}_t(\tilde{\omega})) \circ \tilde{\varphi}(t, \tilde{\omega}, x) = T_{\tilde{Y}(\tilde{\omega})} \tilde{\varphi}(t, \omega, \cdot) \circ \tilde{H}(\tilde{\omega})(x).$$

Now state the natural definition $H(x, \omega) = \tilde{H}(\tilde{\omega})$.

□

Figure 1 illustrates the result of Theorem 4.2. The linearized trajectory $v_t = D\varphi(v_0)$ equals the image $H(x_t, \omega)(y_t)$ of some $y_t = \varphi(t, \omega, y_0)$ with $y_t \in V(x_t, \theta_t \omega)$ for all $t \in [-1, 1]$. It gives a pictorial view of the fact that the linearized cocycle $D\varphi$ together with the random homeomorphism H characterize completely the cocycle φ in a neighbourhood of a hyperbolic stationary trajectory.

For stochastic dynamical systems on a Riemannian manifold, this centralization argument does not work straightforward once the group structure of \mathbb{R}^d is intrinsic in the argument. Nevertheless, by local coordinates, the conjugacy in \mathbb{R}^d is transported to the manifold, namely:

Corollary 4.4 *Let φ be a C^1 -cocycle in a Riemannian manifold M . If $Y(\omega)$ is a hyperbolic stationary point, then, there exist a random local homeomorphisms $H(\omega) : U(\omega) \subset M \rightarrow W(\omega)$ where $W(\omega)$ is a neighbourhood of origin in $T_{Y(\omega)}M$ such that:*

$$\varphi(t, \omega, \cdot) = H^{-1}(\theta_t \omega) \circ D_{Y(\omega)}\varphi(t, \omega, \cdot) \circ H(\omega),$$

for all x and t which are in the domain of the composition (restricted to local coordinate neighbourhoods).




Figure 1: Tubular neighbourhood of conjugacy in the tangent bundle.

We recall the example of Baxendale [4] of a hyperbolic Brownian motion on the flat torus T^2 (moreover, with non-zero rotation number, see Ruffino [16]). The ergodic invariant probability measure is the Lebesgue measure λ . The corollary above states that for $\mathbb{P} \otimes \lambda$ -a.e. (ω, x_0) , given the trajectory $x_t(\omega) = \varphi_t(\omega, x_0)$ of this Brownian motion, there is a ‘traveling’ neighbourhood of $x_t(\omega)$ such that the dynamics around $x_t(\omega)$ in the torus is conjugate to the linearized dynamics in the tangent bundle.

5 Applications

In this section we describe explicitly the applications of Theorems 4.2 and 3.1 on deterministic dynamical systems. We intend rather to show directions of further problems. The theorems below are particular cases of Corollaries 3.2 and 4.3, one just have to lift the Markovian hypothesis and consider the probability space only as an ergodic invariant probability measures in the state space.

Theorem 5.1 (HGT, discrete chaotic systems) *Let ρ be an ergodic invariant probability measure on \mathbf{R}^d for an application $f : U \subset \mathbf{R}^d \rightarrow \mathbf{R}^d$. If the system is hyperbolic ρ -a.s., then, for ρ -a.e. $x \in \mathbf{R}^d$, there exists a homeomorphism $h(x) : U(x) \rightarrow W(x)$ where $U(x)$ is a neighbourhood of x and $W(x)$ is a neighbourhood of the origin in the tangent space $T_x\mathbf{R}^d$, such that:*

$$f(y) = h^{-1}(f(x)) \circ (D_x f(\cdot)) \circ h(x)(y),$$

for all y in the domain of the composition.

Theorem 5.2 (HGT, continuous deterministic systems) *Let μ be an invariant ergodic probability measure on \mathbf{R}^d for the flow φ_t associated to an ODE. Assume hyperbolicity in μ -a.s., then, for μ -a.e. $x \in \mathbf{R}^d$, there exists a homeomorphism $H(x) : U(x) \rightarrow W(x)$ where $U(x)$ is a neighbourhood of x and $W(x)$ is a neighbourhood of the origin in the tangent space $T_x\mathbf{R}^d$ such that:*

$$\varphi_t(y) = H^{-1}(x_t)(\cdot) \circ D_{x_t} \varphi_t(\cdot) \circ H(x)(y),$$

for all y in the domain of the composition, where $x_t = \varphi_t(x)$ and $t \in [0, 1]$.

Among other systems, we recall that the Lorentz attractors fits perfectly for an analysis in view of Theorem 5.2, see e.g. Sparrow [17].

The results above suggest to apply the ideas of Cong [8] on topological classification of linear cocycles to generalize the classical classification of dynamics in a neighbourhood of hyperbolic fixed point to stationary hyperbolic orbits, i.e., with ergodic invariant measures with hyperbolic Lyapunov spectrum.

References

- [1] ARNOLD, L. – *Random Dynamical Systems*. Springer, Berlin (1998).
- [2] ARNOLD, L. and M. SCHEUTZOW – Perfect cocycles through stochastic differential equations. *Prob. Th. Rel. Fields*, **101**, p. 65–88, 1995 .

- [3] ARNOLD, L. AND P. IMKELLER, – Stratonovich calculus with spatial parameters and anticipative problems in multiplicative ergodic theory, *Stochastic Processes and their Applications*, **62**, 19-54, 1996.
- [4] BAXENDALE, P. – Asymptotic behaviour of stochastic flow of diffeomorphisms. In: *Stochastic Processes and their Applications- Proceedings of the International Conference in Nagoya, July 2–6, 1985*. Ed. K. Itô and T. Hida. Lecture Notes Math. **1203**. Springer, 1986.
- [5] BOXLER, P. – A Stochastic Version of Center Manifold Theory. *Probability Theory and Related Fields*. **83**, (1989), 509-545.
- [6] CARVERHILL, A. – Flows of stochastic dynamical systems: Ergodic theory, *Stochastics*, **14** (1985), 273-317.
- [7] COAYLA-TERAN, E.A. AND P.R.C. RUFFINO – Stochastic Versions of Hartman-Grobman theorems. *Stochastics and Dynamics*, **4**(4)(2004), 571-593.
- [8] CONG, N. D. – Topological classification of linear hyperbolic cocycles, *Journal of Dynamics and Differential Equations*, **8**, no. 3 (1996), 427-467.
- [9] GROBMAN, D. M. – Homeomorphisms of systems of differential equations, *Dokl. Akad. Nauk SSSR*, **128** (1959), 880-881.
- [10] HARTMAN, P. – A lemma in the theory of structural stability of differential equations. *Proc. A.M.S.* **11** (1960), 610-620.
- [11] HARTMAN, P. – On local homeomorphisms of Euclidean spaces, *Bol. Soc. Math. Mexicana*, **5** (1960), 220-241.
- [12] KATOK, A. AND B. HASSELBLATT – *Introduction to the modern theory of dynamical systems*. CUP, 1999.
- [13] MOHAMMED, S.-E. A. AND M. SCHEUTZOW – The stable manifold theorem for stochastic differential equations, *Annals of Prob.*, Vol. **27**, no. 2, pp. 615-652, 1999.
- [14] PALIS JR., J. AND W. DE MELO – *Geometric theory of Dynamical systems: an introduction*. Springer-Verlag, New York, Heidelberg, Berlin, 1982.
- [15] RUELLE, D. – Ergodic theory of differentiable dynamical systems, *Publ. IHES* **50** (1979) 275-305.

- [16] RUFFINO, P. R. C. – Rotation number for stochastic dynamical systems, *Stoch. and Stoch. Reports*, **60** (1997), pp. 289-318.
- [17] SPARROW, C. – *The Lorentz Equations: Bifurcations, Chaos and Strange attractors*. Springer-Verlag, New York, 1982.
- [18] STERNBERG, S. – Local contractions and a theorem of Poincaré. *Amer. J. Math.* vol. **79** (1957), 809-824.
- [19] WANNER, T. – Linearization of random dynamical systems. In John, C., Kirchgraaber, U., and Walter, H. O. (Eds.), *Dynamics Reported*, Vol. **4**, Springer, Berlin, Heidelberg, New York (1994).