Conservation Efforts through *In Vitro* Seed Germination of a Hepatoprotective Plant - *Phyllanthus beddomie* (Gamble) Mohanan

Maridass, M¹ and K. Thangavel²

¹Animal Health Research Unit, St. Xavier's College, Palayamkottai, TN, India ²Depart of Biotechnology, Sri Paramakalyani College, Alwarkurichi, Thirunelveli (D.t), Tamil Naud, India Author for correspondence: E-mail: orchidevadass@vahoo.com

Issued 22 June 2008

Introduction

Whole Plants and plant parts used in ethnomedicine contain curative agents that are used in many modern medicines. The *Phyllanthus* genus contains over 600 species of shrubs, trees and annual or biennial herbs distributed throughout the tropical and subtropical regions of both hemispheres (Figueira *et al.*, 2006), and have long been used in traditional medicine to treat chronic liver diseases (Dhiman *et al.*, 2005). *Phyllanthus* appears to be promising in patients with chronic hepatitis B virus (HBV) infection (Thyagarajan *et al.*1988; 1990;1999). Bioactive principles like alkaloids, tannins, flavonoids, lignans, phenols and terpenes have been isolated from various species of *Phyllanthus* and their compounds showed antinociceptive activity (Cechinel Filho et al.1996). Liu *et al.* (2001) published a meta-analysis of the effect on and safety of genus *Phyllanthus* for chronic HBV infection.

In - vitro propagation of endangered plants can offer considerable benefits for the rapid cultivation of species that are at risk, that have limited reproductive capacity and exist in threatened habitats (Fay, 1992). In vitro propagation methods are essential components of plant genetic resources management and they are becoming increasingly important for conservation of rare and endangered plant species (Sudha et al. 1998; Benson et al. 2000; Iankova et al. 2001; Bhatia et al. 2002). The successful use of plant tissue culture techniques for the micropropagation of members of the genus Phyllanthus has been reported for species such as P. emblica, P. urinaria, P. amarus, P. abnormis, P. caroliniensis, P. tenellus, and P. niruri (Unander, 1991; Ishimaru et al., 1992; Santos et al., 1994). Phyllanthus beddomie (Gamble) Mohanan is a sub-shrub found in evergreen forest and the flowering and fruiting season is April - May. Viswanathan et al., (2002) reported the existence of this species (rediscovered) after a lapse of about 73 years in 1998 (collected from Kalakad Mundanthurai Tiger Reserve, Tirunelveli District), and they have treated this species as critically endemic one. Thus, there is an urgent need for the conservation of this species. There are no reports on the *in-vitro* micropropagation of *P. beddomie*. The present study represents a preliminary approach for the development of a rapid and reproducible protocol forin vitro multiplication of *P. beddomie*, which may be helpful in the conservation of this medicinal plant. Moreover, this type of approach may be a vital tool to have a continuous supply of raw materials for the pharmaceutical industry throughout the year to formulate antihepatitic medicines.

Materials and Methods

In vitro culture of Phyllanthus beddomie

Phyllanthus beddomie seeds were collected from Kalakad Mundanthurai Tiger Reserve forest area (Kodhaiyar) and sterilized by immersion in 70% ethanol for 5sec, followed by immersion in 0.5% aqueous solution of sodium hypochlorite with one drop of Tween 20 per 500ml for 20min, and rinsed three times in sterile double distilled water. Surface sterilized seeds were sowed for germination in MS medium. Ten duplicate sets were maintained (one in light and one in complete dark condition). Prior to this, an attempt was made to test the viability of the seeds by sowing them in fertile soil. But there was no appreciable rate of germination.

Sowed seeds in MS medium (Murashige and Skoog, 1962) were incubated for 21 days in darkness in a growth cabinet at 24 °C. Varying levels cytokinins like BAP and Kinetin (0.10 – 4.00 µM) and auxins like NAA, IAA, IBA (Sigma Chemicals, USA) supplemented with MS medium. For all the combinations, 10 duplicates were maintained. During the course of incubation, parameters like shooting, rooting and callusing were observed, average values were calculated and tabulated (Table 1).

Results and Conclusion

Among the tested concentrations and combinations of PGRs, auxins were able to facilitate the rooting phenomenon than that of shooting (Benson, 2000). Even in case of the duplicates supplemented with cytokinins there was no notable percentage of shooting. This may be due to the improper placement of plumule during the initial stage of germination. Among the auxins tested, IBA was able to enhance the rooting ability of the implants at the maximum level (90%) in the concentration of 4μ M. These observations were indicating the indigenous level of hormones especially the auxins in the seeds specifically during the initial phases of germination (Table.1)

Regarding the nature of emerged roots, sturdy nature of the roots was proportional to the increasing concentration of auxins like IBA and NAA (Fig.2). In case of the roots developed from other sets i.e. media combinations supplemented with cytokinins and in low levels of auxins, there appeared fragile and less sturdy ones (Fig.1).

No responses in root and shoot formation of MS medium containing BA, and Kinetin in low concentrations (0.10 – 2.00 μ M). The findings of the present study may be helpful for the development of micropropagation of *P. beddomie*. Further studies are ongoing for the further establishment and ex *situ* conservation of *P. beddomie* and such type of medicinal plants are in need to fulfill the present need of the pharmaceutical industry (Fay, 2002).


Table 1. The frequency (%) of seed germination responses of *P. beddomie* after 21 days on MS medium supplemented with PGRs in different concentrations.

PGRs used	Concentration	Shooting (%)	Rooting (%)	Callusing (%)
BAP	0.10	-	-	-
	0.50	-	-	-
	1.00	-	-	-
	2.00	-	-	-
	3.00	-	-	-
	4.00	-	-	-

I	1	I	1	1
	0.10	_	0	
Kinetin		-		-
	0.50	-	20	-
	1.00	-	40	-
	2.00	-	50	-
	3.00	-	60	-
	4.00	-	60	-
IAA	0.10	-	-	-
	0.50	-	-	-
	1.00	-	-	-
	2.00	-	-	-
	3.00	-	-	-
	4.00	-	-	-
IBA	0.10	-	20	-
	0.50	-	35	-
	1.00	-	40	-
	2.00	-	60	-
	3.00	-	80	-
	4.00	-	90	-
NAA	0.10	-	10	-
	0.50	-	10	_
	1.00	-	20	_
	2.00	-	20	_
	3.00	-	25	_
	4.00	-	40	-



Live endemic medicinal plant Phyllanthus beddomei



Phyllanthus beddomei seed dormancy of ms medium

Phyllanthus beddomei regenerated from seed

Acknowledgement:

The authors wish to thank the Department of Science and Technology, Govt. of India, New Delhi for providing financial assistance.

References

- Benson, E.E., Danaher J.E., Pimbley I.M., Anderson C.T., Wake J.E., Daley S. and Adams, L.K. 2000. *In vitro* micropropagation of *Primula scotica*: a rare Scottish plant. *Biodiversity and Conservation*, **9**: 711–726.
- Bhatia P., Bhatia N.P. and Ashwath N. 2002. *In -vitro* propagation of *Stackhousia tryonii* Bailey (Stackhousiaceae): a rare and serpentine-endemic species of central Queensland, Australia. *Biodiversity and Conservation*, **11**: 1469–1477.
- Cechinel Filho, V., Santos, A.R.S., Campos, R.O.P., Miguel, O.G., Yunes, R.A., Ferrari, F., Messana, J. & Calixto, J.B. 1996. Chemical and pharmacological studies of *Phyllanthus caroliniensis* in mice. *Journal of Pharmacology*, **48**:1231-1236.
- Fay, M. 1992. Conservation of rare and endangered plants using *in -vitro* methods. *In Vitro Cellular Developmental Biology*, **28**: 1-4.
- Gamborg, O.L., Miller, R.A, Ojima, K.1968. Nutrient requirement of suspension cultures of soybean root cells. *Exp. Cell. Res.*, **50**:150–158.
- Iankova E., Cantos M., Lin, J., Robeva P. and Troncoso A. 2001. In vitro propagation of *Angelica pancicii* Vauds., an endangered plant species in Bulgaria. *Seed Science and Technology*, **29**: 477–482.
- Ishimaru, K, Yoshimatsu, K, Yamakawa, T, Kamada, H and Shimomura, K (1992) Phenolic constituents in tissue cultures of *Phyllanthus niruri*. *Phytochemistry*, **34**: 2015 –2016.
- Liu, J., Lin, H, McIntosh, H.2001. Genus *Phyllanthus* for chronic hepatitis B virus infection: a systematic review. *J. Viral Hepat.*, **8**:358–366.
- Murashige T. and Skoog F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, **15**: 473–497.
- Santos, A.R.S., Cechinel Filho, V., Viana, A.M, Moreno, F.N, Campos, M.M, Yunes, R.A and Calixto, J.B.1994. Analgesic effects of callus culture extracts from selected species of *Phyllanthus* in mice. *J. Pharm. Pharmacol.* **46**: 755–759.
- Sudha, C.G., Krishnan, P.N. and Pushpahgadan, P.1998. *In-vitro* propagation of *Holostemma annulare* (Roxb.) K. Schum, a rare medicinal plant. *In Vitro Cellular and Developmental Biology Plant*, **34**: 57–63.
- Thyagarajan, S.P., Subramanian, S., Thirunaksundari, T, Venkateswaran, P.S., Blumberg, B.S.1988. Effects of *Phyllanthus amarus* on chronic carriers of hepatitis B virus. *Lancet*, **2**:764–766.
- Thyagrajan, S.P., Jayaram, S, Valliammae, T, Madangopalan, N., Pal, V.G., Jayaraman, K.1990. *Phyllanthus amarus* and hepatitis B. *Lancet*, **336**:949–950.
- Unander, D.W.1991. Callus induction in *Phyllanthus* species and inhibition of viral DNA polymerase and reverse transcriptase by callus extracts. *Plant Cell Rep.*, **10**: 461-466.
- Viswanathan, M. B., Ramesh, N., Maridass, M. and Manikandan, U. 2002. Rediscovery of a critically endangered species *Phyllanthus beddomei* (Gamble) Mohanan, Euphorbiaceae, from Kalakkad Mundanthurai Tiger Reserve in India. *Bombay Natural History Society*, **99**(3): 560-562.