

1993

Prime Decomposition of Knots in Lorenz-like Templates

Michael C. Sullivan

Southern Illinois University Carbondale, msulliva@math.siu.edu

Follow this and additional works at: http://opensiuc.lib.siu.edu/math_articles

 Part of the [Mathematics Commons](#)

Electronic version, with some figures missing, of an article published as "Prime Decomposition of Knots in Lorenz-like Templates," *Journal of Knot Theory and Its Ramifications*, 2(4), 1993, 453 - 462. DOI: 10.1142/S021821659300026X © copyright World Scientific Publishing Company, <http://www.worldscinet.com/jktr/>

Recommended Citation

Sullivan, Michael C. "Prime Decomposition of Knots in Lorenz-like Templates." (Jan 1993).

This Article is brought to you for free and open access by the Department of Mathematics at OpenSIUC. It has been accepted for inclusion in Articles and Preprints by an authorized administrator of OpenSIUC. For more information, please contact opensiuc@lib.siu.edu.

PRIME DECOMPOSITION OF KNOTS IN LORENZ-LIKE TEMPLATES

MIKE SULLIVAN
UNIVERSITY OF TEXAS AT AUSTIN

ABSTRACT. In [7] R. F. Williams showed that all knots in the Lorenz template are prime. His proof included the cases where any number of positive twists were added to one of the template's branches. However [7] does give an example of a composite knot in a template with a single negative twist. Below we will show that in all the negative cases composite knots do exist, and give a mechanism for producing many examples. This problem was cited in a list of problems in dynamics in [1, problem 4.2].

1. INTRODUCTION

1.1. Flows, Knots and Templates. The periodic orbits of a flow in a 3-manifold form knots. These knots and how they are linked have been studied with the aid of *templates* or *knots holders*, i.e. 2-dimensional branched manifolds with semi-flows [4].



Definition 1. *The $L(m, n)$ templates are shown in Figure 1. The semi-flow comes down from the branch line, β , forming two branches. The X -branch has m half twists, while the Y -branch has n . The sign convention for the twists is that left-handed twists are positive and right-handed ones are negative. The two branches meet tangentially at the branch line.*

$\tilde{L}(m, n)$ is the mirror image of $L(m, n)$.

The symbols defined above will also be used to represent the set of knots formed by the periodic orbits in the semi-flow of the template.

$L(0, 0)$ is the Lorenz template [3]. The remainder will be called Lorenz-like templates. We also note that $L(m, n) = L(n, m)$.

Date: Appeared in Journal of Knot Theory and Its Ramifications, Vol 2, No. 4, 1993, 453–462. (Received 1 August 1991). Current e-mail: msullivan@math.siu.edu.

FIGURE 1. Lorenz-like template, $L(m, n)$, definedFIGURE 2. A square knot in $L(0, -1)$

1.2. Prime and Composite Knots.

Definition 2. A knot $k \subset S^3$ is composite if there exists a tame sphere S^2 such that $S^2 \cap k$ is just two points, p and q , and if γ is any arc on S^2 joining p to q , then the knots

$$k_1 = \gamma \cup (k \cap \text{outside of } S^2) \quad \text{and} \\ k_2 = \gamma \cup (k \cap \text{inside of } S^2),$$

are nontrivial, (i.e. not the unknot). We call k_1 and k_2 factors of k and write

$$k = k_1 \# k_2.$$

If a knot isn't composite then it is prime.

Figure 3 gives an example.




FIGURE 3. A connected sum

Finally we state the all important theorem due to Schubert [6, 5]:

Knots can be factored uniquely into primes, up to order.

2. KNOTS IN $L(m, n)$

Theorem 1. *As sets of knots, $L(0, n - 2) \supseteq L(0, n)$, for all integers n .*

This may be restated as

$$\dots \supseteq L(0, -2) \supset L(0, 0) \supseteq L(0, 2) \supseteq \dots,$$

$$\dots \supseteq L(0, -3) \supseteq L(0, -1) \supset L(0, 1) \supseteq L(0, 3) \supseteq \dots,$$

where the orientable and nonorientable cases are listed separately. We have also noted that two of the set inclusions are proper. Lorenz knots are positive braids [3], but $L(0, -2)$ has a negative braid. (To construct one, trace out the periodic orbit that wraps around the X-branch once and the Y-branch three times.) Knots in $L(0, 1)$ are prime while $L(0, -1)$ has composite knots. It is not known which, if any, of the other set inclusions are proper.

Proof. Recall the equation “writhe = twist”. We apply this in Figure 4. The illustration shows how to take a knot embedded in $L(0, n)$ over to a knot of the same knot type in $L(0, n - 2)$. □

Theorem 2. *Any knot formed by the connected sum of a knot in $L(0, 2)$ and a knot in $\tilde{L}(0, 2)$ is in $L(0, -2)$.*




FIGURE 4. Proof of Theorem 1


Proof. The proof again is pictorial. We start with $L(-2, 0)$, which is identical to $L(0, -2)$. By cutting along certain orbits and gradually deforming the template we will make the knots in question become clearly visible.

Most of the steps in Figure 5(a-w) should be self evident. Figures (a) to (n) show that the subtemplate of (n) lives in $L(-2, 0)$. In going from figure (j) to (k) we have thrown away part of the template to help make the remaining steps clearer. The template of (n) is shown in (o) with lines to cut along. This divides (o) into two parts, (p) and (t). To see that any pair of orbits on (p) and (t) respectively can be summed on (n) just study the example in Figure 6. Following from (p) to (s) we get $\tilde{L}(0, 2)$. Likewise, (t) to (w) gives $L(0, -2)$.


In Figure 6 we show a composite knot on (n) which is the square knot. In Figure 7 we display this knot on the original form of the template. The reader can check this by reversing all the steps while carrying the knot back. $\square$

PRIME DECOMPOSITION OF KNOTS IN LORENZ-LIKE TEMPLATES


a)




b)




c)




d)




e)




f)




g)




h)




i)




j)




k)




l)




m)




n)




o)




p)




t)



q)



u)






FIGURE 5. Figures (a)-(w) prove Theorem 2





FIGURE 6. Composite knot


Thus THEOREMS 1 and 2 imply that for all negative integers n , $L(0, n)$ contains composite knots. In fact THEOREM 2 shows how to construct many examples of composite knots on $L(0, -2)$ and hence

FIGURE 7. Composite knot on $L(-2, 0)$

for any Lorenz-like template with an even number of negative twists on one its branches. The theorem below does this for the odd cases.

Theorem 3. *As sets of knots $L(0, -4)$ and $\tilde{L}(1, -(2n - 1))$, for all $n > 0$, are subsets of $L(0, -1)$.*

Proof. Figure 8 shows that the knots in $L(0, -4)$ are in $L(0, -1)$. Figure 9 completes the proof. $\square$

FIGURE 8. $L(0, -4) \subset L(0, -1)$




FIGURE 9. Rest of Theorem 3

Acknowledgement: Supported in part by NSF grant no. DMS-9001973.

REFERENCES

- [1] M. Barge & M. Brown *Problems in Dynamics on Continua*, Contemporary Mathematics: Continuum Theory and Dynamical Systems, **117**, AMS, 1991.
- [2] J. Birman. *Braids, Links and Mapping Class Groups*, Annals of Math. Studies # 81, Princeton University Press, 1974.
- [3] J. Birman & R. Williams. Knotted Periodic Orbits in Dynamical Systems I: Lorenz Knots. *Topology* **22** (1983), 47-82.
- [4] J. Birman & R. Williams. Knotted Periodic Orbits in Dynamical Systems II: Knot Holders for Fibered Knots. *Contemporary Math.* **20** (1983).
- [5] G. Burde & H. Zieschang. *Knots*, Walter de Gruyter, 1985.
- [6] H. Schubert. Knotten und Vollringe. *Acta Math.* **90** (1953), 131-286.
- [7] R. Williams. Lorenz Knots are Prime. *Ergod. Th. & Dynam. Sys.* **4** (1983), 147-163.
- [8] R. Williams. Preprint: *The braid index of generalized cables*.