

DESCENT CONSTRUCTION FOR GSPIN GROUPS–ODD CASE

JOSEPH HUNDLEY AND EITAN SAYAG

CONTENTS

1. Introduction	1
1.1. Acknowledgements	2
2. The main result	2
3. Notation	4
3.1. General	4
3.2. Various Products	4
3.3. Similitude groups and GSpin groups	5
3.4. Weyl group of $GSpin_{2m}$; it's action on standard Levis and their representations	6
4. Preliminaries	9
4.1. Unramified Correspondence	9
4.2. Unipotent subgroups and their characters	9
4.3. “Unipotent periods”	10
5. Eisenstein series	11
6. Main Results	19
6.1. Descent Construction	19
7. Appendix I: Eisenstein series	23
7.1. Proof of Proposition 5.0.10	23
7.2. Proof of Lemma 5.0.12	24
7.3. Proof of Proposition 5.0.13	24
7.4. Proof of Proposition 5.0.15	27
7.5. Proof of 5.0.17	27
8. Appendix II: Local results on Jacquet Functors	28
9. Appendix III: Identities of Unipotent Periods	32
9.1. A Lemma Regarding Unipotent Periods	32
9.2. A lemma regarding the projection, and a remark	33
9.3. Relations among Unipotent Periods used in Theorem 6.1.4	33
References	38

ABSTRACT. In this paper we provide an extension of the theory of *descent* of Ginzburg-Rallis-Soudry to the context of essentially symplectic representations, that is representations τ with the property that the exterior square L -function twisted by some Hecke character ω has a pole at $s = 1$. Our theory supplements the recent work of Asgari-Shahidi on the functorial lift from the general Spin groups $GSpin_{2n+1}$ to GL_{2n} .

1. INTRODUCTION

The theory of descent for symplectic cuspidal representations of the general linear group $GL_{2n}(\mathbb{A})$ was developed in a sequence of remarkable works [GRS1]-[GRS5]. In these works the authors

Date: February 24, 2009.

Key words and phrases. Langlands functoriality, descent, unipotent integration.

constructed in an explicit way a space $\sigma(\pi)$ of cuspidal automorphic functions on $SO_{2n+1}(\mathbb{A})$ which weakly lifts to a cuspidal self-dual representation π of $GL_{2n}(\mathbb{A})$ with the property that $L(\pi, \wedge^2, s)$ has a pole at $s = 1$. In [C-K-PS-S2] the method of converse theorem is used to show the existence of a weak functorial lift from generic cuspidal automorphic representations of classical groups to automorphic representations of the general linear group. The combination of these methods allows the authors of [GRS4] to describe the image of the functorial lift of [C-K-PS-S1].

Thus, the conjunction of the descent method with the method of the converse theorem provides a very detailed description of the image of functoriality corresponding to the standard embedding of ${}^L G \rightarrow GL_N(\mathbb{C})$ with G a classical group. For an excellent survey we refer the reader to [So1].

Recently, Asgari and Shahidi proved in [Asg-Sha1] the existence of weak functorial lift from $GSpin$ groups to the general linear group. Later, in the special case of $GSp(4)$ they were able to show in [Asg-Sha2] that this weak functorial lift is in fact strong in an appropriate sense.

In this paper we extend the descent method of Ginzburg, Rallis, and Soudry to $GSpin$ groups. As a bonus, for $n \geq 2$ we can provide a “lower bound” on the image of the functorial lift from $GSpin_{2n+1}$ to GL_{2n} constructed by Asgari and Shahidi. For $n = 2$, these results were obtained by another method in [Gan-Tak].

Let us briefly review the method. For simplicity of the exposition we assume that we are trying to construct a descent for a cuspidal representation, τ .

We first relate the property of essential self-duality to the existence of a pole of an L -function of τ , and then construct an Eisenstein series with the L -function appearing in the constant term. In fact there are two possibilities for what the L -function is, and hence two possibilities for the structure of the Eisenstein series, and we only consider one in these notes. Our Eisenstein series will be defined on the group $GSpin_{4n}$ induced from a Levi M isomorphic to $GL_{2n} \times GL_1$. Now, a pole of the relevant L -function allows us to construct a residue representation $\mathcal{E}_{-1}(\tau, \omega)$ of $GSpin_{4n}$, associated to τ . Next, we give an embedding of $GSpin_{2n+1}$ into $GSpin_{4n}$, and construct, using formation of Fourier coefficient, a space of functions $DC_\omega(\tau)$ on this subgroup of $GSpin_{4n}$. We prove that $DC_\omega(\tau)$ is nonzero, and that all of the functions in it are cuspidal. It follows that it decomposes as a direct sum of irreducible automorphic cuspidal representations of $GSpin_{2n+1}$. We then show that each of these irreducible constituents lifts weakly to τ by the functorial lifting associated to the inclusion

$${}^L(GSpin_{2n+1}) = GSp_{2n}(\mathbb{C}) \hookrightarrow GL_{2n}(\mathbb{C}) = {}^L GL_{2n}.$$

In fact in these notes the representation τ may be an isobaric sum of several cuspidal representations τ_1, \dots, τ_r . The main differences are that the residue is a multi-residue, and the notation is more complicated.

1.1. Acknowledgements. The authors wish to thank the following people for helpful conversations: Dubravka Ban, William Banks, Daniel Bump, Wee Teck Gan, Hervé Jacquet, Erez Lapid, Omer Offen, Yiannis Sakellaridis, Gordan Savin, and Freydoon Shahidi. In addition, we wish to thank Mahdi Asgari, Jim Cogdell, Anantharam Raghuram and Freydoon Shahidi for their interest, which stimulated the work.

Without David Ginzburg and David Soudry’s many careful and patient explanations of the “classical” case— $\omega = 1$ —this work would not have been completed. It is important to point out that not all of the arguments shown to us have appeared in print. Nevertheless, in each case the specialization of our arguments to the case $\omega = 1$ may be correctly attributed to Ginzburg, Rallis, Soudry (with any errors or stylistic blemishes introduced being our own responsibility).

This work was undertaken while both authors were in Bonn at the Hausdorff Research Institute for Mathematics, in connection with a series of lectures of Professor Soudry’s. They wish to thank the Hausdorff Institute and Michael Rapoport for the opportunity. Finally, the second author

wishes to thank Prof. Erez Lapid for many enlightening discussions on the subject matter of these notes.

2. THE MAIN RESULT

Let $G = GSpin_{2n+1}$ and let $H = GL_{2n}$. Consider the inclusion

$${}^L G = {}^L (GSpin_{2n+1}) = GSp_{2n}(\mathbb{C}) \hookrightarrow GL_{2n}(\mathbb{C}) = {}^L GL_{2n} = {}^L H.$$

We denote this map r . Also, if $\pi \cong \otimes'_v \pi_v$ is an automorphic representation of a group $G'(\mathbb{A})$, where \mathbb{A} is the ring of adèles of a number field F , then the semisimple conjugacy class in the L -group ${}^L G'$ associated to the local representation π_v at an unramified place v will be denoted t_{π_v} . We say that an automorphic representation σ of $G(\mathbb{A})$ is a weak lift of the automorphic representation τ of $H(\mathbb{A})$ if for almost all places, $r(t_{\sigma_v}) = t_{\tau_v}$.

To formulate our main result we introduce the notion of η symplectic representations. Let τ be an irreducible automorphic cuspidal representation of GL_{2n} . Suppose that τ is essentially self-dual, i.e. that the contragredient $\tilde{\tau}$ of τ is isomorphic to $\tau \otimes \eta$ for some Hecke character η . It follows from [Ja-Sh2] (see remark (4.11) pp. 553-54) that $L(s, \tau \times \tau \otimes \eta)$ has a simple pole at $s = 1$. Now, $L(s, \tau \times \tau \otimes \eta)$ is the Langlands L function of the representation $\tau \boxtimes \eta$ (exterior tensor product) of the group $GL_{2n}(\mathbb{A}) \times GL_1(\mathbb{A})$ associated to the representation of the L group $GL_{2n}(\mathbb{C}) \times GL_1(\mathbb{C})$ (finite Galois form) on $M_{2n \times 2n}(\mathbb{C})$ in which $GL_{2n}(\mathbb{C})$ acts by $g \cdot X = gX {}^t g$ and $GL_1(\mathbb{C})$ acts by scalar multiplication. But this representation is reducible, decomposing into the subspaces of skew-symmetric and symmetric matrices. We denote the associated L functions $L(s, \tau, \wedge^2 \times \eta)$ and $L(s, \tau, sym^2 \times \eta)$ respectively. The local factors at finite ramified places may be defined using the local Langlands classification ([L2],[H-T],[Henn1]) and the definition of an Artin L function attached to a finite dimensional representation of the Weil group [Tate1], or they may be defined as in [Sha2]. By [Henn2] these two definitions agree. Then we have

$$L(s, \tau \times \tau \otimes \eta) = L(s, \tau, \wedge^2 \times \eta) L(s, \tau, sym^2 \times \eta).$$

As both of the L functions on the right-hand side are obtainable via the Langlands-Shahidi method, neither may vanish at $s = 1$ (see [Gel-Sha] §2.6 p. 84). Thus, exactly one of these two L functions has a simple pole at $s = 1$ while the other is holomorphic and nonvanishing. Similarly, if $\tilde{\tau}$ is not isomorphic to $\tau \otimes \eta$ then they are both holomorphic at $s = 1$. (This requires the extension of [Ja-Sh2] remark (4.11) to *completed* L functions— i.e., the statement that none of the local L functions has a pole at $s = 1$. The requisite facts about local L functions are well-known and a proof is reviewed at the end of Theorem 5.0.4.) One may prove the second assertion using results of Langlands via the method explained on p. 840 of [Kim].

We will say that τ is η -symplectic in case $L(s, \tau, \wedge^2 \times \eta)$ has a pole at $s = 1$ and η -orthogonal otherwise. We also define “almost symplectic” to mean “ η -symplectic for some η ,” and “almost orthogonal” similarly.

- Remarks 2.0.1.** (1) *There is another natural notion of “orthogonal/symplectic representation.” Specifically, one could say that an automorphic representation is orthogonal/symplectic if the space it acts on supports an invariant symmetric/skew-symmetric form. The two notions appear to be related, but do not coincide. See [PraRam].*
- (2) *There is a third approach to defining a local factor for $L(s, \tau, \wedge^2 \times \eta)$, which is to apply the “gcd” construction described in [Gel-Sha] section I.1.6, p. 17, to the integrals in [Ja-Sh1]. As far as we know this is not written down anywhere.*
- (3) *An integral representation for $L(s, \tau, sym^2)$ was given in [BG]. The problem of extending this to $L(s, \tau, sym^2 \times \eta)$ has been considered by Banks [Banks1, Banks2]. Nontrivial technical difficulties arise, particularly in the case we consider, when τ is defined on GL_{2n} [Banks3].*

- (4) Let AS denote the functorial lift constructed in [Asg-Sha1]. It is shown in [Asg-Sha1] that $AS(\pi)$ is nearly equivalent to $\widetilde{AS(\pi)} \otimes \omega_\pi$, where ω_π denotes the central character of the representation π . (Of course, this means that they are the same space of functions when $AS(\pi)$ is cuspidal.) Thus, in practice it turns out to make sense to use $\eta = \omega^{-1}(= \bar{\omega})$.

By proposition 2 of [L3], every irreducible automorphic representation of $GL_n(\mathbb{A})$ is isomorphic to a subquotient of $Ind_{P(\mathbb{A})}^{GL_n(\mathbb{A})} \tau_1 |\det_1|^{s_1} \otimes \cdots \otimes \tau_r |\det_r|^{s_r}$ for some real numbers s_1, \dots, s_r and irreducible unitary automorphic cuspidal representations τ_1, \dots, τ_r of $GL_{n_1}(\mathbb{A}), \dots, GL_{n_r}(\mathbb{A})$ respectively, such that $n_1 + \cdots + n_r = n$. Here P is the standard parabolic of GL_n corresponding to the ordered partition (n_1, \dots, n_r) of n .

In the case when $s_i = 0$ for all i , this induced representation is irreducible. (This follows from the irreducibility of all the local induced representations, which is Theorem 3.2 of [Ja].)

Also, the representations obtained by numbering a given set of cuspidal representations in different ways are isomorphic. (This follows from the fact that the standard intertwining operator between them does not vanish, which follows from [MW1], II.1.8 (meromorphically continued in IV.1.9(e)), and IV.1.10(b). In IV.3.12 these elements are combined to prove that the intertwining operator does not have a pole. The proof that it does not have a zero is an easy adaptation.) Furthermore, if two such induced representations are isomorphic, then they are obtained from two numberings of the same set of cuspidal representations ([Ja-Sh3], Theorem 4.4, p.809). An irreducible unitary representation τ of $GL_n(\mathbb{A})$ which is obtained from irreducible unitary cuspidal representations τ_1, \dots, τ_r in this manner is sometimes called the *isobaric sum* of the cuspiduals, and denoted $\tau_1 \boxplus \cdots \boxplus \tau_r$. (A more general notion of “isobaric representation” was introduced in [L4], but we don’t need it.)

Theorem 2.1. *For $r \in \mathbb{N}$, take τ_1, \dots, τ_r to be irreducible unitary automorphic cuspidal representations of $GL_{2n_1}(\mathbb{A}), \dots, GL_{2n_r}(\mathbb{A})$, respectively, and let $\tau = \tau_1 \boxplus \cdots \boxplus \tau_r$. Let ω denote a Hecke character. Suppose that*

- τ_i is $\bar{\omega}$ -symplectic for each i , and
- $\tau_i \cong \tau_j \Rightarrow i = j$.

Then there exists an irreducible generic cuspidal automorphic representation σ of $GSpin_{2n+1}(\mathbb{A})$ such that

- σ weakly lifts to τ , and
- the central character ω_σ of σ is ω .

Remark 2.0.2. *The case $n = 1$ is trivial because $GSpin_3 = GSp_2 = GL_2$, so the inclusion r is simply the identity map. Clearly, r must be one and $\sigma = \tau_1$. Henceforth, we assume $n \geq 2$. The careful reader will find places where this assumption is crucial to the validity of the argument.*

Corollary 2.2. *The image of the functorial lift AS described in Theorem 1.1 (p. 140) of [Asg-Sha1] contains the set of all representations $\tau_1 \boxplus \cdots \boxplus \tau_r$ such that*

- $\tau_i \cong \tau_j \Rightarrow i = j$,
- there is a Hecke character ω such that τ_i is $\bar{\omega}$ -symplectic for each i .

3. NOTATION

3.1. General. Throughout most of the paper, F will denote a number field. In Appendix II, it will be a non-Archimedean local field of characteristic zero.

We denote by J the matrix, of any size, with ones on the diagonal running from upper right to lower left, and by J' the matrix $(-J^J)$ of any even size. We also employ the notation ${}^t g$ for the transpose of g and ${}_t g$ for the “other transpose” $J {}^t g J$. We employ the shorthand $G(F \backslash \mathbb{A}) = G(F) \backslash G(\mathbb{A})$, where G is any F -group.

We denote the Weyl group of the reductive group G by W_G or by W , when the meaning is clear from context.

If π is an automorphic or local representation, then $\tilde{\pi}$ is the contragredient, and ω_π the central character.

3.2. Various Products. Most tensor products will be denoted \otimes . However \boxtimes will sometimes be used to distinguish the “outer” tensor product from the “inner” tensor products and “twisting.” Let us recall these notions.

If (π_1, V_1) and (π_2, V_2) are representations of groups G_1 and G_2 , then one may consider the representation of $G_1 \times G_2$ on $V_1 \otimes V_2$ given on pure tensors by

$$(\pi_1 \otimes \pi_2)(g_1, g_2)v_1 \otimes v_2 = \pi_1(g_1)v_1 \otimes \pi_2(g_2)v_2.$$

If (π_1, V_1) and (π_2, V_2) happen to be two representations of the *same* group G , then this construction yields a representation of $G \times G$. The space $V_1 \otimes V_2$ also supports a natural “tensor product representation” of the group G itself with the action given on pure tensors by

$$(\pi_1 \otimes \pi_2)(g)v_1 \otimes v_2 = \pi_1(g)v_1 \otimes \pi_2(g)v_2.$$

The representation of $G \times G$ on $V_1 \otimes V_2$ is sometimes called the outer tensor product and denoted \boxtimes to avoid ambiguity.

Adding to the mix, the twist of a representation π of $GL_n(\mathbb{A})$ by a Hecke character χ is often denoted $\pi \otimes \chi$. In terms of the constructions above, it is the inner tensor product of π and the representation of $GL_n(\mathbb{A})$ obtained by composing χ with \det . We shall keep to this notation. We shall also need to consider the (outer) tensor product representation of $GL_n(\mathbb{A}) \times GL_1(\mathbb{A})$, for which we employ \boxtimes .

Let us mention that \boxtimes will not be used in the sense of [L4].

In addition to \otimes and \boxtimes , we use \boxplus for “isobaric sum” as described above. We use \times for Cartesian product of sets, groups, etc., and in the notation for various L functions (e.g., $\wedge^2 \times \bar{\omega}$).

3.3. Similitude groups and GSpin groups. We first define the *similitude orthogonal and symplectic groups* to be

$$GO_m = \{g \in GL_m : gJ^t g = \lambda(g)J \text{ for some } \lambda(g) \in \mathbb{G}_m\},$$

$$GSp_{2n} = \{g \in GL_{2n} : gJ'^t g = \lambda(g)J' \text{ for some } \lambda(g) \in \mathbb{G}_m\}.$$

For each of these groups the map $g \mapsto \lambda(g)$ is a rational character called the *similitude factor*. If m is odd then GO_m is in fact isomorphic to $SO_m \times GL_1$. This case will play no further role. The group GO_{2n} is disconnected; indeed the subgroup generated by SO_{2n} and $\left\{ \begin{pmatrix} \lambda I_n & \\ & I_n \end{pmatrix} : \lambda \in \mathbb{G}_m \right\}$ is a connected index two subgroup, which we denote GSO_{2n} .

We shall now define GSpin groups as the groups whose duals are the similitude classical groups $GSp_{2n}(\mathbb{C}), GSO_{2n}(\mathbb{C})$. Thus we write down the based root data, but employ notation appropriate to the application in which what we write down will arise as the dual of something.

The groups GSp_{2n} and GSO_{2n} share a maximal torus, consisting of matrices of the form

$$\text{diag}(t_1, \dots, t_n, \lambda t_n^{-1}, \dots, \lambda t_1^{-1}).$$

The coordinates used just above correspond to a choice of \mathbb{Z} -bases for the lattices of characters and cocharacters. For $i = 1$ to n , let e_i^* denote the character that sends this torus element to t_i for $i = 1$ to n and e_0^* being the map that sends it to the similitude factor, λ . Let $\{e_i : i = 0 \text{ to } n\}$ denote the dual basis for the cocharacter lattice. Let X^\vee denote the character lattice and X the cocharacter lattice. Each similitude classical group has a Borel subgroup equal to the set of upper triangular

matrices which are in it. In each case we employ this choice of Borel, and let Δ^\vee denote the set of simple roots and Δ the set of simple coroots. Then we easily compute that for GSp_{2n}

$$\Delta^\vee = \{e_i^* - e_{i+1}^*, i = 1 \text{ to } n - 1\} \cup \{2e_n^* - e_0^*\}.$$

$$\Delta = \{e_i - e_{i+1}, i = 1 \text{ to } n - 1\} \cup \{e_n\}.$$

and for GSO_{2n}

$$\Delta^\vee = \{e_i^* - e_{i+1}^*, i = 1 \text{ to } n - 1\} \cup \{e_{n-1}^* + e_n^* - e_0^*\}.$$

$$\Delta = \{e_i - e_{i+1}, i = 1 \text{ to } n - 1\} \cup \{e_{n-1} + e_n\}.$$

We now define $GSpin_{2n+1}$ to be the F -split connected reductive algebraic group having based root datum dual to that of GSp_{2n} , and $GSpin_{2n}$ to be the one having based root datum dual to that of GSO_{2n} . We have here used the fact that F -split connected reductive algebraic groups are classified by based root data, for which see p.274 of [Spr].

To save space, the group $GSpin_m$ will usually be denoted G_m .

Observe that in either the odd or even case e_0^* is a generator for the lattice of cocharacters of the center of G_m .

Because we define G_m in the manner we do, it comes equipped with a choice of Borel subgroup and maximal torus, as do various reductive subgroups we shall consider below. In each case, we denote the Borel subgroup of the reductive group G by $B(G)$, and the maximal torus by $T(G)$.

A straightforward adaptation of the proof of Theorem 16.3.2 of [Spr] shows that there exist surjections $G_m \rightarrow SO_m$ defined over F . We fix one such and denote it pr . We require that pr is such that $\text{pr}(B(G_m))$ consists of upper triangular matrices.

An alternative description of the same group as a quotient of $Spin_m \times GL_1$ is given in [Asg]. Proposition 2.4 on p. 678 of [Asg] shows that the two definitions are equivalent.

For those familiar with the construction of $Spin_m$ as a subgroup of the multiplicative group of a Clifford algebra, we remark that there is a third construction of $GSpin_m$ as the slightly larger group obtained by including the nonzero scalars in the Clifford algebra as well. In this guise, it is sometimes referred to as the ‘‘Clifford group.’’ (See, e.g., [I] p.999.) This description will not play a role for us.

We will construct an Eisenstein series on G_{2m} induced from a standard parabolic $P = MU$ such that M is isomorphic to $GL_m \times GL_1$. There are two such parabolics. We choose the one in which we delete the root $e_{m-1} + e_m$ and the coroot $e_{m-1}^* + e_m^* - e_0^*$ from the based root datum. We shall refer to this parabolic as the ‘‘Siegel.’’

Remark 3.3.1.

- We can identify the based root datum of the Levi M with that of $GL_m \times GL_1$ in such a fashion that e_0 corresponds to GL_1 and does not appear at all in GL_m . We can then identify M itself with $GL_m \times GL_1$ via a particular choice of isomorphism which is compatible this and with the usual usage of e_i, e_i^* for characters, cocharacters of the standard torus of GL_m .
- Having made this identification, a Levi M' which is contained in M will be identified with $GL_1 \times GL_{m_1} \times \dots \times GL_{m_k}$, (for some $m_1, \dots, m_k \in \mathbb{N}$ that add up to m) in the natural way: GL_1 is identified with the GL_1 factor of M , and then $GL_{m_1} \times \dots \times GL_{m_k}$ is identified with the subgroup of M corresponding to block diagonal elements with the specified block sizes, in the specified order.
- The lattice of rational characters of M is spanned by the maps $(g, \alpha) \mapsto \alpha$ and $(g, \alpha) \mapsto \det g$. Restriction defines an embedding $X(M) \rightarrow X(T)$, which sends these maps to e_0 and $(e_1 + \dots + e_m)$, respectively. By abuse of notation, we shall refer to the rational character of M corresponding to e_0 as e_0 as well.
- The modulus of P is $(g, \alpha) \rightarrow \det g^{(m-1)}$.

The group G_{2n} has an involution \dagger which reverses the last two simple roots. The effect is such that

$$\text{pr}(\dagger g) = \begin{pmatrix} I_{n-1} & & & \\ & 1 & & \\ & & 1 & \\ & & & I_{n-1} \end{pmatrix} \text{pr}(g) \begin{pmatrix} I_{n-1} & & & \\ & 1 & & \\ & & 1 & \\ & & & I_{n-1} \end{pmatrix}.$$

As is well known, there is a group $Pin_{4n} \supset Spin_{4n}$ such that pr extends to a two-fold covering $Pin_{4n} \rightarrow O_{4n}$. The involution \dagger can be realized as conjugation by a preimage of the above permutation matrix.

We also fix a maximal compact subgroup K_m of $G_m(\mathbb{A})$. Any which satisfies the conditions required in [MW1] (see pages 1 and 4) will do.

3.4. Weyl group of $GSpin_{2m}$; it's action on standard Levis and their representations.

Lemma 3.4.1. *The Weyl group of G_m is canonically identified with that of SO_m .*

Proof. For this lemma only, let T denote the torus of SO_m and \tilde{T} that of G_m . Then the following diagram commutes:

$$\begin{array}{ccc} Z_{G_m}(\tilde{T}) & \longrightarrow & N_{G_m}(\tilde{T}) \\ \downarrow & & \downarrow \\ Z_{SO_m}(T) & \longrightarrow & N_{SO_m}(T). \end{array}$$

Both horizontal arrows are inclusions and both vertical arrows are pr . □

One easily checks that every element of the Weyl group of SO_{2n} is represented by a permutation matrix. We denote the permutation associated to w also by w . The set of permutations w obtained is precisely the set of permutations $w \in \mathfrak{S}_{2n}$ satisfying,

- (1) $w(2n + 1 - i) = 2n + 1 - w(i)$ and
- (2) $\det w = 1$ when w is written as a $2n \times 2n$ permutation matrix.

It is well known that the Weyl group of SO_{2n} (or G_{2n}) is isomorphic to $\mathfrak{S}_n \times \{\pm 1\}^{n-1}$. To fix a concrete isomorphism, we identify $p \in \mathfrak{S}_n$ with an $n \times n$ matrix in the usual way, and then with

$$\begin{pmatrix} p & \\ & {}_t p^{-1} \end{pmatrix} \in SO_{2n}.$$

We identify $\underline{\epsilon} = (\epsilon_1, \dots, \epsilon_{n-1}) \in \{\pm 1\}^{n-1}$ with the permutation p of $\{1, \dots, 2n\}$ such that

$$p(i) = \begin{cases} i & \text{if } \epsilon_i = 1 \\ 2n + 1 - i & \text{if } \epsilon_i = -1, \end{cases}$$

where ϵ_n is defined to be $\prod_{i=1}^{n-1} \epsilon_i$. We then identify $(p, \underline{\epsilon}) \in \mathfrak{S}_n \times \{\pm 1\}^{n-1}$ (direct product of sets) with $p \cdot \underline{\epsilon} \in W_{SO_{2n}}$.

With this identification made,
(3.4.2)

$$(p, \underline{\epsilon}) \cdot \begin{pmatrix} t_1 & & & & & \\ & \ddots & & & & \\ & & t_n & & & \\ & & & t_n^{-1} & & \\ & & & & \ddots & \\ & & & & & t_1^{-1} \end{pmatrix} \cdot (p, \underline{\epsilon})^{-1} = \begin{pmatrix} t_{p^{-1}(1)}^{\epsilon_{p^{-1}(1)}} & & & & & \\ & \ddots & & & & \\ & & t_{p^{-1}(n)}^{\epsilon_{p^{-1}(n)}} & & & \\ & & & t_{p^{-1}(n)}^{-\epsilon_{p^{-1}(n)}} & & \\ & & & & \ddots & \\ & & & & & t_{p^{-1}(1)}^{-\epsilon_{p^{-1}(1)}} \end{pmatrix}.$$

Lemma 3.4.3. *Let $(p, \underline{\epsilon}) \in \mathfrak{S}_n \times \{\pm 1\}^{n-1}$ be identified with an element of $W_{SO_{2m}} = W_{G_{2m}}$ as above. Then the action on the character and cocharacter lattices of G_{2m} is given as follows:*

$$(p, \underline{\epsilon}) \cdot e_i = \begin{cases} e_{p(i)} & i > 0, \epsilon_{p(i)} = 1, \\ -e_{p(i)} & i > 0, \epsilon_{p(i)} = -1, \\ e_0 + \sum_{\epsilon_{p(i)} = -1} e_{p(i)} & i = 0. \end{cases}$$

$$(p, \underline{\epsilon}) \cdot e_i^* = \begin{cases} e_{p(i)}^* & i > 0, \epsilon_{p(i)} = 1, \\ e_0^* - e_{p(i)}^* & i > 0, \epsilon_{p(i)} = -1, \\ e_0^* & i = 0. \end{cases}$$

Remark 3.4.4. *Much of this can be deduced from (3.4.2), keeping in mind that $w \in W_G$ acts on cocharacters by $(w \cdot \varphi)(t) = w\varphi(t)w^{-1}$ and on characters by $(w \cdot \chi)(t) = \chi(w^{-1}tw)$. However, it is more convenient to give a different proof.*

Proof. Let $\alpha_i = e_i - e_{i+1}$, $i = 1$ to $n-1$ and $\alpha_n = e_{n-1} + e_n$. Let s_i denote the elementary reflection in $W_{G_{2n}}$ corresponding to α_i . Then it is easily verified that s_1, \dots, s_{n-1} generate a group isomorphic to \mathfrak{S}_n which acts on $\{e_1, \dots, e_n\} \in X(T)$ and $\{e_1^*, \dots, e_n^*\} \in X^\vee(T)$ by permuting the indices and acts trivially on e_0 and e_0^* . Also

$$s_n \cdot e_i = \begin{cases} e_i & i \neq n-1, n, 0 \\ e_0 + e_n + e_{n-1} & i = 0 \\ -e_n & i = n-1 \\ -e_{n-1} & i = n \end{cases}$$

$$s_n \cdot e_i^* = \begin{cases} e_i^* & i \neq n-1, n \\ e_0^* - e_n^* & i = n-1 \\ e_0^* - e_{n-1}^* & i = n. \end{cases}$$

If $\underline{\epsilon} \in \{\pm 1\}^{n-1}$ is such that $\#\{i : \epsilon_i = -1\} = 1$ or 2 , then $\underline{\epsilon}$ is conjugate to s_n by an element of the subgroup isomorphic to \mathfrak{S}_n generated by s_1, \dots, s_{n-1} . An arbitrary element of $\{\pm 1\}^{n-1}$ is a product of elements of this form, so one is able to deduce the assertion for general $(p, \underline{\epsilon})$. \square

Observe that the \mathfrak{S}_n factor in the semidirect product is precisely the Weyl group of the Siegel Levi.

In the study of intertwining operators and Eisenstein series (e.g., section 5 below), one encounters a certain subset of the Weyl group associated to a standard Levi, M . Specifically,

$$W(M) := \left\{ w \in W_{G_{2n}} \mid \begin{array}{l} w \text{ is of minimal length in } w \cdot W_M \\ wMw^{-1} \text{ is a standard Levi of } G_{2n} \end{array} \right\}.$$

For our purposes, it is enough to consider the case when M is a subgroup of the Siegel Levi. In this case it is isomorphic to $GL_{m_1} \times \cdots \times GL_{m_r} \times GL_1$ for some integers m_1, \dots, m_r which add up to n , and we shall only need to consider the case when m_i is even for each i . (This, of course, forces n to be even as well.)

Lemma 3.4.5. *For each $w \in W(M)$ with M as above, there exist a permutation $p \in \mathfrak{S}_r$ and an element $\underline{\epsilon} \in \{\pm 1\}^r$ such that, if $m \in M = (g, \alpha)$ with $\alpha \in GL_1$ and*

$$g = \begin{pmatrix} g_1 & & \\ & \ddots & \\ & & g_r \end{pmatrix} \in GL_n,$$

then

$$wmw^{-1} = (g', \alpha \cdot \prod_{\epsilon_i = -1} \det g_i) \quad g' = \begin{pmatrix} g'_1 & & \\ & \ddots & \\ & & g'_r \end{pmatrix},$$

where

$$g'_i \approx \begin{cases} g_{p^{-1}(i)} & \text{if } \epsilon_{p^{-1}(i)} = 1, \\ t g_{p^{-1}(i)}^{-1} & \text{if } \epsilon_{p^{-1}(i)} = -1. \end{cases}$$

Here \approx has been used to denote equality up to an inner automorphism. The map $(p, \underline{\epsilon}) \mapsto w$ is a bijection between $W(M)$ and $\mathfrak{S}_r \times \{\pm 1\}^r$. (Direct product of sets: $W(M)$ is not, in general, a group.)

Proof. We first prove that wMw^{-1} is again contained in the Siegel Levi.

The Levi M determines an equivalence relation \sim on the set of indices, $\{1, \dots, n\}$ defined by the condition that $i \sim i+1$ iff $e_i - e_{i+1}$ is a root of M . View w of $W(M)$ as a permutation of $\{1, \dots, 2n\}$. Because w is of minimal length, $i \sim j$, $i < j \Rightarrow w(i) < w(j)$. Because wMw^{-1} is a standard Levi, we may deduce that if $i \sim i+1$ then $w(i+1) = w(i) + 1$, except possibly when $w(i) = n-1$, in which case $w(i+1)$ could, *a priori* be $n+1$. However, it is easy to check that in the special case when all m_i are even, the condition $\det w = 1$ forces $w(i+1) = w(i) + 1$ even if $w(i) = n-1$. It follows that wMw^{-1} is contained in the Siegel Levi.

When viewed as elements of $\mathfrak{S}_n \times \{\pm 1\}^{n-1}$, the elements of $W(M)$ are those pairs $(p, \underline{\epsilon})$ such that $i \sim i+1 \Rightarrow p(i+1) = p(i) + 1$, and $i \sim j \Rightarrow \epsilon_i = \epsilon_j$. This gives the identification with $\mathfrak{S}_r \times \{\pm 1\}^r$.

It is clear that the precise value of g'_i is determined only up to conjugacy by an element of the torus (because we do not specify a particular representative for our Weyl group element). By Theorem 16.3.2 of [Spr], it may be discerned, to this level of precision, by looking at the effect of w on the based root datum of M . The result now follows from Lemma 3.4.3. \square

Corollary 3.4.6. *Let $w \in W(M)$ be associated to $(p, \underline{\epsilon}) \in \mathfrak{S}_r \times \{\pm 1\}^r$ as above. Let τ_1, \dots, τ_r be irreducible cuspidal representations of $GL_{m_1}(\mathbb{A}), \dots, GL_{m_r}(\mathbb{A})$, respectively, and let ω be a Hecke character. Then our identification of M with $GL_{m_1} \times \cdots \times GL_{m_r} \times GL_1$ determines an identification of $\bigotimes_{i=1}^r \tau_i \boxtimes \omega$ with a representation of $M(\mathbb{A})$. Let $M' = wMw^{-1}$. Then M' is also identified, via 3.3.1 with $GL_{m_{p^{-1}(1)}} \times \cdots \times GL_{m_{p^{-1}(r)}} \times GL_1$, and we have*

$$\bigotimes_{i=1}^r \tau_i \boxtimes \omega \circ Ad(w^{-1}) = \bigotimes_{i=1}^r \tau'_i \boxtimes \omega,$$

where

$$\tau'_i = \begin{cases} \tau_{p^{-1}(i)} & \text{if } \epsilon_{p^{-1}(i)} = 1, \\ \tilde{\tau}_{p^{-1}(i)} \otimes \omega & \text{if } \epsilon_{p^{-1}(i)} = -1. \end{cases}$$

Proof. The contragredient $\tilde{\tau}_i$ of τ_i may be realized as an action on the same space of functions as τ_i via $g \cdot \varphi(g_1) = \varphi(g_1 {}_t g^{-1})$. This follows from strong multiplicity one and the analogous statement for local representations, for which see [GK75] page 96, or [BZ1] page 57. Combining this fact with the Lemma, we obtain the Corollary. \square

4. PRELIMINARIES

4.1. Unramified Correspondence.

Lemma 4.1.1. *Suppose that $\tau \cong \otimes'_v \tau_v$ is an $\bar{\omega}$ -symplectic irreducible cuspidal automorphic representation of $GL_{2n}(\mathbb{A})$. Let v be a place such that τ_v is unramified. Let $t_{\tau,v}$ denote the semisimple conjugacy class in $GL_{2n}(\mathbb{C})$ associated to τ_v . Let $r : GSp_{2n}(\mathbb{C}) \rightarrow GL_{2n}(\mathbb{C})$ be the natural inclusion. Then $t_{\tau,v}$ contains elements of the image of r .*

Proof. For convenience in the application, we take GL_{2n} to be identified with a subgroup of the Levi of the Siegel parabolic as in section 3.3. Since τ_v is both unramified and generic, it is isomorphic to $\text{Ind}_{B(GL_{2n})(F_v)}^{GL_{2n}(F_v)} \mu$ for some unramified character μ of the maximal torus $T(GL_{2n})(F_v)$ such that this induced representation is irreducible. (See [Car], section 4, [Z] Theorem 8.1, p. 195.) Let $\mu_i = \mu \circ e_i^*$.

Since $\tau \cong \tilde{\tau} \otimes \omega$, it follows that $\tau_v \cong \tilde{\tau}_v \otimes \omega_v$ and from this we deduce that $\{\mu_i : 1 \leq i \leq 2n\}$ and $\{\mu_i^{-1} \omega : 1 \leq i \leq 2n\}$ are the same set.

By Theorem 1, p. 213 of [Ja-Sh1], we have $\prod_{i=1}^{2n} \mu_i = \omega^n$.

Now, what we need to prove is the following: if S is a set of $2n$ unramified characters of F_v , such that

- (1) $\prod_{i=1}^{2n} \mu_i = \omega^n$
- (2) For each i there exists j such that $\mu_i = \mu_j^{-1} \omega$

then there is a permutation $\sigma : \{1, \dots, 2n\} \rightarrow \{1, \dots, 2n\}$ such that $\mu_{\sigma(i)} = \omega \mu_{2n-\sigma(i)}^{-1}$ for $i = 1$ to n . This we prove by induction on n . When $n = 1$, we know that $\mu_1 = \mu_i^{-1} \omega$ for $i = 1$ or 2 . If $i = 2$ we are done, while if $i = 1$ we use $\omega = \mu_1 \mu_2$ to obtain $\mu_1 = \mu_2$, and the desired assertion. Now, if $n > 1$ it is sufficient to show that there exist $i \neq j$ such that $\mu_i = \mu_j^{-1} \omega$. If there exists i such that $\mu_i \neq \mu_i^{-1} \omega$ then this is clear. On the other hand, there are exactly two unramified characters μ such that $\mu = \mu^{-1} \omega$. The result follows \square

Corollary 4.1.2. *Suppose $\tau = \tau_1 \boxplus \dots \boxplus \tau_r$ with τ_i an $\bar{\omega}$ -symplectic irreducible cuspidal automorphic representation of $GL_{2n_i}(\mathbb{A})$, for each i . Then the same conclusion holds.*

4.2. Unipotent subgroups and their characters. The kernel of pr consists of semisimple elements. In particular, the number of unipotent elements of a fiber is zero or one, and it's one if and only if the element of SO_m is unipotent. In other words, pr yields a bijection of unipotent elements (indeed, an isomorphism of unipotent subvarieties), and we may specify unipotent elements or subgroups by their images under pr . This also defines coordinates for any unipotent element or subgroup, which we use when defining characters. Thus, we write u_{ij} for the i, j entry of $\text{pr}(u)$.

Above we fixed a specific isomorphism of a subgroup of G_{2m} with GL_m . If u is a unipotent element of of this subgroup this identification with an $m \times m$ matrix gives a second definition of u_{ij} . This is not a problem, however, as the two definitions agree.

Most of the unipotent groups we consider are subgroups of the maximal unipotent of G_m consisting of elements u with $\text{pr}(u)$ upper triangular. We denote this group U_{\max} . A complete set of coordinates is $\{u_{ij} : 1 \leq i < j \leq m - i\}$. We denote the opposite maximal unipotent by $\overline{U_{\max}}$. It consists of all unipotent elements of G_m such that $\text{pr}(u)$ is lower triangular.

We fix once and for all a character ψ_0 of \mathbb{A}/F . We use this character together with the coordinates just above to specify characters of our unipotent subgroups.

When specifying subgroups of U_{\max} and their characters, the restriction to $\{(i, j) : 1 \leq i < j \leq m - i\}$ is implicit.

It will also sometimes be necessary to describe unipotent subgroups such that only a few of the entries in the corresponding elements of SO_m are nonzero. For this purpose we introduce the notation $e'_{ij} = e_{ij} - e_{m+1-j, m+1-i}$. One may check that for all $i \neq j$ and $a \in F$, the matrix $I + ae'_{ij}$ is an element of $SO_m(F)$.

4.3. “Unipotent periods”. We now introduce the framework within which, we believe, certain of the computations involved in the descent construction can be most easily understood.

Let G be a reductive algebraic group defined over a number field F . If U is a unipotent subgroup of G and ψ_U is a character of $U(F \backslash \mathbb{A})$, we define the *unipotent period* (U, ψ_U) associated to this pair to be given by the formula

$$\varphi^{(U, \psi_U)}(g) := \int_{U(F \backslash \mathbb{A})} \varphi(ug) \psi_U(u) du.$$

Clearly, φ must be restricted to a space of left $U(F)$ -invariant functions such that the integral is defined (for example, because φ is smooth).

Let \mathcal{U} denote the set of unipotent periods. For V a space of functions defined on $G(\mathbb{A})$, put

$$\mathcal{U}^\perp(V) = \{(U, \psi) \in \mathcal{U} : \varphi^{(U, \psi)} \equiv 0 \ \forall \varphi \in V\}.$$

When V is the space of a representation π we will employ also the notation $\mathcal{U}^\perp(\pi)$. We also employ the notation $(U, \psi) \perp V$ for $(U, \psi) \in \mathcal{U}^\perp(V)$ and similarly $(U, \psi) \perp \pi$.

We also require a vocabulary to express relationships among unipotent periods. We shall say that

$$(U, \psi_U) \in \langle (U_1, \psi_{U_1}), \dots, (U_n, \psi_{U_n}) \rangle$$

if $V \perp (U_i, \psi_{U_i}) \forall i \Rightarrow V \perp (U, \psi_U)$. Clearly, if $(U_1, \psi_{U_1}) \in \langle (U_2, \psi_2), (U_3, \psi_3) \rangle$, and $(U_2, \psi_2) \in \langle (U_4, \psi_4), (U_5, \psi_5) \rangle$ then $(U_1, \psi_1) \in \langle (U_3, \psi_3), (U_4, \psi_4), (U_5, \psi_5) \rangle$.

We also use notation $(U_1, \psi_1) | (U_2, \psi_2)$, or the language “ (U_1, ψ_1) divides (U_2, ψ_2) ,” “ (U_2, ψ_2) is divisible by (U_1, ψ_1) ” for $(U_2, \psi_2) \in \langle (U_1, \psi_1) \rangle$. Finally, $(U_1, \psi_1) \sim (U_2, \psi_2)$ means $(U_1, \psi_1) | (U_2, \psi_2)$ and $(U_2, \psi_2) | (U_1, \psi_1)$. This is an equivalence relation and we shall refer to unipotent periods which are related in this way as “equivalent.”

It is *sometimes* possible to compose unipotent periods. Specifically, if $f^{(U_1, \psi_1)}$ is left-invariant by $U_2(F)$, then one may consider $(f^{(U_1, \psi_1)})^{(U_2, \psi_2)}$. We denote the composite by $(U_2, \psi_2) \circ (U_1, \psi_1)$.

Now, suppose that U is the unipotent radical of a parabolic P of G with Levi M . The choice of ψ_0 gives rise to an identification of the space of characters of $U(F) \backslash U(\mathbb{A})$ with the F points of $\overline{U}/(\overline{U}, \overline{U})$ which is compatible with the action of $M(F)$. Here \overline{U} denotes the unipotent radical of the parabolic \overline{P} of G opposite to P . For ϑ a character, let M^ϑ denote the stabilizer of ϑ (regarded as a point in $\overline{U}/(\overline{U}, \overline{U})(F)$) in M . So M^ϑ is an algebraic subgroup of M defined over F .

Definition 4.3.1. Then we define $FC^\vartheta : C^\infty(G(F \backslash \mathbb{A})) \rightarrow C^\infty(M^\vartheta(F \backslash \mathbb{A}))$ by

$$FC^\vartheta(\varphi)(m) = \varphi^{(U, \vartheta)}(m) = \int_{U(F \backslash \mathbb{A})} \varphi(um) \vartheta(u) du.$$

This is certainly an $M^\vartheta(\mathbb{A})$ -equivariant map.

5. EISENSTEIN SERIES

The main purpose of this section is to construct, for each integer $n \geq 2$ and Hecke character ω , a map from the set of all isobaric representations τ satisfying the hypotheses of theorem 2.1 into the residual spectrum of G_{4n} . We use the same notation $\mathcal{E}_{-1}(\tau, \omega)$ for all n . The construction is given by a multi-residue of an Eisenstein series in several complex variables, induced from the cuspidal

representations τ_1, \dots, τ_r used to form τ . (Note that by [Ja-Sh3], Theorem 4.4, p.809, this data is recoverable from τ .)

Let ω be a Hecke character. Let τ_1, \dots, τ_r be a irreducible cuspidal automorphic representations of $GL_{2n_1}, \dots, GL_{2n_r}$, respectively.

For each i , let V_{τ_i} denote the space of cuspforms on which τ_i acts. Then pointwise multiplication

$$\varphi_1 \otimes \cdots \otimes \varphi_r \mapsto \prod_{i=1}^r \varphi_i$$

extends to an isomorphism between the abstract tensor product $\bigotimes_{i=1}^r V_{\tau_i}$ and the space of all functions

$$\Phi(g_1, \dots, g_r) = \sum_{i=1}^N c_i \prod_{j=1}^r \varphi_{i,j}(g_j) \quad c_i \in \mathbb{C}, \varphi_{i,j} \in V_{\tau_j} \quad \forall i, j.$$

(This is an elementary exercise.) We consider the representation $\tau_1 \otimes \cdots \otimes \tau_r$ of $GL_{2n_1} \times \cdots \times GL_{2n_r}$, realized on this latter space, which we denote $V_{\otimes \tau_i}$.

Let $n = n_1 + \cdots + n_r$.

We will construct an Eisenstein series on G_{4n} induced from the subgroup $P = MU$ of the Siegel parabolic such that $M \cong GL_{2n_1} \times \cdots \times GL_{2n_r} \times GL_1$. Let s_1, \dots, s_r be a complex variables. Using the identification of M with $GL_{2n_1} \times \cdots \times GL_{2n_r} \times GL_1$ fixed in section 3.3 above, we define an action of $M(\mathbb{A})$ on the space of $\tau_1 \otimes \cdots \otimes \tau_r$ by

$$(5.0.2) \quad (g_1, \dots, g_r, \alpha) \cdot \prod_{j=1}^r \varphi_j(h_j) = \left(\prod_{j=1}^r \varphi(h_j g_j) |\det g_j|^{s_j} \right) \omega(\alpha).$$

We denote this representation of $M(\mathbb{A})$, by $(\bigotimes_{i=1}^r \tau_i \otimes |\det_i|^{s_i}) \boxtimes \omega$.

To shorten the notation, we write $\underline{g} = (g_1, \dots, g_r)$. Then (5.0.2) may be shortened to

$$\underline{g} \cdot \Phi(\underline{h}) = \Phi(\underline{h} \cdot \underline{g}) \left(\prod_{j=1}^r |\det g_j|^{s_j} \right) \omega(\alpha).$$

We shall also employ the shorthand $\underline{s} = (s_1, \dots, s_r)$, and $\underline{\tau} = (\tau_1, \dots, \tau_r)$.

For each \underline{s} we have the induced representation $\text{Ind}_{P(\mathbb{A})}^{G_{4n}(\mathbb{A})} (\bigotimes_{i=1}^r \tau_i \otimes |\det_i|^{s_i}) \boxtimes \omega$, (normalized induction) of $G_{4n}(\mathbb{A})$. The standard realization of this representation is action by right translation on the space $V^{(1)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega)$ given by

$$\left\{ \tilde{F} : G_{4n}(\mathbb{A}) \rightarrow V_{\tau}, \text{ smooth} \left| \tilde{F}((\underline{g}, \alpha)h)(\underline{g}') = \tilde{F}(h)(\underline{g}'\underline{g})\omega(\alpha) \prod_{i=1}^r |\det g_i|^{s_i + n - \frac{1}{2} + \sum_{j=i+1}^r n_j - \sum_{j=1}^{i-1} n_j} \right. \right\}.$$

(The factor

$$\prod_{i=1}^r |\det g_i|^{n - \frac{1}{2} + \sum_{j=i+1}^r n_j - \sum_{j=1}^{i-1} n_j}$$

is equal to $|\delta_P|^{\frac{1}{2}}$, and makes the induction normalized.) A second useful realization is action by right translation on

$$V^{(2)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega) = \left\{ f : G_{4n}(\mathbb{A}) \rightarrow \mathbb{C}, \left| f(h) = \tilde{F}(h)(id), \tilde{F} \in V^{(1)}(\underline{s}, \underline{\tau}, \omega) \right. \right\}.$$

(Here id denotes the identity element of $GL_{2n}(\mathbb{A})$.)

These representations fit together into a fiber bundle over \mathbb{C}^r . So a section of this bundle is a function f defined on \mathbb{C}^r such that $f(\underline{s}) \in V^{(i)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega)$ ($i = 1$ or 2) for each \underline{s} . We shall only require the use of flat, K -finite sections, which are defined as follows. Take $f_0 \in V^{(i)}(\underline{0}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega)$ K -finite, and define $f(\underline{s})(h)$ by

$$f(\underline{s})(u(\underline{g}, \alpha)k) = f_0(u(\underline{g}, \alpha)k) \prod_{i=1}^r |\det g_i|^{s_i}$$

for $u \in U(\mathbb{A})$, $\underline{g} \in GL_{2n_1}(\mathbb{A}) \times \cdots \times GL_{2n_r}(\mathbb{A})$, $\alpha \in \mathbb{A}^\times$, $k \in K$. This is well defined. (I.e., although g_i is not uniquely determined in the decomposition, $|\det g_i|$ is. Cf. the definition of m_P on p.7 of [MW1].)

We begin with a flat K finite section of the bundle of representations realized on the spaces $V^{(2)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega)$.

Remark 5.0.3. *Clearly, the function f is determined by $f(\underline{s}^*)$ for any choice of base point \underline{s}^* . In particular, any function of f may be regarded as a function of $f_{\underline{s}^*} \in V^{(2)}(\underline{s}^*, \bigotimes_{i=1}^r \tau_i \boxtimes \omega)$, for any particular value of \underline{s}^* . We have exploited this fact with $\underline{s}^* = 0$ to streamline the definitions. A posteriori it will become clear that the point $\underline{s}^* = \frac{1}{2} := (\frac{1}{2}, \dots, \frac{1}{2})$ is of particular importance, and we shall then switch to $\underline{s}^* = \frac{1}{2}$.*

For such f the sum

$$E(f)(g)(\underline{s}) := \sum_{\gamma \in P(F) \backslash G(F)} f(\underline{s})(\gamma g)$$

converges for all \underline{s} such that $\text{Re}(s_r), \text{Re}(s_i - s_{i+1}), i = 1$ to $r - 1$ are all sufficiently large. ([MW1], §II.1.5, pp.85-86). It has meromorphic continuation to \mathbb{C}^r ([MW1] §IV.1.8(a), IV.1.9(c), p.140). These are our Eisenstein series. We collect some of their well-known properties in the following theorem.

Theorem 5.0.4. (1) *The function*

$$(5.0.5) \quad \prod_{i \neq j} (s_i + s_j - 1) \prod_{i=1}^r (s_i - \frac{1}{2}) E(f)(g)(\underline{s})$$

is holomorphic at $s = \frac{1}{2}$. (More precisely, while $E(f)(g)$ may have singularities, there is a holomorphic function defined on an open neighborhood of $\underline{s} = \frac{1}{2}$ which agrees with (5.0.7) on the complement of the hyperplanes $s_i = \frac{1}{2}$, and $s_i + s_j = 1$.)

(2) *The function (5.0.5) remains holomorphic (in the same sense) when $s_i + s_j - 1$ is omitted, provided $\tau_i \not\cong \omega \otimes \bar{\tau}_j$. It remains holomorphic when $s_i - \frac{1}{2}$ is omitted, provided τ_i is not $\bar{\omega}$ -symplectic. Furthermore, each of these sufficient conditions is also necessary, in that the holomorphicity conclusion will fail, for some f and g , if any of the factors is omitted without the corresponding condition on $\underline{\tau}$ being satisfied. From this we deduce that if*

$$(5.0.6) \quad \text{the representations } \tau_1, \dots, \tau_r \text{ are all distinct and } \bar{\omega}\text{-symplectic,}$$

then the function

$$(5.0.7) \quad \prod_{i=1}^r (s_i - \frac{1}{2}) E(f)(g)(\underline{s})$$

is holomorphic at $s = \frac{1}{2}$ for all f, g and nonvanishing at $s = \frac{1}{2}$ for some f, g .

- (3) Let us now assume condition (5.0.6) holds, and regard f as a function of $f_{\frac{1}{2}} \in V^{(2)}(\frac{1}{2}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega)$. Let $E_{-1}(f_{\frac{1}{2}})(g)$ denote the value of the function (5.0.7) at $\underline{s} = \frac{1}{2}$ (defined by analytic continuation). Then $E_{-1}(f)$ is an L^2 function for all $f_{\frac{1}{2}} \in V^{(2)}(\frac{1}{2}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega)$.
- (4) The function E_{-1} is an intertwining operator from $\text{Ind}_{P(\mathbb{A})}^{G_{4n}(\mathbb{A})}(\bigotimes_{i=1}^r \tau_i \otimes |\det_i|^{\frac{1}{2}}) \boxtimes \omega$ into the space of L^2 automorphic forms.
- (5) If $\mathcal{E}_{-1}(\tau, \omega)$ is the image of E_{-1} , and ψ_{LW} is the character of U_{\max} given by $\psi_{LW}(u) = \psi_0(\sum_{i=1}^{2n-1} u_{i,i+1})$, then $(U_{\max}, \psi_{LW}) \notin \mathcal{U}^\perp(\mathcal{E}_{-1}(\tau, \omega))$.
- (6) The space of functions $\mathcal{E}_{-1}(\tau, \omega)$ does not depend on the order chosen on the cuspidal representations τ_1, \dots, τ_r . Thus it is well-defined as a function of the isobaric representation τ .

Remark 5.0.8. By induction in stages, the induced representation $\text{Ind}_{P(\mathbb{A})}^{G_{4n}(\mathbb{A})}(\bigotimes_{i=1}^r \tau_i \otimes |\det_i|^{\frac{1}{2}}) \boxtimes \omega$, which comes up in part (4) of the theorem can also be written as $\text{Ind}_{P_{\text{Siegel}}(\mathbb{A})}^{G_{4n}(\mathbb{A})} \tau \otimes |\det|^{\frac{1}{2}} \boxtimes \omega$, where $\tau = \tau_1 \boxplus \dots \boxplus \tau_r$ as before, and P_{Siegel} is the Siegel parabolic. (Cf. section 2.) Here, we also exploit the identification of the Levi M_{Siegel} of P_{Siegel} with $GL_{2n} \times GL_1$ fixed in 3.3.1.

Proof. We first review the standard arguments by which the presence or absence of a singularity of an Eisenstein series reduces to the presence or absence of a singularity of a relative rank one intertwining operator. To do so, we recall the set

$$W(M) := \left\{ w \in W_{G_{4n}} \mid \begin{array}{l} w \text{ is of minimal length in } w \cdot W_M \\ wMw^{-1} \text{ is a standard Levi of } G_{4n} \end{array} \right\}.$$

It will be convenient and harmless to treat the elements of $W(M)$ as though they were elements of $G_{4n}(F)$, rather than repeatedly choose representatives and remark the independence of the choice. For each $w \in W(M)$, $\underline{s} \in \mathbb{C}^r$, we define P^w to be the standard parabolic with Levi wMw^{-1} . For \underline{s} such that s_r and $s_i - s_{i+1}$, $i = 1$ to $r - 1$ are all sufficiently large, the integral

$$M(w, \underline{s})f(g) := \int_{U_{\max} \cap w \overline{U_{\max}} w^{-1} (F \backslash \mathbb{A})} f(\underline{s})(w^{-1}ug) du$$

converges ([MW1], II.1.6), defining an operator $M(w, \underline{s})$ from $V^{(2)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega)$ to a space of functions which is easily verified to afford a realization of

$$\text{Ind}_{P^w(\mathbb{A})}^{G_{4n}(\mathbb{A})} \left(\left(\bigotimes_{i=1}^r \tau_i \otimes |\det_i|^{s_i} \right) \boxtimes \omega \right) \circ Ad(w^{-1}).$$

Here, $((\bigotimes_{i=1}^r \tau_i \otimes |\det_i|^{s_i}) \boxtimes \omega) \circ Ad(w^{-1})$, denotes the representation of wMw^{-1} obtained by composing the representation $(\bigotimes_{i=1}^r \tau_i \otimes |\det_i|^{s_i}) \boxtimes \omega$ of M with conjugation by w^{-1} . We denote this latter space of functions by $V_w^{(2)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega)$. Then $M(w, \underline{s})f(g)$ has meromorphic continuation to \mathbb{C}^r . (IV.1.8(b).)

It may be helpful also to review the sorts of singularities which Eisenstein series and intertwining operators have—lying along so-called “root hyperplanes.” (cf. IV.1.6) We defer the notion of “root hyperplane” until later. For now, we allow arbitrary hyperplanes in \mathbb{C}^r , defined by equations of the form $l(\underline{s}) = c$, with l a linear functional $\mathbb{C}^r \rightarrow \mathbb{C}$ and c a constant. Then for any bounded open set $U \subset \mathbb{C}^r$, there exist a finite number of distinct hyperplanes H_1, \dots, H_N , which “carry” the singularities of the Eisenstein series and intertwining operators in U , in the following sense. For each i fix l_i, c_i such that $H_i = \{\underline{s} \in \mathbb{C}^r \mid l_i(\underline{s}) = c_i\}$. Then for each i there is a non-negative integer

$\nu(H_i)$ such that

$$(5.0.9) \quad \prod_{i=1}^N (l_i(\underline{s}) - c_i)^{\nu(H_i)} E(f)(g)(\underline{s})$$

continues to a function holomorphic on all of U . Covering \mathbb{C}^r with bounded open sets and taking a union, we obtain an infinite, but *locally* finite, set of hyperplanes which carry all the singularities of the Eisenstein series and intertwining operators. The same hyperplane H will of course occur more than once. It is easily verified that the minimal exponent $\nu(H)$ appearing in (5.0.9) is the same each time. Thus we may speak of whether an Eisenstein series or intertwining operator does or does not have a pole along H , and of the order of the pole.

One may define “analytic/meromorphic continuation” for functions taking values in Fréchet spaces of locally L^2 functions and the like ([MW1] I.4.9, IV.1.3) of functions and operators. In this case, outside of the domain of convergence, one’s functions are defined only up to L^2 equivalence. However, in view of I.4.10, one has a unique smooth representative for the class. For us it will be more convenient simply to adopt the convention that when we say the Eisenstein series has a pole along H , we mean for some f, g .

Now let us state the relationship between poles of Eisenstein series and intertwining operators, which we prove in an appendix.

Proposition 5.0.10. *For $f \in V^{(2)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega)$, there exists $g \in G_{4n}(\mathbb{A})$ such that $E(f)(g)$ has a pole along H if and only if there exist $w \in W(M), g' \in G_{4n}(\mathbb{A})$ such that $M(w, \underline{s})f(g')$ has a pole along H .*

The same construction can be performed with the Levi M replaced by wMw^{-1} , yielding an operator

$$M_w(w', w \cdot \underline{s}) : V_w^{(2)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega) \rightarrow V_{w'w}^{(2)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega),$$

for each $w' \in W(wMw^{-1})$. Furthermore, one has for all f, g , the equality of meromorphic functions

$$M_w(w', w \cdot \underline{s}) \circ M(w, \underline{s})f(g) = M(w'w, \underline{s})f(g)$$

([MW1], II.1.6, IV.4.1). (For now, the reader may think of “ $w \cdot \underline{s}$ ” simply as a notational contrivance. We shall give it a precise meaning below.)

Next we wish to describe the decomposition of $w \in W(M)$ as a product of elementary symmetries, as in [MW1] I.1.8. The lattice $X(Z_M)$ of rational characters of the center of M has a unique basis $\{e_0, \varepsilon_1, \dots, \varepsilon_r\}$, with the property that for each $i = 1, \dots, m$, there exists $j \in \{1, \dots, r\}$ such that the restriction of e_i as in 3.3 to Z_M is ε_j . The set of restrictions of positive roots of G_{4n} to Z_M is

$$\{0\} \cup \{\varepsilon_i - \varepsilon_j : 1 \leq i < j \leq r\} \cup \{\varepsilon_i + \varepsilon_j : 1 \leq i < j \leq r\} \cup \{2\varepsilon_i : 1 \leq i \leq r\}.$$

We denote the set obtained by excluding zero by $\Phi^+(Z_M)$. For $\alpha \in \Phi^+(Z_M)$, and $w \in W(M)$, one may say “ $w\alpha > 0$ ” or “ $w\alpha < 0$ ” without ambiguity.

Each element $w \in W(M)$ can be decomposed as a product $s_{\alpha_1} \dots s_{\alpha_\ell}$ of elementary symmetries as in [MW1] I.1.8. The element s_{α_ℓ} will be in $W(M)$, while $s_{\alpha_{\ell-1}}$ will be in $W(s_{\alpha_\ell} M s_{\alpha_\ell}^{-1})$ and so on. Each is labeled with the unique restricted root (for the operative Levi) which it reverses. That is $\{\alpha \in \Phi^+(Z_M) : s_{\alpha_\ell} \alpha < 0\}$ is singleton, and α_ℓ is the unique element. (Cf. [MW1] I.1.8, observe that all elements of $\Phi^+(Z_M)$ are indivisible.)

Let $w = s_{\alpha_1} \dots s_{\alpha_\ell}$ be a minimal-length decomposition into elementary symmetries, and put $w_i = s_{\alpha_{i+1}} \dots s_{\alpha_\ell}$. Then

$$\{\alpha \in \Phi^+(Z_M) | w\alpha < 0\} = \{w_i^{-1} \alpha_i | 1 \leq i \leq \ell\}$$

and ℓ is the cardinality of this set (i.e., there is no repetition). Combining this discussion with that of the previous paragraphs, we obtain a decomposition of $M(w, \underline{s})$ as a composite of intertwining operators $M_{w_i}(s_{\alpha_i}, w_i \cdot \underline{s})$, each corresponding naturally to one of the elements of $\{\alpha \in \Phi^+(Z_M) \mid w\alpha < 0\}$.

Let \det_i denote the rational character $(g, \alpha) \mapsto \det g_i$ of M . Then $\{e_0, \det_1, \dots, \det_r\}$ is a basis for the lattice $X(M)$ of rational characters of M . Here, the character e_0 of T introduced in 3.3 has been identified with a character of M as in 3.3.1. Let $\{e_0^*, \det_1^*, \dots, \det_r^*\}$ be the dual basis of the dual lattice. Again, e_0^* is the same as in 3.3. Elements of $X(M)$ may be paired with elements of $X^\vee(T)$ defining a projection from $X^\vee(T)$ onto the dual lattice. For each $i = 1, \dots, m$, there exists unique $j \in \{1, \dots, r\}$ such that e_i^* maps to \det_j^* . If α is a root, then the projection of the coroot α^\vee to the dual lattice of $X(M)$ depends only on the restriction of α to Z_M , and the correspondence is as follows:

$$\begin{aligned} 0 &\leftrightarrow 0, \\ \varepsilon_i - \varepsilon_j &\leftrightarrow \det_i^* - \det_j^*, \\ \varepsilon_i + \varepsilon_j &\leftrightarrow \det_i^* + \det_j^* - e_0^*, \\ 2\varepsilon_i &\leftrightarrow 2\det_i^* - e_0^*. \end{aligned}$$

We denote the element corresponding to $\alpha \in \Phi^+(Z_M)$ by α^\vee (in agreement with [MW1], I.1.11).

We may identify $\underline{s} \in \mathbb{C}^r$ with

$$\sum_{i=1}^r \det_i \otimes s_i \in X(M) \otimes_{\mathbb{Z}} \mathbb{C}.$$

This is compatible with [MW1], I.1.4. Restriction of functions gives a natural injective map $X(M) \rightarrow X(T)$, and hence $X(M) \otimes_{\mathbb{Z}} \mathbb{C} \rightarrow X(T) \otimes_{\mathbb{Z}} \mathbb{C}$, which we use to identify the first space with a subspace of the second. This gives the notation $w \cdot \underline{s}$ a precise meaning, as an element of $X(wMw^{-1}) \otimes_{\mathbb{Z}} \mathbb{C}$, which is compatible with the usage above. In addition, it gives a “meaning” to the set

$$\{s_i - s_j\} \cup \{s_i + s_j\} \cup \{2s_i\},$$

of linear functionals on \mathbb{C}^r , identifying each with an element of $\Phi^+(Z_M)$. Formally,

Definition 5.0.11. *A root hyperplane (relative to the Levi M) is a hyperplane of the form*

$$H = \{s \in \mathbb{C}^r \mid \langle \alpha^\vee, \underline{s} \rangle = c\}$$

for some $\alpha \in \Phi^+(Z_M)$ and $c \in \mathbb{C}$. We say that the hyperplane H is associated to the root α , which is uniquely determined.

The next main statement is

Lemma 5.0.12. *Let $w = s_{\alpha_1} \dots s_{\alpha_\ell}$ be any decomposition of minimal length, and for each i let $w_i = s_{\alpha_{i+1}} \dots s_{\alpha_\ell}$. Then the set of poles of $M(w, \underline{s})$ is the disjoint union of the sets of poles of the operators $M_{w_i}(s_{\alpha_i}, w_i \cdot \underline{s})$. A pole of $M(w, \underline{s})$ comes from $M_{w_i}(s_{\alpha_i}, w_i \cdot \underline{s})$ if and only if it is associated to $w_i^{-1}\alpha_i$. Furthermore, if $\{s \in \mathbb{C}^r \mid \langle \alpha^\vee, \underline{s} \rangle = c\}$ is a pole of $M(w, \underline{s})$, then $c \neq 0$.*

We now prove (1). A root hyperplane passing through $\frac{1}{2}$ is defined by an equation of one of three forms: $s_i = \frac{1}{2}$, $s_i + s_j = 1$, or $s_i - s_j = 0$. The third kind can not support singularities of the Eisenstein series. The first two can, but by [MW1]IV.1.11 (c), they will be without multiplicity, and so the factor of

$$\prod_{i \neq j} (s_i + s_j - 1) \prod_{i=1}^r (s_i - \frac{1}{2})$$

will take care of them.

The operators corresponding to elementary symmetries are called relative rank one because they could be defined without reference G_{4n} , considering M instead as a maximal Levi of another Levi subgroup M_α of G_{4n} , having semisimple rank one greater than that of M . Furthermore, in a suitable sense, the relative rank one operator only “lives on one component of M_α ,” which will allow us to deduce the general case of (2) from the case $r = 1$ and a similar fact about intertwining operators on GL_n . Let us make this more precise.

Fix $\alpha \in \Phi^+(Z_M)$. There is a minimal Levi subgroup M_α of G_{4n} containing M such that α is the restriction of a root of M_α . (It is standard iff α is the restriction of a simple root.) Fix $w \in W(M)$ such that $w\alpha < 0$, and a decomposition $w = s_{\alpha_1} \dots s_{\alpha_\ell}$ of w as into elementary symmetries, which is of minimal length. For some unique i , we have $\alpha = w_i^{-1}\alpha_i$, where w_i is as above. Then $w_i M_\alpha w_i^{-1}$ is a standard Levi of G_{4n} . Different choices of decomposition give different (even conjugate) embeddings of the same reductive group into G_{4n} as a standard Levi.

If $\alpha = \varepsilon_j - \varepsilon_k$, or $\varepsilon_j + \varepsilon_k$, then M_{α_i} is isomorphic to $GL_{2(n_j+n_k)} \times \prod_{l \neq j,k} GL_{2n_l} \times GL_1$. while if $\alpha = 2\varepsilon_j$, it is isomorphic to $G_{4n_j} \times \prod_{k \neq j} GL_{2n_k}$. Let G' denote $GL_{2(n_j+n_k)}$ or G_{4n_j} as appropriate and let ι be a choice of isomorphism with the “new” factor. Then $\iota^{-1}(\iota(G') \cap P^{w_i})$ is a maximal parabolic subgroup $P' = M'U'$ of G' , and $\sigma := (\bigotimes_{i=1}^r \tau \otimes \omega) \circ Ad(w_i) \circ \iota$, is an irreducible unitary cuspidal automorphic representation of $M'(\mathbb{A})$. The map ι also induces a linear projection

$$\iota_* : X(w_i M w_i^{-1}) \otimes_{\mathbb{Z}} \mathbb{C} \rightarrow X(M') \otimes_{\mathbb{Z}} \mathbb{C}.$$

(Recall that we have agreed to think of $w_i \cdot \underline{s}$ as an element of the former space.)

Following, [MW1] I.1.4, define m^μ for $m \in M'(\mathbb{A})$ and μ in $X(M') \otimes_{\mathbb{Z}} \mathbb{C}$, by stipulating that $m^\mu = |\chi(m)|^s$ if $\mu = \chi \otimes s$ and $m^{\mu_1 + \mu_2} = m^{\mu_1} m^{\mu_2}$.

The set $W'_G(M')$, defined analogously to $W(M)$ above, contains a unique nontrivial element. It is the elementary symmetry s_β associated to the restriction to Z'_M of any of the positive roots of G' which are not roots of M' . The map ι identifies s_β with s_{α_i} .

For $\mu \in X(M') \otimes_{\mathbb{Z}} \mathbb{C}$, let $V^{(1)}(\mu, \sigma)$ denote

$$\{h : G'(\mathbb{A}) \rightarrow V_\sigma, \text{ smooth} | h(mg')(m') = h'(g')(m'm)m^{\mu + \rho_{P'}} \quad m, m' \in M'(\mathbb{A}), g' \in G'(\mathbb{A})\},$$

$$V^{(2)}(\mu, \sigma) = \{h : G'(\mathbb{A}) \rightarrow \mathbb{C}, \text{ smooth} | h(g')(e) \in V^{(1)}(\mu, \sigma)\}.$$

There is a standard intertwining operator $M(s_\beta, \mu) : V^{(2)}(\mu, \sigma) \rightarrow V_{s_\beta}^{(2)}(\mu, \sigma)$. One has the identity

$$M_{w_{i-1}}(s_{\alpha_i}, w_i \cdot \underline{s}) f(\iota(h)g) = M(s_\beta, \mu) f(\iota(h)g).$$

That is, if p_g denotes the map

$$V_{w_i}^{(2)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega) \rightarrow V^{(2)}(\mu, \sigma)$$

corresponding to evaluation at $\iota(h)g$ for a fixed g , then, for all g , the following diagram commutes:

$$\begin{array}{ccc} V_{w_i}^{(2)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega) & \xrightarrow{M_{w_i}(s_{\alpha_i}, w_i \cdot \underline{s})} & V_{w_{i-1}}^{(2)}(\underline{s}, \bigotimes_{i=1}^r \tau_i \boxtimes \omega) \\ p_g \downarrow & & p_g \downarrow \\ V^{(2)}(\iota_*(w_i \cdot \underline{s} + \rho_{P_{\alpha_i}}), \sigma) & \xrightarrow{M(s_\beta, \iota_*(w_i \cdot \underline{s} + \rho_{P_{\alpha_i}}))} & V_{s_\beta}^{(2)}(\iota_*(w_i \cdot \underline{s} + \rho_{P_{\alpha_i}}), \sigma). \end{array}$$

Hence $M_{w_i}(s_{\alpha_i}, w_i \cdot \underline{s})$ has a pole along a root hyperplane associated to α iff $M(\iota_*(w_i \cdot \underline{s} + \rho_{P_{\alpha_i}}), \sigma)$ does.

Since the set of poles of $M_{w_i}(s_{\alpha_i}, w_i \cdot \underline{s})$ is equal to the set of poles of $M(w, \underline{s})$ along hyperplanes associated to α , it is independent of the choice of decomposition $w = s_{\alpha_1} \dots s_{\alpha_\ell}$. Hence, for each $\alpha \in \Phi^+(Z_M)$, we may use a decomposition tailored to that α .

First suppose $\alpha = \varepsilon_j - \varepsilon_k$. One may choose a decomposition so that w_i corresponds to the permutation matrix in GL_{2n} (identified with a subgroup of the Siegel Levi) which moves the j th block of M up so that it is immediately after the i th, and otherwise preserves order. It is then easily verified that $\sigma = \tau_i \otimes \tau_j$ and

$$\begin{pmatrix} h_1 & \\ & h_2 \end{pmatrix}^{\iota_*(w_i \cdot s) + \rho_{P\alpha_i}} = |\det h_1|^{s_i + \kappa} |\det h_2|^{s_j + \kappa},$$

where $\kappa = \sum_{k>i, k \neq j} n_k - \sum_{k<i} n_k + n - \frac{1}{2}$.

Next suppose $\alpha = 2\varepsilon_j$. Then we choose a decomposition so that w_i is in the Weyl group of GL_{2n} , and moves the j th block to be last, otherwise preserving order. Then one easily verifies that σ is the representation $\tau_j \boxtimes \omega$ of the Siegel Levi of G_{4n_j} , and that, for (g', α) in the Siegel Levi of G_{4n_j} ,

$$(g', \alpha)^{\iota_*(w_i \cdot s) + \rho_{P\alpha_i}} = |\det g|^{s_j}.$$

Finally, suppose $\alpha = \varepsilon_j + \varepsilon_k$. Then we choose a decomposition so that w_i that projects to a permutation matrix in SO_{4n} of the form

$$\begin{pmatrix} I & & & \\ & I & & \\ & & I & \\ I & & & 1 \end{pmatrix},$$

with the off-diagonal blocks being $2n_j \times 2n_j$, and the first block being $\sum_{k=1}^i 2n_k$. We deduce from Corollary 3.4.6 that $\sigma = \tau_i \otimes (\tilde{\tau}_j \otimes \omega)$, and from Lemma 3.4.5 that

$$\begin{pmatrix} h_1 & \\ & h_2 \end{pmatrix}^{\iota_*(w_i \cdot s) + \rho_{P\alpha_i}} = |\det h_1|^{s_i + \kappa} |\det h_2|^{-s_j + \kappa},$$

where κ is as before.

Thus (2) follows from

Proposition 5.0.13. *Let w denote the unique nontrivial element of $W(M)$, in the case when M is the Levi of the Siegel parabolic. Let τ be a cuspidal representation of GL_m . Then $M(w, s)f(g)$ has a pole at $s = \frac{1}{2}$ for some $f \in \text{Ind}_{P(\mathbb{A})}^{G_{2m}(\mathbb{A})}(\tau \otimes |\det|^s) \boxtimes \omega$, and $g \in G_{2m}(\mathbb{A})$ if and only if τ is $\bar{\omega}$ -symplectic.*

Remark 5.0.14. *Of course, we only need the case $m = 2n$. Furthermore, if m is odd, then τ is never $\bar{\omega}$ -symplectic. We hope to show more clearly how the ideas fit together by stating the result for general m . It is proved also for general m , because the proof is “blind to m .”*

Proposition 5.0.15. *Let $P = MU$ be a maximal standard parabolic of GL_n such that $M \cong GL_k \times GL_{n-k}$. Let f be an element of $\text{Ind}_{P(\mathbb{A})}^{GL_n(\mathbb{A})}(\tau_1 \otimes |\det|^{s_1}) \otimes (\tau_2 \otimes |\det|^{s_2})$. Let w be the unique nontrivial element of $W(M)$. Then $M(w, \underline{s})f(g)$ is singular along the hyperplane $s_1 - s_2 = 1$ for some f, g iff $\tau_2 \cong \tau_1$.*

We defer the proofs to the appendix.

Now, we assume (5.0.6) holds and prove the remaining part of the theorem. Let $N(\underline{s}) = \prod_{i=1}^r (s_i - \frac{1}{2})$.

Item (3) follows from [MW1] I.4.11. The constant term of $E(f)$ along a parabolic $P' = M'U'$ has nontrivial cuspidal component iff M' is conjugate to M . (IV.1.9 (b)(ii)). For such P' it is equal to

$$\sum_{w \in W(M), wMw^{-1} = M'} M(w, \underline{s})f(g).$$

Take $w \in W(M)$, such that $wMw^{-1} = M'$. If $w \cdot (2\varepsilon_i) > 0$ for some i , then $M(w, \underline{s})f(g)$ does not have a pole at $s_i - \frac{1}{2}$, and hence $N(\underline{s})M(w, \underline{s})f(g)$ vanishes at $\frac{1}{2}$. On the other hand, if $w \cdot (2\varepsilon_i) < 0$ for all i , then $M(w, \underline{s})f(g)$ satisfies the criterion of I.4.11.

It follows from [MW1] IV.1.9 (b)(i) applied to $N(\underline{s})E(f)$ (which is valid by IV.1.9 (d)) that the residue is an automorphic form. To complete the proof of (4), let $\rho(g)$ denote right translation. It is clear that for values of s in the domain of convergence, $N(\underline{s})E(\rho(g)f)(\underline{s}) = N(\underline{s})\rho(g)(E(f)(\underline{s}))$. By uniqueness of analytic continuation, the equality also holds at values of s where both sides are defined by analytic continuation, including $\frac{1}{2}$. The action of the Lie algebra at the infinite places is handled similarly.

Next we consider the constant term of $E(f)$ along the Siegel parabolic. By [MW1] II.1.7(ii) it may be expressed in terms of GL_{2n} Eisenstein series, formed using the functions $M(w, \underline{s})f$, corresponding to those $w \in W(M)$ such that $w^{-1}(e_i - e_{i+1}) > 0$ for all i . (Note: we proved in Lemma 3.4.3 that wMw^{-1} is contained in the Siegel Levi for every $w \in W(M)$.) When we pass to $E_{-1}(f)$, the term corresponding to w only survives if $w \cdot (2\varepsilon_i) < 0$ for all i . This condition picks out a unique element, w_0 . It is the shortest element of $W_{GL_{2n}} \cdot w_\ell \cdot W_{GL_{2n}}$, where w_ℓ is the longest element of $W_{G_{4n}}$, and we have identified GL_{2n} with a subgroup of the Siegel Levi as usual. Via corollary 3.4.6 one finds that

$$\left(\bigotimes_{i=1}^r \tau_i \boxtimes \omega\right) \circ Ad(w_0) = \left(\bigotimes_{i=1}^r (\tilde{\tau}_{r+1-i} \otimes \omega) \boxtimes \omega\right) = \left(\bigotimes_{i=1}^r \tau_{r+1-i} \boxtimes \omega\right).$$

For $f \in V^{(2)}(\bigotimes_{i=1}^r \tau_i \boxtimes \omega, \frac{1}{2})$, $M(w_0, \frac{1}{2})f|_{GL_{2n}(\mathbb{A})}$ is an element of the analogue of $V^{(2)}(\bigotimes_{i=1}^r \tau_i \boxtimes \omega, \underline{s})$, for the induced representation

$$\text{Ind}_{\bar{P}^0(\mathbb{A})}^{GL_{2n}(\mathbb{A})} \left(\bigotimes_{i=1}^r \tau_{r+1-i} \otimes |\det_i|^{\frac{n-2}{2}}\right) = |\det|^{\frac{n-2}{2}} \otimes \tau$$

of GL_{2n} . Here $\bar{P}^0 = GL_{2n} \cap P^{w_0}$, and $\tau = \tau_1 \boxplus \cdots \boxplus \tau_r$. Furthermore, since this representation is irreducible, it may be regarded as an arbitrary element. Also, we may regard this representation as induced from τ_1, \dots, τ_r in the usual order. Let \bar{P} denote the relevant parabolic of GL_{2n} .

The representation τ sits inside a fiber bundle of induced representations $\text{Ind}_{\bar{P}(\mathbb{A})}^{GL_{2n}(\mathbb{A})} (\bigotimes_{i=1}^r \tau_i \otimes |\det_i|^{s_i})$. For a flat, K -finite section f let $E^{GL_{2n}}(f)(g)(\underline{s})$ be the GL_{2n} Eisenstein series defined by

$$\sum_{\bar{P}(F) \backslash GL_{2n}(F)} f(\underline{s})(\gamma g)$$

when $s_i - s_{i+1}$ is sufficiently large for each i , and by meromorphic continuation elsewhere.

Let $U_{\max}^{GL_{2n}}$ denote the usual maximal unipotent subgroup of GL_{2n} , consisting of all upper triangular unipotent matrices. Let $\psi_W(u) = \psi_0(u_{1,2} + \cdots + u_{m-1,m})$ be the usual generic character.

To complete the proof of (5), we must prove that

$$(5.0.16) \quad \int_{U_{\max}^{GL_{2n}}(F \backslash \mathbb{A})} E^{GL_{2n}}(f)(ug)(\underline{0})\psi_W(u) du \neq 0$$

for some $f \in \text{Ind}_{\bar{P}(\mathbb{A})}^{GL_{2n}(\mathbb{A})} \bigotimes_{i=1}^r \tau_{r+1-i}$, $g \in GL_{2n}(\mathbb{A})$, i.e., that the space of GL_{2n} Eisenstein series $E^{GL_{2n}}(f)$ is globally ψ_W -generic. Granted this, (5) follows from [MW1]II.1.7(ii) and the discussion just above.

The following proposition follows from work of Shahidi.

Proposition 5.0.17.

$$\int_{U_{\max}^{GL_{2n}}(F \backslash \mathbb{A})} E^{GL_{2n}}(f)(ug)(\underline{s})\psi_W(u) du = \prod_{v \in S} W_v(g_v) \cdot \prod_{v \notin S} W_v^\circ(g_v) \cdot \prod_{i < j} L^S(s_i - s_j + 1, \tau_i \times \tilde{\tau}_j)^{-1},$$

where, for each v , W_v is a Whittaker function in the $\psi_{W,v}$ -Whittaker model of $\text{Ind}_{\bar{P}(F_v)}^{GL_{2n}(F_v)}(\otimes_{i=1}^r \tau_{i,v} \otimes |\det_i|_v^{s_i})$, S is a finite set of places, depending on f , outside of which τ_v is unramified and W_v° is the normalized spherical vector in the the $\psi_{W,v}$ -Whittaker model of $\text{Ind}_{\bar{P}(F_v)}^{GL_{2n}(F_v)}(\otimes_{i=1}^r \tau_{i,v} \otimes |\det_i|_v^{s_i})$. A flat, K -finite section f may be chosen so that, for all $v \in S$, the function W_v is not identically zero at $\underline{s} = \underline{0}$.

We briefly review the steps of the proof in the appendix.

It follows from [Ja-Sh3] Propositions 3.3 and 3.6 that the product of partial L functions appearing in Proposition 5.0.17 does not have a pole at $\underline{s} = \underline{0}$ provided the representations τ_1, \dots, τ_r are distinct. This completes the proof of (5).

Finally, (6) follows from the functional equation of the Eisenstein series ([MW1]IV.1.10(a)), and the fact that τ is equal to an irreducible full induced representation (as opposed to a constituent of a reducible one). \square

6. MAIN RESULTS

6.1. Descent Construction. In this section, we shall make use of remark 5.0.8, and regard $\mathcal{E}_{-1}(\tau, \omega)$ as affording an automorphic realization of the representation induced from the representation $\tau \otimes |\det|^{\frac{1}{2}} \boxtimes \omega$ of the Siegel Levi. Thus we may dispense with the smaller Levi denoted by P in the previous section, and in this section we denote the Siegel parabolic more briefly by $P = MU$.

Next we describe certain unipotent periods of G_{2m} which play a key role in the argument. For $1 \leq \ell < m$, let N_ℓ be the subgroup of U_{\max} defined by $u_{ij} = 0$ for $i > \ell$. (Recall that according to the convention above, this refers only to those i, j with $i < j \leq m - i$.) This is the unipotent radical of a standard parabolic Q_ℓ having Levi L_ℓ isomorphic to $GL_1^\ell \times G_{2m-2\ell}$.

Let ϑ be a character of N_ℓ then we may define

$$DC^\ell(\tau, \omega, \vartheta) = FC^\vartheta \mathcal{E}_{-1}(\tau, \omega).$$

Theorem 6.1.1. *Let ω be a Hecke character. Let $\tau = \tau_1 \boxplus \dots \boxplus \tau_r$ be an isobaric sum of $\bar{\omega}$ -symplectic irreducible cuspidal automorphic representations τ_1, \dots, τ_r , of $GL_{2n_1}(\mathbb{A}), \dots, GL_{2n_r}(\mathbb{A})$, respectively. If $\ell \geq n$, and ϑ is in general position, then*

$$DC^\ell(\tau, \omega, \vartheta) = \{0\}.$$

Proof. By Theorem 5.0.4, (3) the representation $\mathcal{E}_{-1}(\tau, \omega)$ decomposes discretely. Let $\pi \cong \otimes'_v \pi_v$ be one of the irreducible components, and $p_\pi : \mathcal{E}_{-1}(\tau, \omega) \rightarrow \pi$ the natural projection.

Fix a place v_0 such which τ_{v_0} and π_{v_0} are unramified. For any $\xi^{v_0} \in \otimes'_{v \neq v_0} \text{Ind}_{P(F_v)}^{G_{4n}(F_v)} \tau_v \otimes |\det|_v^{\frac{1}{2}} \boxtimes \omega_v$ we define a map

$$i_{\xi^{v_0}} : \text{Ind}_{P(F_{v_0})}^{G_{4n}(F_{v_0})} \tau_{v_0} \otimes |\det|_{v_0}^{\frac{1}{2}} \boxtimes \omega_{v_0} \rightarrow \text{Ind}_{P(\mathbb{A})}^{G_{4n}(\mathbb{A})} \tau \otimes |\det|^{\frac{1}{2}} \boxtimes \omega$$

by $i_{\xi^{v_0}}(\xi_v) = \iota(\xi_{v_0} \otimes \xi^{v_0})$, where ι is an isomorphism of the restricted product $\otimes'_v \text{Ind}_{P(F_v)}^{G_{4n}(F_v)} \tau_v \otimes |\det|_v^{\frac{1}{2}} \boxtimes \omega_v$ with the global induced representation $\text{Ind}_{P(\mathbb{A})}^{G_{4n}(\mathbb{A})} \tau \otimes |\det|^{\frac{1}{2}} \boxtimes \omega$. Clearly

$$\mathcal{E}_{-1}(\tau, \omega) = E_{-1} \circ \iota(\otimes'_v \text{Ind}_{P(F_v)}^{G_{4n}(F_v)} \tau_v \otimes |\det|_v^{\frac{1}{2}} \boxtimes \omega_v).$$

For any decomposable vector $\xi = \xi_{v_0} \otimes \xi^{v_0}$,

$$p_\pi \circ E_{-1} \circ \iota(\xi) = p_\pi \circ E_{-1} \circ i_{\xi^{v_0}}(\xi_{v_0}).$$

Thus, π_{v_0} is a quotient of $\text{Ind}_{P(F_{v_0})}^{G_{4n}(F_{v_0})} \tau_{v_0} \otimes |\det|_{v_0}^{\frac{1}{2}} \boxtimes \omega_{v_0}$, and hence (since we took v_0 such that π_{v_0} is unramified) it is isomorphic to the unramified constituent ${}^{un}\text{Ind}_{P(F_{v_0})}^{G_{4n}(F_{v_0})} \tau_{v_0} \otimes |\det|_{v_0}^{\frac{1}{2}} \boxtimes \omega_{v_0}$.

Denote the isomorphism of π with $\otimes'_{v \neq v_0} \pi_v$ by the same symbol ι . This time, fix $\zeta^{v_0} \in \otimes'_{v \neq v_0} \pi_v$, and define $i_{\zeta^{v_0}} : {}^{un}\text{Ind}_{P(F_{v_0})}^{G_{4n}(F_{v_0})} \tau_{v_0} \otimes |\det|_{v_0}^{\frac{1}{2}} \boxtimes \omega_{v_0} \rightarrow \pi$. It follows easily from the definitions that

$$FC^{\vartheta} \circ i_{\zeta^{v_0}}$$

factors through the Jacquet module $\mathcal{J}_{N_{\ell, \vartheta}}({}^{un}\text{Ind}_{P(F_{v_0})}^{G_{4n}(F_{v_0})} \tau_{v_0} \otimes |\det|_{v_0}^{\frac{1}{2}} \boxtimes \omega_{v_0})$. In appendix 8 we show that this Jacquet module is zero. The result follows. \square

Remark 6.1.2. *A general character of N_{ℓ} is of the form*

$$\psi(c_1 u_{1,2} + \cdots + c_{\ell-1} u_{\ell-1, \ell} + d_1 u_{\ell, \ell+1} + \cdots + d_{4n-2\ell} u_{\ell, 4n-\ell}).$$

The Levi L_{ℓ} acts on the space of characters (cf. section 4.3). Over an algebraically closed field there is an open orbit, which consists of all those elements such that $c_i \neq 0$ for all i and ${}^t \underline{d} J \underline{d} \neq 0$. Here, \underline{d} is the column vector ${}^t(d_1, \dots, d_{4n-2\ell})$, and J is defined as in 3.1. Over a general field two such elements are in the same F -orbit iff the two values of ${}^t \underline{d} J \underline{d}$ are in the same square class.

Let ψ_{ℓ} be the character of N_{ℓ} defined by

$$\psi_{\ell}(u) = \psi_0(u_{12} + \cdots + u_{\ell-1, \ell} + u_{\ell, 2n} - u_{\ell, 2n+1}).$$

It is not hard to see that

- the stabilizer $L_{\ell}^{\psi_{\ell}}$ (cf. M^{ϑ} in definition 4.3.1) has two connected components,
- the one containing the identity is isomorphic to $G_{4n-2\ell-1}$,
- there is an “obvious” choice of isomorphism $\text{inc} : G_{4n-2\ell-1} \rightarrow (L_{\ell}^{\psi_{\ell}})^0$ having the following property: if $\{e_i^* : i = 0 \text{ to } 2n\}$ is the basis for the cocharacter lattice of G_{4n} as in section 3.3, and $\{\bar{e}_i^*, i = 0 \text{ to } 2n - \ell - 1\}$ is the basis for that of $G_{4n-2\ell-1}$, then

$$(6.1.3) \quad \text{inc} \circ \bar{e}_i^* = \begin{cases} e_0^*, & i = 0 \\ e_{\ell+i}^*, & i = 1 \text{ to } 2n - \ell - 1. \end{cases}$$

In the case when $\ell = 2n - 1$, $N_{\ell} = U_{\max}$, and ψ_{ℓ} is a generic character. The above remarks remain valid with the convention that $G_1 = GL_1$.

Let

$$DC_{\omega}(\tau) = FC^{\psi_{n-1}} \mathcal{E}_{-1}(\tau, \omega).$$

It is a space of smooth functions $G_{2n+1}(F \backslash \mathbb{A}) \rightarrow \mathbb{C}$, and affords a representation of the group $G_{2n+1}(\mathbb{A})$ acting by right translation, where we have identified G_{2n+1} with the identity component of $L_{n-1}^{\psi_{n-1}}$.

Theorem 6.1.4. *Let ω be a Hecke character. Let $\tau = \tau_1 \boxplus \cdots \boxplus \tau_r$ be an isobaric sum of $\bar{\omega}$ -symplectic irreducible cuspidal automorphic representations τ_1, \dots, τ_r , of $GL_{2n_1}(\mathbb{A}), \dots, GL_{2n_r}(\mathbb{A})$, respectively. The space $DC_{\omega}(\tau)$ is a nonzero cuspidal representation of $G_{2n+1}(\mathbb{A})$, which supports a nonzero Whittaker integral. If σ is any irreducible automorphic representation contained in $DC_{\omega}(\tau)$, then σ lifts weakly to τ under the map r . Also, the central character of σ is ω .*

Remark 6.1.5. *Since $DC_{\omega}(\tau)$ is nonzero and cuspidal, there exists at least one irreducible component σ . In the case of orthogonal groups, one may show ([So1], pp. 8-9, item 4) that all of the components are generic using the Rankin-Selberg integrals of [Gi-PS-R], [So2]. On the other hand, in the odd case, one may also show ([GRS4], Theorem 8, p. 757, or [So1] page 9, item 6) using the results of [Ji-So] that $DC_{\omega}(\tau)$ is irreducible.*

Proof. The statements are proved by combining relationships between unipotent periods and knowledge about $\mathcal{E}_{-1}(\tau, \omega)$.

For genericity, let (U_1, ψ_1) denote the unipotent period obtained by composing the one which defines the descent with the one which defines the Whittaker function on G_{2n+1} embedded into G_{4n} as the stabilizer of the descent character. Thus U_1 is the subgroup of the standard maximal unipotent defined by the relations $u_{i,2n} = u_{i,2n+1}$ for $i = n$ to $2n - 1$, and

$$\psi_1(u) = \psi(u_{1,2} + \cdots + u_{n-2,n-1} + u_{n-1,2n} - u_{n-1,2n+1} + u_{n,n+1} + \cdots + u_{2n-1,2n}).$$

Next, let U_2 denote the subgroup of the standard maximal unipotent defined by $u_{i,i+1} = 0$ for i even and less than $2n$. (One may also put $\leq 2n$: the equation $u_{2n,2n+1} = 0$ is automatic for any element of U_{\max} .) The character ψ_2 depends on whether n is odd or even. If n is even, it is

$$\psi(u_{1,3} + u_{2,4} + \cdots + u_{2n-1,2n+1}),$$

while, if n is odd, it is

$$\psi(u_{1,3} + u_{2,4} + \cdots + u_{2n-3,2n-1} + u_{2n-2,2n+1} + u_{2n-1,2n}),$$

Finally, let U_3 denote the maximal unipotent, and ψ_3 denote

$$\psi_3(u) = \psi(u_{1,2} + \cdots + u_{2n-1,2n}).$$

Thus (U_3, ψ_3) is the composite of the unipotent period defining the constant term along the Siegel parabolic, and the one which defines the Whittaker functional on the Levi of this parabolic. By Theorem 5.0.4 (5) this period is *not* in $\mathcal{U}^\perp(\mathcal{E}_{-1}(\tau, \omega))$.

In the appendices, we show

- (1) $(U_1, \psi_1)|(U_2, \psi_2)$, in Lemma 9.3.1, and
- (2) $(U_3, \psi_3) \in \langle (U_2, \psi_2), \{(N_\ell, \vartheta) : n \leq \ell < 2n \text{ and } \vartheta \text{ in general position.}\} \rangle$ in Lemma 9.3.2.

By Theorem 6.1.1 $(N_\ell, \vartheta) \in \mathcal{U}^\perp(\mathcal{E}_{-1}(\tau, \omega))$ for all $n \leq \ell < 2n$ and ϑ in general position. It follows that $(U_1, \psi_1) \notin \mathcal{U}^\perp(\mathcal{E}_{-1}(\tau, \omega))$. This establishes genericity (and hence nontriviality) of the descent.

Turning to cuspidality, we prove in the appendices an identity relating:

- Constant terms on G_{2n+1} embedded as $(L_{n-1}^{\psi_{n-1}})^0$,
- Descent periods in G_{4n} ,
- Constant terms on G_{4n} ,
- Descent periods on G_{4n-2k} , embedded in G_{4n} as a subgroup of a Levi.

To formulate the exact relationship we introduce some notation for the maximal parabolics of GSpin groups.

The group G_{2n+1} has one standard maximal parabolic having Levi $GL_i \times G_{2n-2i+1}$ for each value of i from 1 to n . Let us denote the unipotent radical of this parabolic by V_i^{2n+1} . We denote the trivial character of any unipotent group by $\mathbf{1}$.

The group G_{4n} has one standard maximal parabolic having Levi $GL_k \times G_{4n-2k}$ for each value of k from 1 to $2n - 2$. We denote the unipotent radical of this parabolic by V_k .

(The group G_{4n} also has two parabolics with Levi isomorphic to $GL_{2n} \times GL_1$, but since they will not come up in this discussion, we do not need to bother over a notation to distinguish them.)

We prove in Lemma 9.3.4 that $(V_k^{2n+1}, \mathbf{1}) \circ (N_{n-1}, \psi_{n-1})$ is contained in

$$\langle (N_{n+k-1}, \psi_{n+k-1}), \{(N_{n+j-1}, \psi_{n+j-1})^{(4n-2k+2j)} \circ (V_{k-j}, \mathbf{1}) : 1 \leq j < k\} \rangle,$$

where $(N_{n+j-1}, \psi_{n+j-1})^{(4n-2k+2j)}$ denotes the descent period, defined as above, but on the group $G_{4n-2k+2j}$, embedded into G_{4n} as a component of the Levi with unipotent radical V_{k-j} .

By Theorem 6.1.1 $(N_{n+k-1}, \psi_{n+k-1}) \in \mathcal{U}^\perp(\mathcal{E}_{-1}(\tau, \omega))$ for $k = 1$ to n . Furthermore, for k, j such that $1 \leq j < k \leq n$, the function $E(f)(s)^{(V_{k-j}, \mathbf{1})}$ may be expressed in terms of Eisenstein series on GL_{k-j} and $G_{4n-2k+2j}$ using Proposition II.1.7 (ii) of [MW1]. What we require is the following:

Lemma 6.1.6. For all $f \in V^{(2)}(\underline{s}, \bigotimes_{i=1}^r \tau \boxtimes \omega)$

$$E_{-1}(f)^{(V_{k-j}, \mathbf{1})} \Big|_{G_{4n-2k+2j}(\mathbb{A})} \in \bigoplus_S \mathcal{E}_{-1}(\tau_S, \omega),$$

where the sum is over subsets S of $\{1, \dots, r\}$ such that $\sum_{i \in S} 2n_i = 2n - k + j$, and, for each such S , $\mathcal{E}_{-1}(\tau_S, \omega)$ is the space of functions on $G_{4n-2k+2j}(\mathbb{A})$ obtained by applying the construction of $\mathcal{E}_{-1}(\tau, \omega)$ to $\{\tau_i : i \in S\}$, instead of $\{\tau_i : 1 \leq i \leq r\}$.

Once again, this is immediate from [MW1] Proposition II.1.7 (ii).

Applying Theorem 6.1.1, with τ replaced by τ_S and $2n$ by $2n - k + j$, we deduce

$$(N_{n+j-1}, \psi_{n+j-1})^{(4n-2k+2j)} \in \mathcal{U}^\perp(\mathcal{E}_{-1}(\tau_S, \omega)) \quad \forall S,$$

and hence $(N_{n+j-1}, \psi_{n+j-1})^{(4n-2k+2j)} \circ (V_{k-j}, \mathbf{1}) \in \mathcal{U}^\perp(\mathcal{E}_{-1}(\tau, \omega))$. This shows that any nonzero function appearing in any of the spaces $DC_\omega^a(\tau)$ must be cuspidal. Such a function is also easily seen to be of uniformly moderate growth, being the integral of an automorphic form over a compact domain. In addition, such a function is easily seen to have central character ω , and any function with these properties is necessarily square integrable modulo the center ([MW1] I.2.12). It follows that each of the spaces $DC_\omega^a(\tau)$ decomposes discretely.

Now, suppose $\sigma \cong \otimes'_v \sigma_v$ is an irreducible representation which is contained in $DC_\omega(\tau)$. Let p_σ denote the natural projection $DC_\omega(\tau) \rightarrow \sigma$. Once again, by Theorem 5.0.4, (3) the representation $\mathcal{E}_{-1}(\tau, \omega)$ decomposes discretely. Let π be an irreducible component of $\mathcal{E}_{-1}(\tau, \omega)$ such that the restriction of $p_\sigma \circ FC$ to π is nontrivial. As discussed previously in the proof of Theorem 6.1.1, at all but finitely many v , τ is unramified at v and furthermore, π_v is the unramified constituent ${}^{un}Ind_{P(F_v)}^{G_{4n}(F_v)} \tau_v \boxtimes \omega_v \otimes |\det|_{\frac{1}{2}}$ of ${}^{un}Ind_{P(F_v)}^{G_{4n}(F_v)} \tau_v \boxtimes \omega_v \otimes |\det|_{\frac{1}{2}}$. If v_0 is such a place, the map $p_\sigma \circ FC \circ i_{\zeta v_0}$, with $i_{\zeta v_0}$ defined as in Theorem 6.1.1, factors through $\mathcal{J}_{N_{n-1}, \psi_{n-1}} \left({}^{un}Ind_{P(F_v)}^{G_{4n}(F_v)} \tau_v \otimes |\det|_{\frac{1}{2}} \boxtimes \omega_v \right)$, and gives rise to a $G_{2n+1}(F_{v_0})$ -equivariant map from this Jacquet-module onto σ_{v_0} .

To pin things down precisely, assume that τ_v is the unramified component of $Ind_{B(GL_{2n})(F_v)}^{GL_{2n}(F_v)} \mu$, and let μ_1, \dots, μ_{2n} be defined as in the proof of Lemma 4.1.1. By Lemma 4.1.1, we may assume without loss of generality that $\mu_{2n+1-i} = \omega \mu_i^{-1}$ for $i = 1$ to n .

We also need to refer to the elements of the basis of the cocharacter lattice of G_{2n+1} fixed in section 3.3. As in the remarks preceding the definition of $DC_\omega(\tau)$, we denote these $\bar{e}_0^*, \dots, \bar{e}_n^*$.

In the appendices, we show that

$$\mathcal{J}_{N_{n-1}, \psi_{n-1}} \left({}^{un}Ind_{P(F_v)}^{G_{4n}(F_v)} \tau_v \boxtimes \omega_v \otimes |\det|_{\frac{1}{2}} \right)$$

is isomorphic as a $G_{2n+1}(F_v)$ -module to $Ind_{B(G_{2n+1})(F_v)}^{G_{2n+1}(F_v)} \chi$ for χ the unramified character of $B(G_{2n+1})(F_v)$ such that

$$\chi \circ \bar{e}_i^* = \mu_i, i = 1 \text{ to } n, \chi \circ \bar{e}_0^* = \omega_v.$$

It follows that τ is a weak lift of σ associated to the map r . □

7. APPENDIX I: EISENSTEIN SERIES

In this appendix we complete the proofs of several intermediate statements used in the proof of Theorem 5.0.4. As far as we know, all of these results are well-known to the experts, but do not appear in the literature in the precise form we need.

7.1. Proof of Proposition 5.0.10. First, suppose that a set D of hyperplanes carries all the singularities of all the intertwining operators $M(w, \underline{s})f$. Then it follows from [MW1] II.1.7, IV.1.9 (b) that it carries all the singularities of the cuspidal components of all the constant terms of $E(f)(g)(\underline{s})$. By I.4.10, it therefore carries the singularities of the Eisenstein series itself.

On the other hand, it is clear that a set which carries the singularities of the Eisenstein series carries those of all of its constant terms. Thus, what we need to prove is:

Lemma 7.1.1. *Fix M' a standard Levi which is conjugate to M and $\alpha \in \Phi^+(Z_M)$. Let H be the root hyperplane given by $\langle \alpha^\vee, \underline{s} \rangle = c$, $c \neq 0$. Consider the family of functions $M(w, \underline{s})f$ corresponding to $\{w \in W(M) | wMw^{-1} = M'\}$. If any one or them has a pole along H , then the constant term of the Eisenstein series along P' does as well. In other words, it is not possible for two poles to cancel one another.*

Proof. Clearly, it is enough to prove this under the additional hypothesis that $M' = M$.

Let A_M^+ denote the group isomorphic to $(\mathbb{R}_+^\times)^{r+1}$, embedded diagonally at the infinite places, which is inside the center of M .

The Lie algebra of A_M^+ is naturally identified with the real dual of $X(M) \otimes_{\mathbb{Z}} \mathbb{R}$. Recall that above we identified \underline{s} with an element of $X(M) \otimes_{\mathbb{Z}} \mathbb{C}$. So, there is a natural pairing $\langle X, \underline{s} \rangle$, $X \in \mathfrak{a}_M^+$, given as follows. Write \det_i for the determinant of the i th block of an element of M , regarded as a $2n \times 2n$ matrix via the identification with $GL_m \times GL_1$ fixed above. Then we have

$$\prod_{i=1}^r |\det_i \exp(\log y \cdot X)|^{s_i} = y^{\langle X, \underline{s} \rangle}.$$

It follows that

$$|M(w, \underline{s})f(\exp(\log y \cdot X)g)| = y^{\operatorname{Re}(\langle w^{-1}X, \underline{s} \rangle)} \cdot \delta_P^{\frac{1}{2}}(w^{-1} \exp(\log y \cdot X)w) \cdot |M(w, \underline{s})f(g)|.$$

Here δ_P is the modular quasicharacter of P .

Let

$$W_{\text{sing}}(M, H) = \{w \in W(M), wMw^{-1} = M, M(w, \underline{s}) \text{ has a pole along } H\}.$$

Suppose that this set is nonzero. Choose $w_0 \in W_{\text{sing}}(M, H)$ such that the order of the pole of $M(w_0, \underline{s})$ is of maximal order. Let $\nu(H)$ denote the order. Choose $X \in \mathfrak{a}_M^+$ such that the points $w^{-1} \cdot X, w \in W_{\text{sing}}(M, H)$ are all distinct. Consider the family of functions

$$(\langle \alpha^\vee, \underline{s} \rangle - c)^{\nu(H)} M(w, \underline{s})f(\exp(\log y \cdot X)g), \quad w \in W_{\text{sing}}(M, H).$$

They have singularities carried by a locally finite set of root hyperplanes not containing H . Assume g has been chosen so that $(\langle \alpha^\vee, \underline{s} \rangle - c)^{\nu(H)} M(w_0, \underline{s})f(g) \neq 0$. For \underline{s} restricted to an open subset of H not intersecting any of the singular hyperplanes we obtain a family of holomorphic functions, at least one of which is nonzero. If we further exclude the intersection of H with the hyperplanes

$$\langle w_1^{-1}X - w_2^{-1}X, \underline{s} \rangle = 0, \quad w_1, w_2 \in W_{\text{sing}}(M, H),$$

(which can not coincide with H because $c \neq 0$), then at every point \underline{s} , those functions which are nonzero all have distinct orders of magnitude as functions of y . Hence they can not possibly cancel one another. \square

7.2. Proof of Lemma 5.0.12. Regarding $w_i \cdot \underline{s} + \rho_{P_{\alpha_i}}$ as an element of $X(w_i M w_i^{-1}) \otimes_{\mathbb{Z}} \mathbb{C}$, we may decompose it as $\mu_1 + \langle \alpha_i^\vee, w_i \cdot \underline{s} \rangle \tilde{\alpha}_i$, where $\tilde{\alpha}_i$ is defined by the property that

$$\langle \alpha^\vee, \tilde{\alpha}_i \rangle = \delta_{\alpha, \alpha_i}, \quad \text{for } \alpha \in \Phi^+(Z_{w_i M w_i^{-1}}).$$

Then it follows easily from the definitions that μ_1 is in the image of the natural projection $X(M_{\alpha_i}) \otimes_{\mathbb{Z}} \mathbb{C} \rightarrow X(w_i M w_i^{-1}) \otimes_{\mathbb{Z}} \mathbb{C}$ corresponding to restriction of characters of $M_{\alpha_i}(\mathbb{A})$ to $w_i M w_i^{-1}(\mathbb{A})$.

Take f a K -finite flat section of $\text{Ind}_{P^{w_i}(\mathbb{A})}^{G_{4n}(\mathbb{A})}(\otimes_{j=1}^r \tau_j \otimes |\det_j|^{s_j} \boxtimes \omega) \circ \text{Ad}(w_i^{-1})$. Then $M_{w_i}(s_{\alpha_i}, w_i \cdot \underline{s})f$ resides in a finite dimensional subspace of $\text{Ind}_{P^{w_{i-1}}(\mathbb{A})}^{G_{4n}(\mathbb{A})}(\otimes_{j=1}^r \tau_j \otimes |\det_j|^{s_j} \boxtimes \omega) \circ \text{Ad}(w_{i-1}^{-1})$, corresponding to a finite set of K -types determined by f . Write $M_{w_i}(s_{\alpha_i}, w_i \cdot \underline{s})f$ in terms of a basis of flat K -finite sections. The coefficients are functions of \underline{s} , but it follows easily from the integral definition where this is valid, and by meromorphic continuation elsewhere, that in fact they are independent of μ_1 (which corresponds to a character of $M_{\alpha_1}(\mathbb{A})$ and may be pulled out of the integration). Thus, they depend only on $\langle w_i \cdot \underline{s}, \alpha_i^\vee \rangle = \langle \underline{s}, w_i^{-1} \alpha_i^\vee \rangle$.

The first two assertions are now clear. A proof that $c \neq 0$ is obtained by a straightforward modification of the opening paragraph of [MW1], IV.3.12.

7.3. Proof of Proposition 5.0.13. In this section, we denote by $V^{(i)}(s, \tau, \omega)$, $i = 1, 2$, the spaces of functions previously introduced in section 5 as $V^{(i)}(\underline{s}, \otimes_{i=1}^r \tau_i \boxtimes \omega)$, in the special case when $r = 1$.

Let $\tilde{M}(s)$ denote the analogue of $M(w, s)$ defined using $V^{(1)}(s, \tau, \omega)$. It maps into the space $V^{(3)}(-s, \tilde{\tau} \otimes \omega, \omega)$ given by

$$\left\{ \tilde{F} : G_{2m}(\mathbb{A}) \rightarrow V_\tau, \text{ smooth } \left| \tilde{F}((g, \alpha)h)(g_1) = \omega(\alpha \det g) |\det g|^{-s + \frac{(m-1)}{2}} \tilde{F}(h)(g_1 {}_t g^{-1}) \right. \right\}.$$

Fix realizations of the local induced representations τ_v and an isomorphism $\iota : \otimes'_v \tau_v \rightarrow \tau$. Define, for each v , $V^{(1)}(s, \tau_v, \omega_v)$ to be

$$\left\{ \tilde{F}_v : G_{2m}(F_v) \rightarrow V_{\tau_v}, \text{ smooth } \left| \tilde{F}_v((g, \alpha)h) = \omega_v(\alpha) |\det g|_v^{s + \frac{(m-1)}{2}} \tau_v(g) \tilde{F}_v(h) \right. \right\},$$

and $V^{(3)}(s, \tilde{\tau}_v \otimes \omega_v, \omega_v)$ to be

$$\left\{ \tilde{F}_v : G_{2m}(F_v) \rightarrow V_{\tau_v}, \text{ smooth } \left| \tilde{F}_v((g, \alpha)h) = \omega_v(\alpha \det g) |\det g|_v^{s + \frac{(m-1)}{2}} \tau_v({}_t g^{-1}) \tilde{F}_v(h) \right. \right\}.$$

Then the formula

$$\tilde{\iota}(\otimes'_v \tilde{F}_v)(g) = \iota(\otimes'_v \tilde{F}_v(g_v))$$

defines maps

$$\otimes'_v V^{(1)}(s, \tau_v, \omega_v) \rightarrow V^{(1)}(s, \tau, \omega),$$

$$\otimes'_v V^{(3)}(s, \tilde{\tau}_v \otimes \omega_v, \omega_v) \rightarrow V^{(3)}(s, \tilde{\tau} \otimes \omega, \omega),$$

both of which we denote by $\tilde{\iota}$.

It is known that each map is, in fact, an isomorphism. For the benefit of the reader we sketch an argument. On pp. 307 of [Sha1] certain explicit elements of (a generalization of) $V^{(1)}(s, \tau, \omega)$ are constructed as integrals involving matrix coefficients. Using Schur orthogonality, one may check that \tilde{F} is expressible in this form iff both the K -module it generates and the $K \cap M(\mathbb{A})$ -module it generates are irreducible. It is clear that such vectors span the space of all K -finite vectors. On the other hand the (finite dimensional) space of matrix coefficients of this irreducible representation of K is spanned by those that factor as a product of matrix coefficients of local representations, and these are clearly in the image of $\tilde{\iota}$.

For $\tilde{F}_v \in V^{(1)}(s, \tau_v, \omega_v)$, let

$$A_v(s) \tilde{F}_v(g) = \int_{U_w(F_v)} \tilde{F}_v(wug) du.$$

Then the following diagram commutes

$$\begin{array}{ccc} \otimes'_v V^{(1)}(s, \tau_v, \omega_v) & \xrightarrow{A(s)} & \otimes'_v V^{(1)}(-s, \tau_v, \omega_v) \\ \tilde{\iota} \downarrow & & \tilde{\iota} \downarrow \\ V^{(1)}(s, \tau, \omega) & \xrightarrow{\tilde{M}(s)} & V^{(1)}(-s, \tau, \omega) \end{array}$$

with $A(s) := \otimes_v A_v(s)$.

Now, $M(w, s)f(s)$ has a pole (i.e., there exists $g \in G_{4n}(\mathbb{A})$ such that $M(w, s)f(s)(g)$ has a pole) if and only if $\tilde{M}(s)\tilde{F}(s)$ has a pole (i.e., there exist $g \in G_{4n}(\mathbb{A})$ and $m \in M(\mathbb{A})$ such that $\tilde{M}(s)\tilde{F}(s)(g)(m)$ has a pole), where \tilde{F} is the element of $V^{(1)}(s, \tau, \omega)$ such that $f(g) = \tilde{F}(g)(id)$.

We wish to show that there exists \tilde{F} such that this is the case iff τ is $\bar{\omega}$ -symplectic. Clearly, we may restrict attention to \tilde{F} of the form $\tilde{\iota}(\otimes_v \tilde{F}_v)$.

Recall that for all but finitely many non-archimedean v , the space V_{τ_v} comes equipped with a choice of $GL_{2n}(\mathfrak{o}_v)$ -fixed vector ξ_v° used to define the restricted tensor product.

If $\tilde{F} = \tilde{\iota}(\otimes_v \tilde{F}_v) \in V^{(1)}(s, \tau, \omega)$, then there is a finite set S of places, such that if $v \notin S$ then v is non-archimedean, τ_v is unramified, and $\tilde{F}_v(s) = \tilde{F}_{(s, \tau_v, \omega_v)}^\circ$ is the unique element of $V^{(1)}(s, \tau_v, \omega_v)$ satisfying $\tilde{F}_{(s, \tau_v, \omega_v)}^\circ(k) = \xi_v^\circ$ for all $k \in G_{4n}(\mathfrak{o}_v)$.

Now

$$A_v(s)\tilde{F}_{(s, \tau_v, \omega_v)}^\circ = \frac{L_v(2s, \tau_v, \wedge^2 \times \bar{\omega}_v)}{L_v(2s+1, \tau_v, \wedge^2 \times \bar{\omega}_v)} \tilde{F}_{(-s, \tilde{\tau}_v \otimes \omega_v, \omega_v)}^\circ.$$

(A proof of this appears in [L1], albeit not in this precise language. See especially pp. 25-27.) Thus,

$$A(s)\tilde{\iota}(\otimes_v \tilde{F}_v) = \frac{L^S(2s, \tau, \wedge^2 \times \bar{\omega})}{L^S(2s+1, \tau, \wedge^2 \times \bar{\omega})} \tilde{\iota} \left(\left(\bigotimes_{v \in S} A_v(s)\tilde{F}_v(s) \right) \otimes \left(\bigotimes_{v \notin S} \tilde{F}_{-s, \tilde{\tau}_v \otimes \omega_v, \omega_v} \right) \right).$$

To complete the proof we must show:

- (i): $A_v(s)$ is holomorphic and nonvanishing (i.e., not the zero operator) on $Ind_{P(\mathbb{A})}^{G_{2m}(\mathbb{A})} \tau \otimes |\det|^s \boxtimes \omega$ at $s = \frac{1}{2}$, for all τ .
- (ii): $L_v(s, \tau_v, \wedge^2 \times \bar{\omega}_v)$ is holomorphic and nonvanishing at $s = 1$, for all τ_v .
- (iii): $L^S(s, \tau, \wedge^2 \times \bar{\omega})$ is holomorphic and nonvanishing at $s = 2$.

Item (iii) is covered by Proposition 7.3 of [Kim-Sh].

Items (i) and (ii) are essentially contained in Proposition 3.6, p. 153 of [Asg-Sha1]. Since what we need is *part* of the same information, presented differently, we repeat the part of the arguments we are using.

The nonvanishing part of (i) is a completely general fact (i.e., holds at least for any Levi of any split reductive group). For example, the only element of the arguments made on p. 813 of [GRS3] which is particular to the situation they consider there (the Siegel of Sp_{4n}) is the precise ratio of L functions appearing in the constant term.

Similarly, local L functions never vanish. At a finite prime the local L function is $P(q_v^{-s})^{-1}$ for some polynomial P , while at an infinite prime it is given in terms of the Γ function and functions of exponential type.

We turn to holomorphicity.

Lemma 7.3.1. *Let π_v be any representation of $GL_m(F_v)$, which is irreducible, generic, and unitary. Then there exist*

- integers k_1, \dots, k_r of such that $k_1 + \dots + k_r = m$,
- real numbers $\alpha_1, \dots, \alpha_r \in (-\frac{1}{2}, \frac{1}{2})$,

• discrete series representations δ_i of $GL_{k_i}(F_v)$ for $i = 1$ to r
such that

$$\pi_v \cong \text{Ind}_{P_{(k)}(F_v)}^{GL_m(F_v)} \bigotimes_{i=1}^r (\delta_i \otimes |\det_i|^{\alpha_i}).$$

Here $P_{(k)}$ denotes the standard parabolic of GL_m with Levi consisting of block diagonal matrices with the block sizes k_1, \dots, k_r (in that order), and \det_i denotes the determinant of the i th block.

Remark 7.3.2. In fact, one may prove a much more precise statement, but the above is what is needed for our purposes.

Proof. This follows from the main theorem of [Tad2] (see p. 3) together with the fact that the representation denoted $u(\delta, m)$ in that paper is only generic if $m = 1$. For this latter statement see the “Proof of (a) \Rightarrow (f)” on p. 93 of [Vog] in the Archimedean case and Theorem 8.1 on p. 195 of [Z] in the non-Archimedean case. (For the notion of “highest derivative” see p. 452 of [BZ2]: a representation is generic iff its “highest derivative” is the trivial representation of the trivial group, which corresponds to the empty multiset under the Zelevinsky classification.) \square

Continuing with the proof of Proposition 5.0.13, let $(k) = (k_1, \dots, k_r)$, $\delta = (\delta_1, \dots, \delta_r)$ and $\alpha = (\alpha_1, \dots, \alpha_r)$ be obtained from τ_v as just above, and let $\tilde{P}_{(k)}$ denote the standard parabolic of G_{2m} which is contained in the Siegel parabolic P such that $\tilde{P}_{(k)} \cap M = P_{(k)}$.

Then

$$\text{Ind}_{P(F_v)}^{G_{2m}(F_v)} \tau_v \otimes |\det|_v^s \boxtimes \omega_v^s \cong \text{Ind}_{\tilde{P}_{(k)}(F_v)}^{G_{2m}(F_v)} \bigotimes_{i=1}^r (\delta_i \otimes |\det_i|_v^{s+\alpha_i}) \boxtimes \omega_v.$$

This family (as s varies) of representations lies inside the larger family,

$$\text{Ind}_{\tilde{P}_{(k)}(F_v)}^{G_{2m}(F_v)} \bigotimes_{i=1}^r (\delta_i \otimes |\det_i|^{s_i}) \boxtimes \omega_v \quad s = (s_1, \dots, s_r) \in \mathbb{C}^r,$$

and our intertwining operator $A_v(s)$ is the restriction, to the line $s_i = s + \alpha_i$ of the standard intertwining operator for this induced representation, which we denote $A_v(\underline{s})$. This operator is defined, for all $\text{Re}(s_i)$ sufficiently large, by the same integral as $A_v(s)$.

A result of Harish-Chandra says that “ $\text{Re}(s_i)$ sufficiently large” can be sharpened to “ $\text{Re}(s_i) > 0$.” (This is because all δ_i are discrete series, although tempered would be enough.) This result is given in the p -adic case as [Sil] Theorem 5.3.5.4, and in the Archimedean case, [Kn] Theorem 7.22, p. 196.

Hence, the integral defining $A_v(s)$ converges for $s > \max_i(-\alpha_i)$, and in particular converges at $\frac{1}{2}$.

From the relationship between the local L functions and the so-called local coefficients, it follows that the local L functions are also holomorphic in the same region. For this relationship see [Sha3] for the Archimedean case and [Sha2], p. 289 and p. 308 for the non-Archimedean case.

This completes the proof of (i) and (ii).

7.4. Proof of Proposition 5.0.15. The proof is the same as the previous proposition, except that the ratio of partial L function which emerges from the intertwining operators at the unramified places is

$$\frac{L^S(s_1 - s_2, \tau_1 \times \tilde{\tau}_2)}{L^S(s_1 - s_2 + 1, \tau_1 \times \tilde{\tau}_2)}.$$

Convergence of local L functions and intertwining operators at $s_1 - s_2 = 1$ follows again from Lemma 7.3.1. The only difference is the reference for (iii), which in this case is Theorem 5.3 on p. 555 of [Ja-Sh2].

7.5. **Proof of 5.0.17.** As noted, this material is mostly due to Shahidi.

Since the statement is true (with the same proof) for general m , not only $m = 2n$, we prove it in that setting.

In this subsection only, we write τ for the irreducible unitary cuspidal representation $\bigotimes_{i=1}^r \tau_i$ of $M(\mathbb{A})$ (as opposed to the isobaric representation $\tau_1 \boxplus \cdots \boxplus \tau_r$).

First, observe that the integral in question is clearly absolutely and uniformly convergent, and as such defines a meromorphic function of \underline{s} for each g with poles contained in the set of poles of the Eisenstein series itself.

For \underline{s} in the domain of convergence

$$(7.5.1) \quad \int_{U_{\max}^{GL_m}(F \backslash \mathbb{A})} E^{GL_m}(f)(ug)(\underline{s}) \psi_W(u) du = \int_{U_{w_1}(\mathbb{A}) \cdot U^{w_1}(F \backslash \mathbb{A})} f(\underline{s})(w_1^{-1}ug) \psi_W(u) du,$$

where w_1 is the longest element of $W_{GL_m}(\bar{M})$ (defined analogously to $W(M)$ above), $U_{w_1} = U_{\max}^{GL_m} \cap w_1 \bar{U}_{\max}^{GL_m} w_1^{-1}$ and $U^{w_1} = U_{\max}^{GL_m} \cap w_1 U_{\max}^{GL_m} w_1^{-1}$.

Indeed,

$$\bar{P}(F) \backslash GL_m(F) = \coprod_w w^{-1} U_w(F),$$

where the union is over w of minimal length in $w W_{\bar{M}}$. Telescoping, we obtain a sum of terms similar to the right hand side of (7.5.1) for these w . Let $U_{\max}^M = M \cap U_{\max}$. Observe that $w U_{\max}^M w^{-1} \subset U_{\max}$ for all such w . The restriction of ψ_W to $w U_{\max}^M w^{-1}$ is a generic character iff $w M w^{-1}$ is a standard Levi. If it is not, the term corresponding to w vanishes by cuspidality of τ .

On the other hand, $f(w^{-1}ug)$ vanishes if $w^{-1}U_\alpha w$ is contained in the unipotent radical of \bar{P} (which we denote $U_{\bar{P}}$) for any simple root α . Here U_α denotes the one-dimensional unipotent subgroup corresponding to the root α . The element w_1 is the only element of $W_{GL_m}(\bar{M})$ such that this does not hold for any α .

Let λ denote the Whittaker functional on V_τ given by

$$\varphi \mapsto \int_{U_{\max}^M(F \backslash \mathbb{A})} \varphi(u) \psi_W(w_1 u w_1^{-1}) du.$$

Then (7.5.1) equals

$$(7.5.2) \quad \int_{U_{w_1}(\mathbb{A})} \lambda(\tilde{f}(\underline{s})(ug)) \psi_W(u) du,$$

where $\tilde{f} : GL_m(\mathbb{A}) \rightarrow V_{\otimes \tau_i}$ is given by $\tilde{f}(g)(m) = f(mg) \delta_{\bar{P}}^{-\frac{1}{2}}$. (I.e., \tilde{f} is the element of the analogue of $V^{(1)}(\bigotimes_{i=1}^r \tau_i \boxtimes \omega, \underline{s})$, corresponding to f .)

For each place v there exists a Whittaker functional λ_v on the local representation τ_v such that $\lambda(\otimes_v \xi_v) = \prod_v \lambda_v(\xi_v)$. (A finite product because $\lambda_v(\xi_v^c) = 1$ for almost all v . Cf. [Sha1], §1.2.) The induced representation $\text{Ind}_{\bar{P}(\mathbb{A})}^{GL_m(\mathbb{A})}(\bigotimes_{i=1}^r \tau_i | \det_i |^{s_i})$ is isomorphic to a restricted tensor product of local induced representations $\otimes_v' \text{Ind}_{\bar{P}(F_v)}^{GL_m(F_v)}(\bigotimes_{i=1}^r \tau_{i,v} | \det_i |^{s_i})$. (Cf. section 7.3.) Consider an element \tilde{f} which corresponds to a pure tensor $\otimes_v \tilde{f}_v$ in this factorization. So $\tilde{f}_v(\underline{s})$ is a smooth function $GL_m(F_v) \rightarrow V_{\otimes \tau_{i,v}}$ for each \underline{s} .) Then (7.5.2) equals

$$(7.5.3) \quad \prod_v \int_{U_{w_1}(F_v)} \lambda_v(\tilde{f}(\underline{s})(u_v g_v)) \psi_W(u_v) du_v,$$

whenever each of the local integrals is convergent, and the infinite product is convergent (cf [Tate2] Theorem 3.3.1). By Propositions 3.1 and 3.2 of [Sha4], all of the local integrals are always convergent. (See also Lemma 2.3 and the remark at the end of section 2 of [Sha3].)

It is an application of Theorem 5.4 of [C-S] that the term corresponding to an unramified non-archimedean place v in (7.5.2) is equal to $W_v^\circ(g_v) \cdot \prod_{i < j} L_v(s_i - s_j + 1, \tau_{i,v} \otimes \tilde{\tau}_{j,v})^{-1}$. The convergence of the infinite product is then an elementary exercise, as is the main equation in the statement of our present theorem.

The fact that f may be chosen so that the local Whittaker functions at the places in S do not vanish follows again from Propositions 3.1 and 3.2 of [Sha4] (see also the remark at the end of section 2 of [Sha3]).

8. APPENDIX II: LOCAL RESULTS ON JACQUET FUNCTORS

In this appendix, F is a non-archimedean local field, on which we place the additional technical hypothesis

$$(8.0.4) \quad B(G_{2n-1})(F)G_{2n-1}(\mathfrak{o}) = G_{2n-1}(F),$$

which is known (see [Tits], 3.9, and 3.3.2) to hold at all but finitely many non-Archimedean completions of a number field. Here, G_{2n-1} is identified with $(L_{n-1}^\psi)^0$ is defined as in (6.1.3), and \mathfrak{o} denotes the ring of integers of F .

Proposition 8.0.5. *Let $\tau = \text{Ind}_{B(GL_{2n})(F)}^{GL_{2n}(F)} \mu$, where μ satisfies $\mu \circ e_i^* = \omega \mu \circ e_{2n+1-i}^*$, and let P denote the Siegel parabolic subgroup. Then for $\ell \geq n$ and ϑ in general position, the Jacquet module $\mathcal{J}_{N_\ell, \vartheta}(\text{un Ind}_{P(F)}^{G_{4n}(F)} \tau \otimes |\det|^{\frac{1}{2}} \boxtimes \omega)$ is trivial.*

Proof. First, let $\mu_i : F \rightarrow \mathbb{C}$ be the unramified character given by $\mu_i = \mu \circ e_i^*$. By induction in stages,

$$\text{un Ind}_{P(F)}^{G_{4n}(F)} \tau \otimes |\det|^{\frac{1}{2}} \boxtimes \omega = \text{un Ind}_{B(G_{4n})(F)}^{G_{4n}(F)} \tilde{\mu},$$

where $\tilde{\mu} \circ e_i^*(x) = |x|^{\frac{1}{2}} \mu_i(x)$, for $i = 1$ to $2n$ and $\tilde{\mu} \circ e_0^* = \omega$. By the definition of the unramified constituent

$$\text{un Ind}_{B(G_{4n})(F)}^{G_{4n}(F)} \tilde{\mu} = \text{un Ind}_{B(G_{4n})(F)}^{G_{4n}(F)} \tilde{\mu}',$$

where $\tilde{\mu}' \circ e_{2i-1}^*(x) = \mu_i(x)|x|^{\frac{1}{2}}$, and $\tilde{\mu}' \circ e_{2i}^*(x) = \mu_i(x)|x|^{-\frac{1}{2}}$, for $i = 1$ to n , and $\tilde{\mu}' \circ e_0^* = \omega$. Now, it is well known that

$$\text{un Ind}_{B(GL_2)(F)}^{GL_2(F)} \mu |^{\frac{1}{2}} \otimes \mu |^{-\frac{1}{2}} = \mu \circ \det.$$

It follows that

$$\text{un Ind}_{B(G_{4n})(F)}^{G_{4n}(F)} \tilde{\mu}' = \text{un Ind}_{P_{2n}(F)}^{G_{4n}(F)} \hat{\mu},$$

where P_{2n} is the parabolic of G_{4n} having Levi isomorphic to $GL_2^n \times GL_1$, such that the roots of this Levi are $e_1 - e_2, e_3 - e_4, \dots, e_{2n-1} - e_{2n}$, and $\hat{\mu}$ is the character given by $\hat{\mu} \circ e_{2i-1}^* = \hat{\mu} \circ e_{2i}^* = \mu_i$, $\hat{\mu} \circ e_0^* = \omega$.

The space $\text{Ind}_{P_{2n}(F)}^{G_{4n}(F)} \hat{\mu}$ has a filtration as a $Q_\ell(F)$ -module, in terms of $Q_\ell(F)$ -modules indexed by the elements of $(W \cap P_{2n}) \backslash W / (W \cap Q_\ell)$. For any element x of $P_{2n}(F)wQ_\ell(F)$ the module corresponding to w is isomorphic to $c - \text{ind}_{x^{-1}P_{2n}(F)x \cap Q_\ell(F)}^{Q_\ell(F)} \hat{\mu} \delta_{P_{2n}}^{\frac{1}{2}} \circ \text{Ad}(x)$. Here $\text{Ad}(x)$ denotes the map given by conjugation by x . It sends $x^{-1}P_{2n}(F)x \cap Q_\ell(F)$ into $P_{2n}(F)$. Also, here and throughout $c - \text{ind}$ denotes non-normalized compact induction. (See [Cass], section 6.3.)

Recall from 3.4 that the Weyl group of G_{4n} is identified (canonically after the choice of pr) with the set of permutations $w \in \mathfrak{S}_{4n}$ satisfying,

- (1) $w(4n+1-i) = 4n+1-w(i)$ and
- (2) $\det w = 1$ when w is written as a $4n \times 4n$ permutation matrix.

As representatives for the double cosets $(W \cap P_{2n}) \backslash W / (W \cap Q_\ell)$ we choose the element of minimal length in each. As permutations, these elements have the properties

- (3) $w^{-1}(2i) > w^{-1}(2i-1)$ for $i = 1$ to $2n$, and
(4) If $\ell \leq i < j \leq 4n+1-\ell$ and $w(i) > w(j)$, then $i = 2n$ and $j = 2n+1$.
Let I_w be the $Q_\ell(F)$ -module obtained as

$$c - \text{ind}_{\dot{w}^{-1}P_{2n}(F)\dot{w} \cap Q_\ell(F)}^{Q_\ell(F)} \hat{\mu} \delta_{P_{2n}}^{\frac{1}{2}} \circ \text{Ad}(\dot{w})$$

using any element \dot{w} of $\text{pr}^{-1}(w)$.

A function f in I_w will map to zero under the natural projection to $\mathcal{J}_{N_\ell, \vartheta}(I_w)$ iff there exists a compact subgroup N_ℓ^0 of $N_\ell(F)$ such that

$$\int_{N_\ell^0} f(hn) \overline{\vartheta(n)} dn = 0 \quad \forall h \in Q_\ell(F).$$

(See [Cass], section 3.2.) Let $\vartheta^h(n) = \vartheta(hnh^{-1})$. It is easy to see that the integral above vanishes for suitable N_ℓ^0 whenever

$$(8.0.6) \quad \vartheta^h|_{N_\ell(F) \cap w^{-1}P_{2n}(F)w} \text{ is nontrivial.}$$

Furthermore, the function $h \mapsto \vartheta^h$ is continuous in h , (the topology on the space of characters of $N_\ell(F)$ being defined by identifying it with a finite dimensional F -vector space, cf. section 4.3) so if this condition holds for all h in a compact set, then N_ℓ^0 can be made uniform in h .

Now, ϑ is in general position. Hence, so is ϑ^h for every h . So, if we write

$$\vartheta^h(u) = \psi_0(c_1 u_{1,2} + \cdots + c_{\ell-1} u_{\ell-1,\ell} + d_1 u_{\ell,\ell+1} + \cdots + d_{2m-2\ell} u_{\ell,2m-\ell}),$$

we have that $c_i \neq 0$ for all i and ${}^t d w d \neq 0$.

Clearly, the condition (8.0.6) holds for all h unless

$$(5) \quad w(1) > w(2) > \cdots > w(\ell).$$

Furthermore, because ${}^t d w d \neq 0$, there exists some i_0 with $\ell+1 \leq i_0 \leq 2n$ such that $d_{i_0-\ell} \neq 0$ and $d_{4n+1+\ell-i_0} \neq 0$. From this we deduce that the condition (8.0.6) holds for all h unless w has the additional property

$$(6) \quad \text{There exists } i_0 \text{ such that } w(\ell) > w(i_0) \text{ and } w(\ell) > w(4n+1-i_0).$$

However, if $\ell \geq n$ it is easy to check that no permutations with properties (1),(3) (5) and (6) exist.

Thus $\mathcal{J}_{N_\ell, \vartheta}(I_w) = \{0\}$ for all w and hence $\mathcal{J}_{N_\ell, \vartheta}({}^{un} \text{Ind}_{P(F)}^{G_{4n}(F)} \tau \otimes |\det|^{\frac{1}{2}} \boxtimes \omega) = \{0\}$ by exactness of the Jacquet functor. \square

Proposition 8.0.7. *Let $\tau = \text{Ind}_{B(GL_{2n})(F)}^{GL_{2n}(F)} \mu$, where μ satisfies $\mu \circ e_i^* = \omega \mu \circ e_{2n+1-i}$. Then the Jacquet module*

$$\mathcal{J}_{N_{n-1}, \psi_{n-1}} \left({}^{un} \text{Ind}_{P(F)}^{G_{4n}(F)} \tau \otimes |\det|^{\frac{1}{2}} \boxtimes \omega \right)$$

is isomorphic as a $G_{2n+1}(F)$ -module to a subquotient of $\text{Ind}_{B(G_{2n+1})(F)}^{G_{2n+1}(F)} \chi$ for χ the unramified character of $B(G_{2n+1})(F)$ such that

$$\chi \circ \bar{e}_i^* = \mu_i, i = 1 \text{ to } n, \chi \circ \bar{e}_0^* = \omega.$$

Proof. As before, we have

$${}^{un} \text{Ind}_{P(F)}^{G_{4n}(F)} \tau \otimes |\det|^{\frac{1}{2}} \boxtimes \omega = {}^{un} \text{Ind}_{P_{2n}}^{G_{4n}(F)} \hat{\mu},$$

and we filter $\text{Ind}_{P_{2n}}^{G_{4n}(F)} \hat{\mu}$ in terms of $Q_{n-1}(F)$ -modules I_w . This time, $\mathcal{J}_{N_{n-1}, \psi_{n-1}}(I_w) = \{0\}$ for all w except one. This one Weyl element, which we denote w_0 , corresponds to the unique permutation

satisfying (1),(2),(3),(4) of the previous result, together with $w(i) = 4n - 2i + 1$ for $i = 1$ to $n - 1$. Exactness yields

$$\mathcal{J}_{N_{n-1}, \psi_{n-1}} \left({}^{un} \text{Ind}_{P(F)}^{G_{4n}(F)} \tau \otimes |\det|^{\frac{1}{2}} \boxtimes \omega \right) \cong \mathcal{J}_{N_\ell, \vartheta}(I_{w_0}).$$

(This is an isomorphism of $Q_{n-1}^{\psi_{n-1}}(F)$ -modules, where $Q_{n-1}^{\psi_{n-1}} = N_{n-1} \cdot L_{n-1}^{\psi_{n-1}} \subset Q_{n-1}$, is the stabilizer of ψ_{n-1} in Q_{n-1} (cf. L^ϑ above).)

Now, recall that for each $h \in Q_{n-1}(F)$ the character $\psi_{n-1}^h(u) = \psi_{n-1}(huh^{-1})$ is a character of N_{n-1} in general position, and as such determines coefficients c_1, \dots, c_{n-2} and d_1, \dots, d_{2n+2} as in remark 6.1.2. Clearly,

$$Q_{n-1}^o := \{h \in Q_{n-1}(F) \mid d_i \neq 0 \text{ for some } i \neq n+1, n+2\}$$

is open. Moreover, one may see from the description of w_0 that for h in this set 8.0.6 is satisfied. We have an exact sequence of $Q_{n-1}^{\psi_{n-1}}(F)$ -modules

$$0 \rightarrow I_{w_0}^* \rightarrow I_{w_0} \rightarrow \bar{I}_{w_0} \rightarrow 0,$$

where I_w^* consists of those functions in I_w whose compact support happens to be contained in Q_{n-1}^o , and the third arrow is restriction to the complement of Q_{n-1}^o . This complement is slightly larger than $Q_{n-1}^{\psi_{n-1}}(F)$ in that it contains the full torus of $Q_{n-1}(F)$, but restriction of functions is an isomorphism of $Q_{n-1}^{\psi_{n-1}}(F)$ -modules,

$$\bar{I}_{w_0} \rightarrow c - \text{ind}_{Q_{n-1}^{\psi_{n-1}}(F) \cap w_0^{-1} P_{2n}(F) w_0}^{Q_{n-1}^{\psi_{n-1}}(F)} \hat{\mu} \delta_{P_{2n}}^{\frac{1}{2}} \circ \text{Ad}(w_0).$$

Clearly $\mathcal{J}_{N_{n-1}, \psi_{n-1}}(I_{w_0}^*) = \{0\}$, and hence

$$\mathcal{J}_{N_{n-1}, \psi_{n-1}} \left(\text{Ind}_{P_{2n}(F)}^{G_{4n}(F)} \hat{\mu} \right) \cong \mathcal{J}_{N_{n-1}, \psi_{n-1}} \left(c - \text{ind}_{Q_{n-1}^{\psi_{n-1}}(F) \cap w_0^{-1} P_{2n}(F) w_0}^{Q_{n-1}^{\psi_{n-1}}(F)} \hat{\mu} \delta_{P_{2n}}^{\frac{1}{2}} \circ \text{Ad}(w_0) \right).$$

Now let \mathcal{W} denote

$$\left\{ f : Q_{n-1}^{\psi_{n-1}}(F) \rightarrow \mathbb{C} \left| \begin{array}{l} f(uq) = \psi_{n-1}(u) f(q) \quad \forall u \in N_{n-1}(F), q \in Q_{n-1}^{\psi_{n-1}}(F), \\ f(bm) = \chi(b) \delta_{B(G_{2n+1})}^{\frac{1}{2}} f(m) \quad \forall b \in B(L_{n-1}^{\psi_{n-1}}(F)), m \in L_{n-1}^{\psi_{n-1}}(F) \end{array} \right. \right\}.$$

For $f \in c - \text{ind}_{Q_{n-1}^{\psi_{n-1}}(F) \cap w_0^{-1} P_{2n}(F) w_0}^{Q_{n-1}^{\psi_{n-1}}(F)} \hat{\mu} \delta_{P_{2n}}^{\frac{1}{2}} \circ \text{Ad}(w_0)$, let

$$W(f)(q) = \int_{N_{n-1}(F) \cap w_0^{-1} \overline{U_{\max}(F)} w_0} f(uq) \bar{\psi}_{n-1}(u) du.$$

Then W maps $c - \text{ind}_{Q_{n-1}^{\psi_{n-1}}(F) \cap w_0^{-1} P_{2n}(F) w_0}^{Q_{n-1}^{\psi_{n-1}}(F)} \hat{\mu} \delta_{P_{2n}}^{\frac{1}{2}} \circ \text{Ad}(w_0)$ into \mathcal{W} . That is, the functions in $c - \text{ind}_{Q_{n-1}^{\psi_{n-1}}(F) \cap w_0^{-1} P_{2n}(F) w_0}^{Q_{n-1}^{\psi_{n-1}}(F)} \hat{\mu} \delta_{P_{2n}}^{\frac{1}{2}} \circ \text{Ad}(w_0)$ are left equivariant with respect to the group $B(G_{2n+1})(F)$,

and a quasicharacter of this group that differs from $\chi \delta_{B(G_{2n+1})}^{\frac{1}{2}}$ by the Jacobian of $\text{Ad}(b)$, $b \in B(G_{2n+1})(F)$, acting on $N_{n-1}(F) \cap w_0^{-1} \overline{U_{\max}(F)} w_0$.

Let us denote

$$c - \text{ind}_{Q_{n-1}^{\psi_{n-1}}(F) \cap w_0^{-1} P_{2n}(F) w_0}^{Q_{n-1}^{\psi_{n-1}}(F)} \hat{\mu} \delta_{P_{2n}}^{\frac{1}{2}} \circ \text{Ad}(w_0)$$

by V and denote by $V(N_{n-1}, \psi_{n-1})$ the kernel of the linear map $V \rightarrow \mathcal{J}_{N_{n-1}, \psi_{n-1}}(V)$.

It is easy to show that $V(N_{n-1}, \psi_{n-1})$ is contained in the kernel of W . In the next lemma, we show that in fact, they are equal. Restriction from $Q_{n-1}^{\psi_{n-1}}(F)$ to $L_{n-1}^{\psi_{n-1}}(F)$ is clearly an isomorphism $\mathcal{W} \rightarrow \text{Ind}_{B(G_{2n+1})(F)}^{G_{2n+1}(F)} \chi$. \square

Lemma 8.0.8. *With notation as in the previous proposition, we have $\text{Ker}(W) \subset V(N_{n-1}, \psi_{n-1})$.*

Proof. For this proof, we denote the Borel of $L_{n-1}^{\psi_{n-1}}$ by B . Also, let $N^{w_0} = N_{n-1} \cap w_0^{-1} P_{2n} w_0$, and $N_{w_0} = N_{n-1} \cap w_0^{-1} \overline{U_{\max}} w_0$,

We consider a smooth function $f : Q_{n-1}^{\psi_{n-1}}(F) \rightarrow \mathbb{C}$ which is compactly supported modulo $Q_{n-1}^{\psi_{n-1}}(F) \cap w_0^{-1} P_{2n}(F) w_0$, and satisfies

$$f(bm) = \chi \delta_B^{\frac{1}{2}}(b) f(m) \quad \forall b \in B(F),$$

and

$$f(uq) = f(q) \quad \forall u \in N^{w_0}(F) \text{ and } q \in Q_{n-1}^{\psi_{n-1}}(F).$$

We assume that

$$\int_{N_{w_0}(F)} f(uq) \bar{\psi}_{n-1}(u) du = 0,$$

for all $q \in Q_{n-1}^{\psi_{n-1}}(F)$. What must be shown is that there is a compact subset C of $N_{n-1}(F)$ such that

$$\int_C f(gu) \bar{\psi}_{n-1}(u) du = 0,$$

for all $q \in Q_{n-1}^{\psi_{n-1}}(F)$.

Consider first $m \in L_{n-1}^{\psi_{n-1}}(\mathfrak{o})$. Let \mathfrak{p} denote the unique maximal ideal in \mathfrak{o} . If U is a unipotent subgroup and M an integer, we define

$$U(\mathfrak{p}^M) = \{u \in U(F) : u_{ij} \in \mathfrak{p}^M \forall i \neq j\}.$$

Observe that for each $M \in \mathbb{N}$, $N_{n-1}(\mathfrak{p}^M)$ is a subgroup of $N_{n-1}(F)$ which is preserved by conjugation by elements of $L_{n-1}^{\psi_{n-1}}(\mathfrak{o})$. We may choose M sufficiently large that $\text{supp}(f) \subset N^{w_0}(F) N_{w_0}(\mathfrak{p}^{-M}) L_{n-1}^{\psi_{n-1}}(F)$.

Then we prove the desired assertion with $C = N_{n-1}(\mathfrak{p}^{-M})$. Indeed, for $m \in L_{n-1}^{\psi_{n-1}}(\mathfrak{o})$, we have

$$\int_{N_{n-1}(\mathfrak{p}^{-M})} f(mu) \bar{\psi}_{n-1}(u) du = \int_{N_{n-1}(\mathfrak{p}^{-M})} f(um) \bar{\psi}_{n-1}(u) du,$$

because $Ad(m)$ preserves the subgroup $N_{n-1}(\mathfrak{p}^{-M})$, and has Jacobian 1. Let $c = \text{Vol}(N^{w_0}(\mathfrak{p}^{-M}))$, which is finite. Then by N^{w_0} -invariance of f , the above equals

$$= c \int_{N_{w_0}(\mathfrak{p}^{-M})} f(um) \bar{\psi}_{n-1}(u) du.$$

This, in turn, is equal to

$$= c \int_{N_{w_0}(F)} f(um) \bar{\psi}_{n-1}(u) du,$$

since none of the points we have added to the domain of integration are in the support of f , and this last integral is equal to zero by hypothesis.

Next, suppose $q = u_1 m$ with $u_1 \in N_{n-1}(F)$ and $m \in L_{n-1}^{\psi_{n-1}}(\mathfrak{o})$. If $u_1 \in N_{n-1}(F) - N_{n-1}(\mathfrak{p}^{-M})$ then qu is not in the support of f for any $u \in N_{n-1}(\mathfrak{p}^{-M})$. On the other hand, if $u_1 \in N_{n-1}(\mathfrak{p}^{-M})$, then

$$\int_{N_{n-1}(\mathfrak{p}^{-M})} f(u_1 m u) \bar{\psi}_{n-1}(u) du = \int_{N_{n-1}(\mathfrak{p}^{-M})} f(u_1 u m) \bar{\psi}_{n-1}(u) du$$

$$= \psi_{n-1}(u_1) \int_{N_{n-1}(\mathfrak{p}^{-M})} f(um) \bar{\psi}_{n-1}(u) du,$$

and now we continue as in the case $u_1 = 1$.

The result for general q now follows from the left-equivariance properties of f and (8.0.4). \square

9. APPENDIX III: IDENTITIES OF UNIPOTENT PERIODS

9.1. A Lemma Regarding Unipotent Periods. We return briefly to the general setting of section 4.3. There is a natural action of $G(F)$ on \mathcal{U} given by $\gamma \cdot (U, \psi) = (\gamma U \gamma^{-1}, \gamma \cdot \psi)$ where $\gamma \cdot \psi(u) = \psi(\gamma^{-1} u \gamma)$. We shall refer to this action as “conjugation.” Obviously, unipotent periods which are conjugate are equivalent.

In the special case $G = G_{4n}$, it is convenient to allow ourselves to conjugate our unipotent periods by elements of the slightly larger group Pin_{4n} . We may allow the involution \dagger to act on unipotent periods by $f^{\dagger(U, \psi_U)}(g) = f^{(U, \psi_U)}(\dagger g)$. Denoting the action of $\text{Pin}_{4n}(F)$ on \mathcal{U} by $\gamma \cdot (U, \psi_U)$, we have

$$\gamma \cdot (U, \psi_U) \sim \begin{cases} (U, \psi_U) & \text{when } \det \text{pr } \gamma = 1, \\ \dagger(U, \psi_U) & \text{when } \det \text{pr } \gamma = -1. \end{cases}$$

Observe that in general $\dagger(U, \psi_U)$ is *not* equivalent to (U, ψ_U) . For example, it is not difficult to verify that $\dagger(U_{\max}, \psi_{LW}) \in \mathcal{U}^{\perp}(\mathcal{E}_{-1}(\tau, \omega))$.

Lemma 9.1.1. *Suppose $U_1 \supset U_2 \supset (U_1, U_1)$ are unipotent subgroups of a reductive algebraic group G . Suppose H is a subgroup of G and let f be a smooth left $H(F)$ -invariant function on $G(\mathbb{A})$. Suppose ψ_2 is a character of U_2 such that $\psi_2|_{(U_1, U_1)} \equiv 0$. Then the set $\text{res}^{-1}(\psi_2)$ of characters of U_1 such that the restriction to U_2 is ψ_2 is nontrivial. (Here “res” is for “restriction” not “residue”.) The elements of $\text{res}^{-1}(\psi_2)$ are permuted by the action of $N_H(U_1)(F)$. The following are equivalent.*

- (1) $f^{(U_2, \psi_2)} \equiv 0$
- (2) $f^{(U_1, \psi_1)} \equiv 0 \forall \psi_1 \in \text{res}^{-1}(\psi_2)$
- (3) For each $N_H(U_1)(F)$ -orbit \mathcal{O} in $\text{res}^{-1}(\psi_2)$ $\exists \psi_1 \in \mathcal{O}$ with $f^{(U_1, \psi_1)} \equiv 0$

Proof. It is obvious that 1 implies 2 and 3, and that 2 and 3 are equivalent. Consider

$$f^{(U_2, \psi_2)}(u_1 g) = \int_{U_2(F \backslash \mathbb{A})} f(u_2 u_1 g) \psi_2(u_2) du_2,$$

regarded as a function of u_1 . It is left u_2 invariant and hence gives rise to a function of the compact abelian group $U_2(\mathbb{A})U_1(F) \backslash U_1(\mathbb{A})$. Denote this function by $\phi(u_1)$. Then

$$\phi(0) = \sum_{\chi} \int_{U_2(\mathbb{A})U_1(F) \backslash U_1(\mathbb{A})} \phi(u_1) \chi(u_1) du_1,$$

where “0” denotes the identity in $U_2(\mathbb{A})U_1(F) \backslash U_1(\mathbb{A})$, and the sum is over characters of $U_2(\mathbb{A})U_1(F) \backslash U_1(\mathbb{A})$. This, in turn, is equal to

$$\sum_{\chi} \int_D \int_{U_2(F \backslash \mathbb{A})} f(u_2 u_1 g) \psi_2(u_2) du_2 \chi(u_1) du_1,$$

for D a fundamental domain for the above quotient in $U_1(\mathbb{A})$. The group $U_1/(U_1, U_1)(F)$ is an F -vector space (cf. sections:uniper) which can be decomposed into $U_2/(U_1, U_1)(F)$ and a complement. The F -dual of this vector space is identified, via the choice of ψ_0 , with the space of characters of $U_1(\mathbb{A})$ which are trivial on $U_1(F)$. It follows that the sum above is equal to

$$= \sum_{\psi_1 \in \text{res}^{-1}(\psi_2)} \int_{U_1(F \backslash \mathbb{A})} f(u_1 g) \psi_1(u_1) du_1.$$

The matter of replacing the sum over χ by one over $\psi_1 \in \text{res}^{-1}(\psi_2)$ is clear from regarding $U_1/(U_1, U_1)(F)$ as a vector space which can be decomposed into $U_2/(U_1, U_1)$ and a complement. Now $2 \Rightarrow 1$ is immediate. \square

Corollary 9.1.2. *If $N_G(H)$ permutes the elements of $\text{res}^{-1}(\psi_2)$ transitively, then $(U_2, \psi_2) \sim (U_2, \psi_1)$ for every $\psi_1 \in \text{res}^{-1}(\psi_2)$.*

Definition 9.1.3. *Many of the applications of the above corollary are of a special type, and it will be convenient to introduce a term for them. The special situation is the following: one has three unipotent periods (U_i, ψ_i) for $i = 1, 2, 3$, such that $U_2 = U_1 \cap U_3$ and $\psi_1|_{U_2} = \psi_3|_{U_2} = \psi_2$. Furthermore, U_1 normalizes U_3 and permutes transitively, the set of characters ψ'_3 such that $\psi'_3|_{U_2}$, and the same is true with the roles of 1 and 3 reversed. In this situation, the identity*

$$(U_1, \psi_1) \sim (U_2, \psi_2) \sim (U_3, \psi_3),$$

*(which follows from Corollary 9.1.2) will be called a **swap**, and we say that (U_1, ψ_1) “may be swapped for” (U_3, ψ_3) , and vice versa.*

9.2. A lemma regarding the projection, and a remark.

Lemma 9.2.1. *The action of G_m on itself by conjugation factors through pr .*

Proof. One has only to check that the kernel of pr is in the center of G_m . When we regard G_m as a quotient of $\text{Spin}_m \times GL_1$, the quotient of pr is precisely the image of the GL_1 factor in the quotient. \square

Corollary 9.2.2. *Let u be a unipotent element of $G_m(\mathbb{A})$ and g any element of $G_m(\mathbb{A})$. Then $\text{pr}(gug^{-1})$ is a unipotent element of $SO_m(\mathbb{A})$ and gug^{-1} is the unique unipotent element of its preimage in $G_m(\mathbb{A})$.*

Remark 9.2.3. *This fact, combined with the fact that pr is an isomorphism of varieties when restricted to the subvariety of unipotent elements of G_m , means that many statements may be proved for $G\text{Spin}$ groups simply by taking the proof of the corresponding statement for special orthogonal groups and inserting the words “any preimage of” here and there.*

9.3. Relations among Unipotent Periods used in Theorem 6.1.4. Before we proceed with the proofs it will be convenient to formulate the statements in a slightly different way, making use of the involution \dagger .

We shall let (U_1, ψ_1) and (U_3, ψ_3) be defined as in the proof of 6.1.4. We also keep the definition of the group U_2 . However, we now define the character ψ_2 by the formula

$$\psi_2(u) = \psi(u_{13} + \cdots + u_{2n-1, 2n+1}),$$

regardless of the parity of n . (This agrees with the previous definition if n is even; if n is odd they differ by an application of \dagger .)

Lemma 9.3.1. *Let (U_1, ψ_1) be defined as in Theorem 6.1.4, and (U_2, ψ_2) defined as just above. Then $(U_1, \psi_1)|(U_2, \psi_2)$ and $(U_1, \psi_1)|^\dagger(U_2, \psi_2)$.*

Proof. We define some additional unipotent periods which appear at intermediate stages in the argument. Let U_4 be the subgroup defined by $u_{n-1, j} = 0$ for $j = n$ to $2n - 2$ and $u_{2n-1, 2n} = u_{2n-1, 2n+1}$. We define a character ψ_4 of U_4 by the same formula as ψ_1 . Then (U_1, ψ_1) may be swapped for (U_4, ψ_4) . (See definition 9.1.3.)

Now, for each k from 1 to n , define $(U_5^{(k)}, \psi_5^{(k)})$ as follows. First, for each k , the group $U_5^{(k)}$ is contained in the subgroup of U_{\max} defined by, $u_{2n-1, 2n} = u_{2n-1, 2n+1}$. In addition, $u_{n+k-2, j} = 0$ for

$j < 2n - 1$, and $u_{i,i+1} = 0$ if $n - k \leq i < n + k$ and $i \equiv n - k \pmod{2}$, and $\psi_5^{(k)}(u)$ equals

$$\psi_0 \left(\sum_{i=1}^{n-k-1} u_{i,i+1} + \sum_{i=n-k}^{n+k-3} u_{i,i+2} + u_{n+k-2,2n} + u_{n+k-2,2n+1} + \sum_{i=n+k-1}^{2n-1} u_{i,i+1} \right).$$

(Note that one or more of the sums here may be empty.)

Next, let $U_6^{(k)}$ be the subgroup of U_{\max} defined by the conditions $u_{2n-1,2n} = u_{2n-1,2n+1}$, $u_{n+k-2,j} = 0$ for $j < 2n - 1$, and $u_{i,i+1} = 0$ if $n - k \leq i < n + k - 2$ and $i \equiv n - k + 1 \pmod{2}$. The same formula which defines $\psi_5^{(k)}$ also defines a character of $U_6^{(k)}$. We denote this character by $\psi_6^{(k)}$.

We make the following observations:

- $(U_5^{(1)}, \psi_5^{(1)})$ is precisely (U_4, ψ_4) .
- For each k , $(U_5^{(k)}, \psi_5^{(k)})$ is conjugate to $(U_6^{(k+1)}, \psi_6^{(k+1)})$. The conjugation is accomplished by any preimage of the permutation matrix which transposes i and $i + 1$ for $n - k \leq i < n + k$ and $i \equiv n - k \pmod{2}$.
- $(U_6^{(k)}, \psi_6^{(k)})$ may be swapped for $(U_5^{(k)}, \psi_5^{(k)})$.

Thus $(U_4, \psi_4) \sim (U_5^{(n)}, \psi_5^{(n)})$.

Now, let ψ_2' be the character of U_2 which is defined by

$$\psi_2'(u) = \psi(u_{1,3} + \cdots + u_{2n-2,2n} - u_{2n-2,2n+1} + u_{2n-1,2n+1}).$$

Then $U_5^{(n)}$ is the subgroup of U_2 defined by $u_{2n-1,2n} = u_{2n-1,2n+1}$ and $\psi_5^{(n)}$ is the restriction of ψ_2' to this group. Thus $(U_5^{(n)}, \psi_5^{(n)}) | (U_2, \psi_2')$. (It is because of this step that $(U_1, \psi_1) \not\sim (U_2, \psi_2)$.)

Finally, (U_2, ψ_2) and (U_2, ψ_2') are conjugate by the unipotent element which projects to $I_{4n} - \sum_{i=2}^n e'_{2i-1,2i-2}$

To obtain $\dagger(U_2, \psi_2)$, we use

$$\psi_2''(u) := \psi(u_{1,3} + \cdots + u_{2n-2,2n} - u_{2n-2,2n+1} + u_{2n-1,2n})$$

instead of ψ_2' . □

Lemma 9.3.2. *Let (U_3, ψ_3) be defined as in Theorem 6.1.4, and let (U_2, ψ_2) be defined as in the previous lemma. Then*

$$(U_3, \psi_3) \in \langle \dagger^n(U_2, \psi_2), \{(N_\ell, \vartheta) : n \leq \ell < 2n \text{ and } \vartheta \text{ in general position.}\} \rangle.$$

Here \dagger^n indicates that we apply \dagger a total of n times, with the effect being \dagger if n is odd and trivial if n is even.

Proof. To prove this assertion we introduce some additional unipotent periods. For $k = 1$ to $2n - 1$ let $U_7^{(k)}$ denote the subgroup of U_{\max} defined by $u_{i,i+1} = 0$ for $i > k$ and $i \equiv k + 1 \pmod{2}$. We use two characters of this group:

$$\begin{aligned} \tilde{\psi}_7^{(k)} &= \psi_0 \left(\sum_{1 \leq i \leq k-1} u_{i,i+1} + \sum_{k \leq i \leq 2n-1} u_{i,i+2} \right), \\ \psi_7^{(k)} &= \psi_0 \left(\sum_{1 \leq i \leq k} u_{i,i+1} + \sum_{k+1 \leq i \leq 2n-1} u_{i,i+2} \right). \end{aligned}$$

Then $(U_7, \psi_7^{(k)})$ is conjugate to $(U_7, \tilde{\psi}_7^{(k)})$ by any preimage of the permutation matrix which transposes i and $i + 1$ for $k < i < 4n - k$ and $i \equiv k + 1 \pmod{2}$. This matrix has determinant -1 iff k is odd.

If k is odd then $(U_7^{(k)}, \psi_7^{(k)})$ may be swapped for $(U_7^{(k+1)}, \tilde{\psi}_7^{(k+1)})$, while if k is even, it may be swapped for $(U_8^{(k+1)}, \tilde{\psi}_8^{(k+1)})$, where $U_8^{(k+1)}$ is the subgroup of $U_7^{(k+1)}$ defined by $u_{2n-1,2n} = 0$, and $\tilde{\psi}_8^{(k+1)}$ is the restriction of $\tilde{\psi}_7^{(k+1)}$ to this group.

Now, for $a \in F^\times$ define a character $\tilde{\psi}_7^{(k+1,a)}$ of $U_7^{(k+1)}$ by

$$\tilde{\psi}_7^{(k+1,a)} = \psi(u_{1,2} + \cdots + u_{k-1,k} + u_{k,k+2} + \cdots + u_{2n-1,2n+1} + au_{2n-1,2n}).$$

Then a Fourier expansion along $U_{2n-1,2n}$ shows that

$$(U_8^{(k+1)}, \tilde{\psi}_8^{(k+1)}) \in \langle (U_7^{(k+1)}, \tilde{\psi}_7^{(k+1)}), \{(U_7^{(k+1)}, \tilde{\psi}_7^{(k+1,a)}) : a \in F^\times\} \rangle.$$

Here $U_{ij} = \{u \in U_{\max} : u_{k,\ell} = 0, \forall (k,\ell) \neq (i,j)\}$.

In Lemma 9.3.3 below we prove that for k even and $a \in F^\times$,

$$(N_{n+\frac{k}{2}}, \psi_{n+\frac{k}{2},a})|(U_7^{(k+1)}, \tilde{\psi}_7^{(k+1,a)}),$$

where

$$\psi_{\ell,a}(u) = \psi(u_{1,2} + \cdots + u_{\ell-1,\ell} + au_{\ell,2n} + u_{\ell,2n+1}).$$

The present lemma then follows from the following observations:

- $(U_7^{(1)}, \tilde{\psi}_7^{(1)}) = (U_2, \psi_2)$, (with ψ_2 defined as at the beginning of this section).
- $(U_7^{(2n-1)}, \tilde{\psi}_7^{(2n-1)}) = (U_3, \psi_3)$
- If one applies \dagger to both sides of a relation among unipotent periods, it remains valid.
- The character $\psi_{n+\frac{k}{2},a}$ of $N_{n+\frac{k}{2}}$ is in general position. (Cf. remarks 6.1.2)
- The set $\{(N_\ell, \vartheta) : n \leq \ell < 2n \text{ and } \vartheta \text{ in general position.}\}$ is stable under \dagger .
- The number of times we conjugate by the preimage of an element of determinant minus 1 in passing from $(U_7^{(k)}, \tilde{\psi}_7^{(k)})$ back to $(U_7^{(k)}, \psi_7^{(k)})$ is precisely n .

□

Lemma 9.3.3. *Let $(N_{n+\frac{k}{2}}, \psi_{n+\frac{k}{2},a})$ and $(U_7^{(k+1)}, \tilde{\psi}_7^{(k+1,a)})$ be defined as in the previous lemma. Then*

$$(N_{n+\frac{k}{2}}, \psi_{n+\frac{k}{2},a})|(U_7^{(k+1)}, \tilde{\psi}_7^{(k+1,a)}).$$

Proof. We regard a as fixed for the duration of this argument, and omit it from the notation. We need still more unipotent periods. Specifically, for each k, ℓ define $U_9^{(k,\ell)}$ to be the subgroup of U_{\max} defined by requiring that $u_{ij} = 0$ under any of the following conditions:

$$\begin{aligned} k < i \leq k + 2\ell, \quad i \equiv k + 1 \pmod{2} \text{ and } j = i + 1 \\ i > k + 2\ell \\ i = k + 2\ell - 1, \text{ and } j \neq 4n + 1 - k - 2\ell, \\ i = k + 2\ell \text{ and } j < 2n. \end{aligned}$$

The formula

$$\psi(u_{1,2} + \cdots + u_{k-1,k} + u_{k,k+2} + u_{k+1,k+3} + \cdots + u_{k+2\ell-2,k_{k+2\ell}} + au_{k+2\ell,2n} + u_{k+2\ell,2n+1})$$

defines a character of this group which we denote $\psi_9^{(k,\ell)}(u)$. Also, let $U_{10}^{(k,\ell)}$ denote the subgroup of U_{\max} defined by requiring that $u_{ij} = 0$ under any of the following conditions:

$$\begin{aligned} k < i \leq k + 2\ell, \quad i \equiv k + 1 \pmod{2} \text{ and } j = i + 1 \\ i > k + 2\ell - 1 \\ i = k + 2\ell - 1 \text{ and } j > 2n, 2n + 1. \end{aligned}$$

The formula

$$\psi(u_{1,2} + \cdots + u_{k,k+1} + u_{k+1,k+3} + \cdots + u_{k+2\ell-2,k+2\ell} + au_{k+2\ell-1,2n} + u_{k+2\ell-1,2n+1})$$

defines a character of this group which we denote $\psi_{10}^{(k,\ell)}(u)$. The period $(U_9, \psi_9^{(k,\ell)})$ is conjugate to $(U_{10}, \psi_{10}^{(k,\ell)})$.

Let $U_{11}^{(k,\ell)}$ denote the subgroup of U_{\max} defined by requiring that $u_{ij} = 0$ under any of the following conditions:

$$\begin{aligned} k < i \leq k + 2\ell, \quad i \equiv k \pmod{2} \text{ and } j = i + 1 \\ i > k + 2\ell - 1 \\ i = k + 2\ell - 1 \text{ and } j > 2n, 2n + 1. \end{aligned}$$

Then $(U_{10}, \psi_{10}^{(k,\ell)})$ may be swapped for $(U_{11}, \psi_{11}^{(k,\ell)})$, where $\psi_{11}^{(k,\ell)}$ is defined by the same formula as $\psi_{10}^{(k,\ell)}$.

Also, $(U_{11}, \psi_{11}^{(k,\ell)})$, is clearly divisible by $(U_9, \psi_9^{(k+1,\ell-1)})$: to pass from the former to the latter one simply drops the integration over $u_{k+2\ell-2,j}$, for $j \neq 4n - k - 2\ell + 2$.

To complete the argument: for k even the period $(U_9^{(k+1,n-\frac{k}{2}-1)}, \psi_9^{(k+1,n-\frac{k}{2}-1)})$ divides the period $(U_7^{(k+1)}, \tilde{\psi}_7^{(k+1,a)})$. Indeed the only difference between the two is that in the former, we omit integration over $u_{2n-2,2n}$.

It follows that $(U_9^{(k+1,n-\frac{k}{2}-1)}, \psi_9^{(k+1,n-\frac{k}{2}-1)})$ is divisible by $(U_{10}^{n+\frac{k}{2}-1,1}, \psi_{10}^{n+\frac{k}{2}-1,1})$. Finally, every extension of $\psi_{10}^{n+\frac{k}{2}-1,1}$ to a character of $N_{n+\frac{k}{2}}$ is in the same orbit as $\psi_{n+\frac{k}{2},a}$. (See Remarks 6.1.2.) Hence

$$(U_{10}^{n+\frac{k}{2}-1,1}, \psi_{10}^{n+\frac{k}{2}-1,1}) \sim (N_{n+\frac{k}{2}}, \psi_{n+\frac{k}{2},a}).$$

The result follows. \square

Lemma 9.3.4. *As in Theorem 6.1.4, let V_i denote the unipotent radical of the standard parabolic of G_{4n} having Levi isomorphic to $GL_i \times G_{4n-2i}$ (for $1 \leq i \leq 2n - 2$). Let $V_i^{4n-2m-1}$ denote the unipotent radical of the standard maximal parabolic of G_{2n+1} (embedded into G_{4n} as $L_{n-1}^{\psi_{n-1}}$) having Levi isomorphic to $GL_i \times G_{2n-2i+1}$ (for $1 \leq i \leq n$). Let (N_ℓ, ψ_ℓ) be the period used to define the descent, as usual, and let $(N_\ell, \psi_\ell)^{(4n-2k)}$ denote the analogue for G_{4n-2k} , embedded into G_{4n} inside the Levi of a maximal parabolic.*

Then, $(V_k^{2n+1}, \mathbf{1}) \circ (N_{n-1}, \psi_{n-1})$ is an element of

$$\langle (N_{n+k-1}, \psi_{n+k-1}), \{(N_{n+j-1}, \psi_{n+j-1})^{(4n-2k+2j)} \circ (V_{k-j}, \mathbf{1}) : 1 \leq j < k\} \rangle.$$

Proof. In this proof, we shall not need to refer to any of the unipotent periods defined previously. On the other hand we will need to consider several new unipotent periods. For convenience, we start the numbering over from one.

Thus, let $(U_1, \psi_1) = (V_k^{2n+1}, \mathbf{1}) \circ (N_{n-1}, \psi_{n-1})$. To describe this group and character in detail, U_1 is the subgroup defined by $u_{ij} = 0$ if $n - 1 < i \leq n - 1 + k < j$, or $n - 1 + k < i$ and $u_{i,2n} = u_{i,2n+1}$ if $n - 1 < i \leq n - 1 + k$, and ψ_1 is given by

$$\psi_1(u) = \psi_0(u_{1,2} + \cdots + u_{n-2,n-1} + u_{n-1,2n} - u_{n-1,2n+1}).$$

Next, let U_2 denote the subgroup of U_1 defined by the additional conditions $u_{ij} = 0$ for $1 \leq i \leq n - 1 < j \leq n - 1 + k$. Let ψ_2 denote the restriction of ψ_1 to this subgroup.

Next, let U_3 denote the subgroup defined by $u_{ij} = 0$ for $i \leq k, j \leq n - 1 + k$, and $i > n - 1 + k$, and $u_{i,2n} = u_{i,2n+1}$ for $i \leq k$. Let

$$\psi_3(u) = \psi(u_{k+1,k+2} + \cdots + u_{k+n-2,k+n-1} + u_{k+n-1,2n} - u_{k+n-1,2n+1}).$$

Then (U_2, ψ_2) is conjugate to (U_3, ψ_3) , by any element of $G_{4n}(F)$ which projects to

$$\begin{pmatrix} & & & I_k \\ & & & \\ I_{n-1} & & & \\ & & I_{4n-2m-2k} & \\ & & & & I_{n-1} \\ & & & & & I_k \end{pmatrix}$$

(cf. subsection 9.2).

Finally, let $U_4 \supset U_3$ denote the subgroup of U_{\max} given by $u_{ij} = 0$ if $j \leq k+1$, or $i \geq n+k$. Then take ψ_4 defined by the same formula as ψ_3

Certainly $(U_2, \psi_2)|(U_1, \psi_1)$, and $(U_2, \psi_2) \sim (U_3, \psi_3)$. In Lemma 9.3.5 we prove that $(U_3, \psi_3) \sim (U_4, \psi_4)$. It follows that $(U_4, \psi_4)|(U_1, \psi_1)$. In fact, one may prove by an argument similar to the proof of Lemma 9.3.5 that in fact $(U_2, \psi_2) \sim (U_1, \psi_1)$ and hence $(U_4, \psi_4) \sim (U_1, \psi_1)$. But this is not needed for our purposes.

Next, let $U^{(r)}$ denote the subgroup of U_{\max} defined by $u_{ij} = 0$ for $j \leq r$, or $i \geq n+k$. So, $U_4 = U^{(k+1)}$, and $N_{n+k-1} = U^{(1)}$.

Let $\psi^{(r)}$ denote the character of $U^{(r)}$ defined by

$$\psi^{(r)}(u) = \psi_0 \left(\sum_{i=r}^{n-2+k} u_{i,i+1} + u_{n-1+k,2n} + u_{n-1+k,2n+1} \right).$$

Then $(U_4, \psi_4) = (U^{(k+1)}, \psi^{(k+1)})$, and $(N_{n+k-1}, \psi_{n+k-1}) = (U^{(1)}, \psi^{(1)})$. It is an easy consequence of Lemma 9.1.1 that

$$(U^{(r)}, \psi^{(r)}) \in \langle (U^{(r-1)}, \psi^{(r-1)}), (N_{n+k-r}, \psi_{n+k-r})^{(4n-2r+2)} \circ (V_{r-1}, \mathbf{1}) \rangle.$$

The result follows. \square

Lemma 9.3.5. *Let (U_3, ψ_3) and (U_4, ψ_4) be defined as in the previous lemma. Then $(U_4, \psi_4) \sim (U_3, \psi_3)$.*

Proof. It's clear that $(U_3, \psi_3)|(U_4, \psi_4)$, so we only need to prove that $(U_4, \psi_4)|(U_3, \psi_3)$. The proof involves a family of groups defining intermediate stages. For ℓ such that $1 \leq \ell \leq n-1$ we define $U_4^{(\ell)}$ to be the subgroup of U_4 defined by the condition that for $i \leq k$ the coordinate u_{ij} must be zero for $j \leq k+\ell$. Thus $U_4 = U_4^{(1)} \supset U_4^{(2)} \supset \dots \supset U_4^{(n-1)} \supset U_3$. For each of these groups we consider the period defined using the restriction of ψ_4 .

We must show that $(U_4^{(n-1)}, \psi_4)|(U_3, \psi_3)$ and that $(U_4^{(i)}, \psi_4)|(U_4^{(i-1)}, \psi_4)$. In each case, all that is involved is an invocation of Lemma 9.1.1. For the first application, what must be checked is that the normalizer of $U_4(F)$ in $G(F)$ permutes $\{\psi'_4 : \psi'_4|_{U_3} = \psi_3\}$ transitively. Let $y(\underline{r}) = y(r_1, \dots, r_k)$ denote the unipotent element in $G_{4n}(F)$ which projects to $I + r_1 e'_{1,2n} + \dots + r_k e'_{k,2n}$. Then every element of $U_4^{(m)}$ is uniquely expressible as $u_3 y(\underline{r})$, for $u_3 \in U_3$ and $\underline{r} \in \mathbb{G}_a^k$. Hence a map ψ'_4 as above is determined by its composition with y , which defines a character of $(F \setminus \mathbb{A})^k$, and hence is of the form

$$(r_1, \dots, r_k) \mapsto \psi(a_1 r_1 + \dots + a_k r_k)$$

for some $a_1, \dots, a_k \in F$. Consider the unipotent element $z(a_1, \dots, a_k)$ of G_{4n} which projects to $I + a_1 e'_{k+n-1,1} + \dots + a_k e'_{k+n-1,k}$. We claim first that it normalizes $U_4^{(n-1)}$, and second that $\psi_4(z(a)y(r)z(a)^{-1}) = \psi(a_1 r_1 + \dots + a_k r_k)$. As noted in 9.2 this may be checked by a matrix multiplication in SO_{4n} .

The proof that $(U_4^{(i)}, \psi_4)|(U_4^{(i-1)}, \psi_4)$ is entirely similar, with the role of $y(r)$ played by $y^{(i)}(r)$ which projects to $I + r_1 e'_{1,k+i+1} + \cdots + r_k e'_{k,k+i+1}$ and that of $z(\underline{a})$ played by $z^{(i)}(\underline{a})$ which projects to $I + a_1 e'_{k+i,1} + \cdots + a_k e'_{k+i,k}$. \square

REFERENCES

- [Asg] M. Asgari, Local L -functions for Split Spinor Groups, *Canad. J. Math.* Vol 54 (4), 2002 pp. 673-693.
- [Asg-Sha1] M. Asgari and F. Shahidi, *Generic Transfer For general Spin Groups*, *Duke Math. Journal.* Vol. 132, No. 1, 2006, pp. 137- 190.
- [Asg-Sha2] M. Asgari and F. Shahidi, *Generic Transfer From $GSp(4)$ to $GL(4)$* , *Compos. Math.* 142 (2006), no. 3, 541–550.
- [Banks1] W. Banks, Twisted symmetric-square L -functions and the nonexistence of Siegel zeros on $GL(3)$.
- [Banks2] W. Banks, Exceptional Representations on the Metaplectic Group. Dissertation, Stanford University 1994
- [Banks3] W. Banks, private communication.
- [BG] D. Bump, D. Ginzburg Symmetric square L -functions on $GL(r)$. *Ann. of Math. (2)* 136 (1992), no. 1, 137–205.
- [BZ1] I. N. Bernšteĭn and A. V. Zelevinskiĭ. Representations of the group $GL(n, F)$, where F is a local non-Archimedean field. *Uspehi Mat. Nauk*, 31(3(189)):5–70, 1976.
- [BZ2] I. N. Bernšteĭn and A. V. Zelevinskiĭ, Induced representations of reductive p -adic groups. I. *Ann. Sci. École Norm. Sup. (4)* 10 (1977), no. 4, 441–472.
- [Car] P. Cartier, Representations of p -adic groups: a survey. *Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977)*, Part 1, pp. 111–155, *Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979*. Available online at http://www.ams.org/online_bks/pspum331/
- [Cass] W. Casselman, Introduction to the theory of admissible representations of p -adic reductive groups, Available at <http://www.math.ubc.ca/cass/research.html>
- [C-S] W. Casselman, J. Shalika, The unramified principal series of p -adic groups. II. The Whittaker function. *Compositio Math.* 41 (1980), no. 2, 207–231.
- [C-K-PS-S1] J. Cogdell, H. Kim, I. Piatetski-Shapiro, F. Shahidi, *On lifting from classical groups to GL_N* , *Publ. Math. Inst. Hautes Études Sci.*, 93 (2001), pp.5-30.
- [C-K-PS-S2] J. Cogdell, H. Kim, I.I. Piatetski-Shapiro, F. Shahidi, Functoriality for the classical groups. *Publ. Math. Inst. Hautes Études Sci.* No. 99 (2004), 163–233.
- [Gan-Tak] W.T. Gan, S. Takeda, The Local Langlands Conjecture for $GSp(4)$, preprint, available at <http://arxiv.org/abs/0706.0952>
- [Gel-Sha] S. Gelbart, F. Shahidi, *Analytic properties of automorphic L -functions*. *Perspectives in Mathematics*, 6. Academic Press, Inc., Boston, MA, 1988.
- [GK75] I. M. Gelfand and D. A. Kajdan. Representations of the group $GL(n, K)$ where K is a local field. In *Lie groups and their representations (Proc. Summer School, Bolyai János Math. Soc., Budapest, 1971)*, pages 95–118. Halsted, New York, 1975.
- [Gi-PS-R] D. Ginzburg, I.I. Piatetski-Shapiro, and S. Rallis, L functions for the orthogonal group. *Mem. Amer. Math. Soc.* 128 (1997), no. 611, MR1357823 (98m:11041)
- [GRS1] D. Ginzburg, S. Rallis, and D. Soudry, *On a correspondence between cuspidal representations of GL_{2n} and \widetilde{Sp}_{2n}* , *J. Amer. Math. Soc.*,12(1999), n.3, pp. 849-907
- [GRS2] D. Ginzburg, S. Rallis, and D. Soudry, *Lifting cusp forms on GL_{2n} to \widetilde{Sp}_{2n} : the unramified correspondence*, *Duke Math. J.*, 100 (1999),n. 2, pp. 243-266
- [GRS3] D. Ginzburg, S. Rallis and D. Soudry, *On explicit lifts from cusp forms from GL_m to classical groups*, *Annals of Math.*, 150(1999) 807-866. Available online at <http://arxiv.org/pdf/math.NT/9911264>
- [GRS4] D. Ginzburg, S. Rallis, D. Soudry, *Generic automorphic forms on $SO(2n + 1)$: functorial lift to $GL(2n)$, endoscopy, and base change*, *Internat. Math. Res. Notices*, 14 (2001), pp.729-764.
- [GRS5] D. Ginzburg, S. Rallis, and D. Soudry, *Endoscopic representations of \widetilde{Sp}_{2n}* , *J. Inst. Math. Jussieu*, vol. 1 2002, no. 1, pp. 77-123.
- [GRS6] D. Ginzburg, S. Rallis, D. Soudry, On the automorphic theta representation for simply laced groups. *Israel Journal of Mathematics* 100 (1997), 61-116.
- [H-T] M. Harris, and R. Taylor, *The geometry and cohomology of some simple Shimura varieties*. With an appendix by Vladimir G. Berkovich. *Annals of Mathematics Studies*, 151. Princeton University Press, Princeton, NJ, 2001.

- [Henn1] G. Henniart, Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p -adique. *Invent. Math.* 139 (2000), no. 2, 439–455.
- [Henn2] G. Henniart, Correspondance de Langlands et fonctions L des carrés extérieur et symétrique, IHES preprint M03-20, available at <http://www.ihes.fr/PREPRINTS/M03/M03-20.pdf>
- [I] J. Igusa, A Classification of Spinors Up to Dimension Twelve, *Amer. J. Math.*, Vol. 92, No. 4 (1970), 997-1028.
- [Ja] H. Jacquet, Generic representations, in *Non-commutative harmonic analysis (Actes Colloq., Marseille-Luminy, 1976)*, pp. 91–101. *Lecture Notes in Math.*, Vol. 587, Springer, Berlin, 1977.
- [Ja-Sh1] H. Jacquet, J. Shalika, Exterior square L functions, *Automorphic Forms, Shimura Varieties and L -Functions*, Vol.2 Academic Press (1990), 143-226.
- [Ja-Sh2] H. Jacquet, J. Shalika, On Euler products and the classification of automorphic representations. I. *Amer. J. Math.* 103 (1981), no. 3, 499–558.
- [Ja-Sh3] H. Jacquet, J. Shalika, On Euler products and the classification of automorphic representations. II. *Amer. J. Math.* 103 (1981), no. 4, 777–815.
- [Ji-So] D. Jiang, D. Soudry, The local converse theorem for $SO(2n + 1)$ and applications. *Ann. of Math. (2)* 157 (2003), no. 3, 743–806.
- [Kim] H. Kim, Langlands-Shahidi Method and Poles of Automorphic L -functions: Application to Exterior Square L -functions, *Canad. J. Math.* Vol. 51 (4), 1999, pp. 835-849.
- [Kim-Sh] H. Kim, F. Shahidi, On simplicity of poles of automorphic L -functions. *J. Ramanujan Math. Soc.* 19 (2004), no. 4, 267–280.
- [Kn] A. Knapp, *Representation theory of semisimple groups. An overview based on examples*. Princeton Mathematical Series, 36. Princeton University Press, Princeton, NJ, 1986.
- [L1] R. Langlands, Euler Products, Yale University Press, (1971). Available online at <http://www.sunsite.ubc.ca/DigitalMathArchive/Langlands/>
- [L2] R. Langlands, “On the classification of irreducible representations of real algebraic groups,” *Representation theory and harmonic analysis on semisimple Lie groups*, pp. 101–170, *Math. Surveys Monogr.*, 31, Amer. Math. Soc., Providence, RI, 1989. Available online at <http://www.sunsite.ubc.ca/DigitalMathArchive/Langlands/>
- [L3] R. P. Langlands, On the notion of an automorphic representation, in *Proc. Sympos. Pure Math., XXXIII, Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1*, pp. 189–207, Amer. Math. Soc., Providence, R.I., 1979. Available online at <http://www.ams.org/online.bks/pspum331/> and <http://www.sunsite.ubc.ca/DigitalMathArchive/Langlands/>
- [L4] R. P. Langlands, Automorphic representations, Shimura varieties, and motives. Ein Mrchen, in *Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2*, pp. 205–246, *Proc. Sympos. Pure Math.*, XXXIII, Amer. Math. Soc., Providence, R.I., 1979. Available online at <http://www.ams.org/online.bks/pspum332/> and <http://www.sunsite.ubc.ca/DigitalMathArchive/Langlands/>
- [MW1] C. Mœglin and J.-L. Waldspurger *Spectral Decomposition and Eisenstein Series*, Cambridge University Press, 1995.
- [MW2] C. Mœglin, J.-L. Waldspurger, Le spectre résiduel de $GL(n)$, *Ann. Sci. École Norm. Sup.* 22 (1989), 605–674. MR 91b:22028
- [PraRam] D. Prasad and D. Ramakrishnan, preprint, 2006. Available online at <http://www.math.caltech.edu/people/dinakar.html>.
- [Sha1] F. Shahidi, Langlands-Shahidi method. *Automorphic forms and applications*, P. Sarnak, F. Shahidi ed. IAS/Park City Math. Ser., 12, Amer. Math. Soc., Providence, RI, 2007, pp.299–330
- [Sha2] F. Shahidi, A proof of Langlands’ conjecture on Plancherel measures; complementary series for p -adic groups. *Ann. of Math. (2)* 132 (1990), no. 2, 273-330.
- [Sha3] F. Shahidi, Local coefficients as Artin factors for real groups. *Duke Math. J.* 52 (1985), no. 4, 973–1007.
- [Sha4] F. Shahidi, On certain L -functions. *Amer. J. Math.* 103 (1981), no. 2, 297–355.
- [Sha5] F. Shahidi, Whittaker models for real groups, *Duke Math. J.* 47 (1980), no. 1, 99–125.
- [Sil] A. Silberger, *Introduction to harmonic analysis on reductive p -adic groups. Based on lectures by Harish-Chandra at the Institute for Advanced Study, 1971–1973*. *Mathematical Notes*, 23. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1979
- [Spr] T.A. Springer, *Linear Algebraic Groups*, 2nd Ed Birkhäuser, 1998.
- [So1] D. Soudry, On Langlands functoriality from classical groups to GL_n . *Automorphic forms. I. Astérisque* No. 298 (2005), 335–390. MR2141707 (2006e:11070)

- [So2] D. Soudry, Rankin-Selberg convolutions for $SO_{2l+1} \times GL_n$: local theory. Mem. Amer. Math. Soc. 105 (1993), no. 500
- [Tad1] M. Tadić, Classification of unitary representations in irreducible representations of general linear group (non-Archimedean case). Ann. Sci. cole Norm. Sup. (4) 19 (1986), no. 3, 335–382.
- [Tad2] M. Tadić, An external approach to unitary representations. Bull. Amer. Math. Soc. (N.S.) 28 (1993), no. 2, 215–252.
- [Tate1] J. Tate, Number theoretic background. Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, pp. 3–26, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979. Available online at <http://www.ams.org/online.bks/pspum332/>
- [Tate2] J. Tate, Fourier analysis in number fields, and Hecke’s zeta-functions. 1967 *Algebraic Number Theory (Proc. Instructional Conf., Brighton, 1965)* Cassels, Frohlich ed., Thompson, Washington, D.C., pp. 305–347.
- [Tits] J. Tits, Reductive groups over local fields. Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1, pp. 29–69, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979. Available online at <http://www.ams.org/online.bks/pspum332/>
- [Vog] D. Vogan, Gelfand-Kirillov dimension for Harish-Chandra modules. Invent. Math. 48 (1978), no. 1, 75–98.
- [Z] A. V. Zelevinskiĭ, Induced representations of reductive \mathfrak{p} -adic groups. II. On irreducible representations of $GL(n)$. Ann. Sci. École Norm. Sup. (4) 13 (1980), no. 2, 165–210.

E-mail address: jhundley@math.siu.edu

MATH. DEPARTMENT, MAILCODE 4408, SOUTHERN ILLINOIS UNIVERSITY CARBONDALE, 1245 LINCOLN DRIVE CARBONDALE, IL 62901

E-mail address: eitan.sayag@gmail.com

INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL