

Callus Induction and Plant Regeneration of *Vigna mungo* (L.) Hepper via Half Seed Explant

*Harisaranraj, R., S. Saravana Babu and Suresh, K.

Department of Plant Biology and Plant Biotechnology
Chikkaiah Naicker College, Erode. (T.N.) INDIA
*hari83biotech@yahoo.com

Issued 11 August 2008

Abstract

The present study optimized the regeneration protocol by using half seed explant in *Vigna mungo* (L.) Hepper organogenesis. Half seed explants were inoculated onto B5 medium supplemented with kinetin (4.7 μ M to 23.5 μ M), 6- benzyladenine (4.4 μ M to 22.2 μ M), naphthaleneacetic acid (5.4 μ M to 27.0 μ M), indolebutyric acid (4.9 μ M to 24.5 μ M) and 2,4- dichlorophenoxyacetic acid (4.5 μ M to 22.5 μ M). Callus initiation was observed in all media evaluated and the highest cell proliferation was obtained from explants cultivated in the presence of 13.3 μ M BAP and 13.5 μ M 2,4-D. Shoot induction was obtained from callus induced on 13.3 μ M BAP and 13.5 μ M 2,4-D at 6 weeks after transferring the callus to a B5 medium supplemented with 13.3 μ M BAP. Roots were induced from shoots on B5 media with indolebutyric acid (IBA, 14.7 μ M) and then regenerated plants were hardened and acclimated in greenhouse conditions.

Key Words: *Vigna mungo*, callus induction, half seed, rooting, shoot regeneration.

Introduction

Vigna mungo (L.) Hepper (Black gram) is considered to have been domesticated in India from its wild ancestral form *V.mungo* var.*silvestris* (Lukoki, Marechal & Otoul, 1980). Center of genetic diversity is found in India (Zeven and de Wet. 1982). Natural distribution of *V.mungo* var.*silvestris* ranges from India to Myanmar (Tateishi. 1996). *Vigna mungo* (L.) Hepper (Black Gram or Urd Bean) is one of the most widely used pulse crop in India. It is a highly prized pulse, very rich in phosphoric acid. It is cultivated as fallow crop after rice cultivation in India. It is grown in various agro-ecological conditions and cropping systems with diverse agricultural practices (Sanjeev Gupta *et al.*). The improvement and optimization of *Vigna mungo* characteristics, such as increasing resistance to pests, lowering allergenic protein levels in seeds, drought and salt tolerance is therefore desirable.

The crop improvement was done by breeding methods in early days. However, breeding is difficult due to the fact that *Vigna mungo* is self-pollinating crop and the genetic variation among the Black gram varieties is narrow. The regeneration system used to generate genetically modified plants was somatic embryogenesis from immature seeds (Christou *et al.*, 1989) or organogenesis from cotyledonary nodes (Finer and McMullen, 1991; Shan *et al.*, 2005). In *Vigna mungo*, genetic modification is also based on organogenesis from the cotyledonary nodes, a high efficient regeneration system was obtained by sub culturing nodes on Thidiazuron (TDZ) supplemented medium (Tzitzikas *et al.*, 2004). Medium components with quantity of plant growth regulator influence the regeneration of plants (Shan *et al.*, 2005). Callus regeneration is advantageous over direct regeneration for transformation since effective selection of transgenic cells can be achieved. However, the efforts made to regenerate plants from callus have yielded poor results since plants could not be regenerated from any type of *Vigna mungo* callus (Hu and Wang, 1999). In attaining this goal, we owe much to transformation techniques for producing new breeding materials that would not be available in the germplasm among cross-compatible species. A number of successful regeneration protocols have been developed, especially with a view to

facilitate genetic transformation.

The earlier reports focused on the cotyledonary node since the morphogenetic potentiality is confined to that region (Hu and Wang, 1999). Eventhough plants have been successfully transformed by using cotyledonary nodal explants, the major problem of chimerism still persists (Meurer *et al.*, 1998). This is primarily due to pre-existing meristematic shoot buds, which continue to grow effectively on a medium containing a selectable agent like antibiotic or herbicide. Christou *et al.* (1988) could effectively select transgenic calli after bombarding protoplasts but they failed to produce transgenic plants. The callus induction and plant regeneration of Indian soybean (*Glycine max* (L.) Merr. cv. CO3) via half seed explant culture was carried out by (B. D. Ranjithakumari *et al.*). In this present investigation was to standardize the optimum protocol for callus and shoot regeneration from half seed explant of the *Vigna mungo*.

Materials and methods

Matured *Vigna mungo* seeds were utilized in the efficient production of *in vitro Vigna mungo* plant production experiments. *Vigna mungo* seeds were washed under continuous flashing of running tap water for 30 min and then treated with a solution of the Tween 20 (5% v/v) for 10 min and finally surface sterilized with HgCl₂ (0.1% w/v) for 10 min. Lastly, the seeds were washed three times with autoclaved distilled water to remove any trace of

Half seed explants preparation

Disinfected seeds were soaked in sterile distilled water for about 4 h and a longitudinal cut along the hilum was made to separate the cotyledons, and the seed coat was removed. The embryonic axis found at the junctions of the hypocotyl and cotyledon was excised to obtain the half- seed explants.

Callus induction

Initial explants consisted of half seed (ie, the part of the explant from where the embryonic axis was removed) was cultured in flate side up in the medium with the base of the explant was embedded condition. The callus initiation medium contained B5 salts and vitamins (Gamborg *et al.*, 1968), 30 g l-1 sucrose. The pH was adjusted to 5.8 and solidified with 0.7% agar before autoclaving at 121°C for 20 min. The effect of growth regulators was tested using KIN (kinetin), BAP (6-benzyladenine), NAA (naphthaleneacetic acid), IBA (indolebutyric acid), 2,4-D (2,4-dichlorophenoxyacetic acid) as follows: KIN (4.7 µ M to 23.5 µ M), BAP (4.4 µ M to 22.2 µ M) and NAA (5.4 µ M to 27.0 µ M) or IBA (4.9 µ M to 24.5 µ M) or 2,4-D (4.5 µ M to 22.5) in combination with BAP (4.4 µ M to 22.2 µ M). Medium without plant growth regulators was used as a control. Cultures were maintained under light 16h or dark 8 h conditions at 26±2°C. 50 half seed explants were inoculated per treatment and repeated 3 times and the frequency of callus formation was determined 4 weeks after culture initiation.

Shoot induction and multiplication

To induce and proliferate shoots, the calli were transferred to shoot induction medium containing B5 salts, B5 vitamins, 30 g l-1 sucrose, pH 5.8, 0.7% agar and fortified with BAP (4.4 µ M to 22.2 µ M). Calli were kept under the same light/ darkness conditions from the callus initiation for 4 weeks. Depending on the treatment, the number of shoots per treatment was recorded at 6 and 10 weeks after transferring the calli to shoot induction medium.

Rooting and Hardening of plantlets

For root induction, 14.7 µ M IBA was tested. Elongated shoots (1.5 to 2.5 cm long) were isolated and transferred to B5 medium, supplemented with B5 vitamins, 30 g l-1 sucrose, pH 5.8, 0.7% agar and 14.7 µ M IBA. After 3 weeks, rooted plantlets were rinsed with water to wash off the agar medium and then, transplanted to soil containing cups for hardening. The hardened plants were maintained at 24±2°C with 18 h photoperiod (140 µmoles s-1) for 2 weeks then, transferred to the greenhouse.

Statistical analysis

The various concentrations of plant growth regulators (KIN, BAP, BAP+ NAA, BAP+ IBA and BAP+ 2, 4 – D) were tested in callus induction and plant regenerations. Frequency of callus and shoot initiation, number of shoots were analyzed by SPSS software, in which statistical significance was determined at the 0.05 probability level.

Results and discussion

Half seed explant is an efficient source for the *Vigna mungo* callus initiation and shoot regeneration. The half seed derived cotyledonary nodal callus could serve as an ideal starting material for developing an efficient *Vigna mungo* transformation system. (Sairam *et al.*, 2003).

Callus initiation

The regeneration of callus induction and plant regeneration were influenced by different kinds of plant growth regulator (KIN, BAP, NAA, IBA and 2,4–D). Callus was induced in all media tested and significant differences were observed in the induction frequency between different plant growth regulators (Fig. 1 to 5). Among the various plant growth regulators, the highest frequency of callus induction was observed to be when the half seed explants were cultured in the presence of BAP (13.3 μ M) + 2,4–D (13.5 μ M) (Fig. 5). These observations indicate that the morphogenetic potential is confined to only the cotyledonary nodal cells as reported by Hu and Wang (1999).




Fig. 1. The effect of different concentrations of KIN on callus regenerations.




Fig. 2. The effect of different concentrations of BAP on callus regenerations.

The low frequency of callus induction was observed in KIN (14.1 μ M) exposed explants cultured on the B5 medium. The explants grown in B5 medium supplemented with BAP (4.4 μ M to 22.2 μ M) and different auxin combination such as NAA (5.4 μ M to 27.0 μM), IBA (4.9 μ M to 24.5 μ M) and 2,4-D (4.5 μ M to 22.5) were given the better yield than that of KIN (4.7 μM to 23.5 μ M). Barwale *et al.* (1986a) reported that the organogenic callus cultures were obtained from immature soybean embryos grown on the medium with a high 6-benzylaminopurine (BAP) concentration (13.3 μ M) and 0.2 μ M NAA.




Fig. 3. The effect of different concentrations of BAP + NAA on callus regenerations.

Plantlet regeneration

Half seed explant derived calli were cultured on the medium fortified with various concentrations of BAP supplemented B5 medium (SI medium). In *Vigna mungo*, BAP was found to be the most efficient in shoot formation when excised parts of mature and immature cotyledones were used (Sairam *et al.*, 2003). 13.3 μ M BAP + 13.5 μ M 2, 4 -D treated callus produced the maximum number of shoots than other concentrations of BAP alone (Table 1). The marked difference observed between the average number of shoots obtained from callus induced from different concentrations and combinations of BAP with 2, 4 - D could be related to the media used for callus induction and multiplication. Callus induced on 13.3 μ M BAP + 13.5 μ M 2, 4 -D formed the highest number of shoots after 6 weeks in the presence of BAP treated SI medium (Table 1). The former calluses were completely covered with shoot

buds were easily disintegrated during manipulation. However, the shoots could be excised from their surface. The calli could be kept for six months of subcultures producing an average of 6 shoots per callus in every month. The procedures that have been established for shoot induction and multiplication of *Glycine* species have reported an average from 2 to 5.5 shoots, depending on the explant used (Mederos *et al.*, 1997; Moura, 1998). These calli could be maintained in a long term process of shoot induction. The maximum number of shoots was regenerated to 13.3 μ M BAP supplemented with SI medium. It is in contrast with BAP, alone or in combination with NAA, has previously been reported as being efficient in promoting shoot differentiation in several species (Blakesley and Constantine, 1992).

Table 1

B5 medium+(BAP and 2,4 - D) for callus induction	B5 medium+(BAP) for shoot regeneration	No. of shoots per callus culture
BAP		
4.4 μ M	4.4 μ M	21.5 \pm 5.2
8.8 μ M	8.8 μ M	22.6 \pm 2.3
13.3 μ M	13.3 μ M	25.4 \pm 1.8
17.7 μ M	17.7 μ M	20.9 \pm 2.2
22.2 μ M	22.2 μ M	20.3 \pm 6.1
BAP+2,4 - D		
4.4 μ M + 4.5 μ M	4.4 μ M	22.8 \pm 2.2
8.8 μ M +9.0 μ M	8.8 μ M	26.1 \pm 4.2
13.3 μ M + 13.5 μ M	13.3 μ M	32.4 \pm 2.9
17.7 μ M +18.0 μ M	17.7 μ M	28.8 \pm 6.2
22.2 μ M +22.5 μ M	22.2 μ M	30.8 \pm 1.8

Use of 5.0 μ M BAP significantly increased the regeneration frequency of shoots from the *Vigna mungo* cotyledonary node. Wright *et al.* (1986) and Hinchee *et al.* (1988) also obtained shoot regeneration on cotyledonary explants at 5.0 μ M BAP. Use of 7.5 μ M BAP [as routinely used in cotyledonary node protocol; Olhoff *et al.*, (2003); Paz *et al.* (2004)] gave the lowest regeneration rate when applied to the half- seed system. However, for *H. canariense* and *H. foliosum*, the best results for shoot induction were obtained directly from apical or axillary buds cultured in media supplemented with BAP and NAA (Mederos, 1991; Moura, 1998).




Fig. 4. The effect of different concentrations of BAP + IBA on callus regenerations.

Barwale *et al.* (1986b) reported that, the best multiple shoot induction was observed in B5 medium supplemented with 1.0 or 5.0 μ M BAP in several *Vigna mungo* genotypes. Paz *et al.* (2006) suggested that culture condition optimized in the cotyledonary node system were not immediately applicable to the half-seed protocol and by improving an optimum hormonal environment to the explant *in vitro*, the ability of transformed cells to regenerate into plants may be improved. In *Vigna mungo*, elongated shoots obtained from callus were rooted in B5 medium with 14.7 μ M IBA. The IBA was an efficient auxin to produce the shoots (Paz *et al.*, 2006).




Fig. 5. The effect of different concentrations of BAP + 2,4 -D on callus regenerations.

Similar results have been described for some species of *Hypericum* (Mederos, 1991; Moura, 1998). The presence of IBA on the rooting media seemed to have an effect on the number of roots induced per shoot. Three weeks were enough to develop strong and healthy plantlets with good root systems, ready to be transplanted and acclimated. The plantlets developed *in vitro* were transferred to sterile soil and maintained in controlled conditions in a growth chamber for one week. In this period of time, the pots were watered regularly. After 2 week, the plants were taken to the greenhouse. In conclusion, using plant growth regulators, the efficient callus mediated regeneration from half seed explant of *Vigna mungo* has been standardized. The half seed callus could serve as an ideal starting material for developing an efficient *Vigna mungo* transformation system.

Acknowledgment

Authors are gratefully thankful to Prof. K. Sekar, Lecturer (SG), A.A Government Arts College, Namakkal, Tamilnadu, India for constant encouragement and timely help during the present work. We are also grateful to the Directors, BIIOGENEIC – Institute of Biotechnology, Namakkal, Tamilnadu, India, for providing laboratory facilities.

References

1. Barwale, U. B., Kerns, H. R. and Widholm, J. M. (1986a). Plant regeneration from callus cultures of several soybean genotypes via embryogenesis and organogenesis. *Planta* **167**: 473 – 481.
2. Barwale, U. B., Meyer, M. M. and Widholm, J. M. (1986b). Screening of *Glycine max* and *Glycine soja* genotypes for multiple shoot formation at the cotyledonary node. *Theoretical and Applied Genetics* **72**: 423-428.
3. Blakesley, D. and Constantine, D. (1992). Uptake and metabolism of 6- benzyladenine in shoot cultures of a

- range of species. *Plant Cell, Tissue and Organ Culture* **28**: 183–186.
4. Christou, P., McCabe, D. E. and Swain, W. F. (1988). Stable transformation of soybean callus by DNA coated gold particles. *Plant Physiology* **87**: 671–674.
 5. Christou, P., Swain, W. F., Yang, N. S. and McCabe, D. E. (1989). Inheritance and expression of foreign genes in transgenic soybean plants. *Proceedings of the National Academy of Sciences of the United States of America* **86**: 7500-7504.
 6. Dan, Y. and Reichert, N. A. (1998). Organogenic regeneration of soybean from hypocotyle explans. *In Vitro Cellular & Developmental Biology-Plant* **34**: 14-21.
 7. Donaldson, P. A. and Simmonds, D. H. (2000). Susceptibility of *Agrobacterium tumefaciens* and cotyledonary node transformation in short-season soybean. *Plant Cell Reports* **19**: 478–484.
 8. Finer, J. F. and McMullen, D. (1991). Transformation of soybean via particle bombardment of embryogenic suspension culture tissue. *In Vitro Cellular & Developmental Biology- Plant* **27**: 175-182.
 9. Finer, J. J., Cheng, T. S. and Verma, D. P. S. (1996). Soybean transformation: technologies and progress. *Soybean: Genetics, molecular biology and biotechnology* **11**: 249–262.
 10. Gamborg, O. L., Miller, R. A. and Ojima, K. (1968). Nutrient requirements of suspensioncultures of soybean root cells. *Experimental Cell Research* **50**: 151-158.
 11. Hinchee, M. A. W., Conor-Ward, D. V., Newell, C. A., McDonnell, R. E., Sato, S. J., Gasser, C. S., Fischhoff, D. A., Re, D. B., Fraley, R. T. and Horsch, R. B. (1988). Production of transgenic soybean plants using *Agrobacterium* mediated DNA transfer. *Biotechnology* **6**: 915–922.
 12. Kaneda, Y., Tabei, Y., Nishimura, S., Harada, K., Akihama, T. and Kitamura, K. (1997). Combination of Thidiazuron and basal media with low salt concentrations increases the frequency of shoot organogenesis in soybean (*Glycine max* (L.) Merrill). *Plant Cell Reports* **17**: 8-12.
 13. Klein, T. M. and Jones, T. J. (1999). Methods of genetic transformation: the gene gun. In: Vasil I. K. (ed) *Molecular Improvement of Cereal Crops* (pp. 21–42). Kluwer Academic Publishers, The Netherlands.
 14. Komatsuda, T. and Ohyama, K. (1988). Genotypes of high competence for somatic embryogenesis and plant regeneration in soybean *Glycine max*. *Plant Cell Reports* **75**: 695–700.
 15. Komatsuda, T., Lee, W. and Oka, S. (1992). Maturation and germination of somatic embryos as affected by sucrose and plant growth regulators in soybeans *Glycine gracilis*. Skvortz and *Glycine max* (L.) Merr. *Plant Cell, Tissue and Organ Culture* **28**: 103–113.
 16. Lukoki, L., R. Marechal and E. Otoul. (1980). Les ancetres sauvages des haricots cultives: *Vigna radiata* (L.) Wilczek et *V. mungo* (L.) Hepper. *Bull. Jard. Bot. Nat. Belgique*. **50**: 385-391.
 17. Mederos, M. S. (1991). *In vitro* growth and multiplication of *Hypericum canariense* L. *Acta Horticultirae* **289**: 133–135.
 18. Mederos, S., San-Andrés, L. and Luis, J. G. (1997). Rosmanol controls explants browning of *Hypericum canariense* L. during *in vitro* establishment of shoots. *Acta Societatis Botanica Poloniae* **66**: 347–349.
 19. Meurer, C. A., Dinkins, R. D. and Collins, G. B. (1998). Factors affecting soybean cotyledonary node transformation. *Plant Cell Reports* **18**: 180–186.

20. Moura, M. (1998). Conservation of *Hypericum foliosum* aiton, an endemic azorean species, by micropropagation. *In Vitro Cellular & Developmental Biology-Plant* **34**: 244–248.
21. Olhoft, P. M., Fligel, L. E., Donovan, C. M., Somers, D. A. (2003). Efficient soybean transformation using hygromycin B selection in the cotyledonary-node method. *Planta*. **216**: 723- 735.
22. Parrott, W. A., Hoffman, L. M., Hildebrand, P., Williams, E. G. and Collins, G. B. (1989). Recovery of primary transformants of soybean. *Plant Cell Reports* **7**: 615–617.
23. Paz, M. M., Martinez, J. C., Kalvig, A. B., Fonger, T. M. and Wang, K. (2006). Improved cotyledonary node method using an alternative explant derived from mature seed for efficient *Agrobacterium* mediated soybean transformation. *Plant Cell Reports* **25**: 206 - 213.
24. Paz, M. M., Shou, H., Guo, Z., Zhang, Z., Banerjee, A. K. and Wang, K. (2004). Assessment of conditions affecting *Agrobacterium* mediated soybean transformation using the cotyledonary node explant. *Euphytica*. **136**: 167 – 179.
25. Radhakrishnan, R. and Ranjithakumari, B.D. (2007). Callus induction and plant regeneration of Indian soybean (*Glycine max* (L.) Merr. cv. CO3) via half seed explant culture. *Journal of Agricultural Technology* **3(2)**: 287-297.
26. Raemakers, K., Emmanouil, N., Tzitzikas, Z. M. A., Richard, G. and Visser, F. (2005). Development of a high efficient, repetitive system of organogenesis in soybean (*Glycine max* (L.) Merr). *Plant Cell Reports* **24**: 507-512.
27. Sairam, R. V., Frankin, G., Hassel, R., Smith, B., Meeker, K., Kashikar, N., Parani, M., Abed, D., Ismail, S., Berry, K. and Goldman, S. L. (2003). A study on effect of genotypes, plant growth regulators and sugars on promoting plant regeneration via organogenesis from soybeancotyledonary nodal callus. *Plant Cell, Tissue and Organ Culture* **75**: 79-85.
28. Saka, H., Voqui-Dinh, T. H. and Cheng, T. (1980). Stimulation of multiple shoot formation on soybean stem nodes in culture. *Plant Sciences Letter* **19**: 193- 201. Shan, Z., Sanjeev Gupta, Gupta, S. R., Dikshit, H. K. and Singh, R. A., *Plant Gene. Resour. Newsl.*, 2001, **127**, 20–24.
29. Tateishi, Y. (1996). Systematics of the species of *Vigna* subgenus *Ceratotropis*. In "Mungbean Germplasm : Collection, Evaluation and Utilization for Breeding Program" JIRCAS Working Report No.2. pp.9-24. Japan International Research Center for Agricultural Science (JIRCAS), Japan.
30. Tzitzikas, E. N., Bergervoet, M., Raemakers, K., Vincken, J. P., Lammeren, A. and Visser, R. G. F. (2004). Regeneration of pea (*Pisum sativum* L.) by a cyclic organogenic system. *Plant Cell Reports* **23**: 453-461.
31. Wright, M. S., Koehler, S. M., Hinchee, M. A. and Carnes, M. G. (1986). Plant regeneration by organogenesis in *Glycine max* (L.) Merr. *Plant Cell Reports* **5**: 150-154.
32. Wright, M. S., Ward, D. V., Hinchee, M. A., Carnes, M. G. and Kaufman, R. J. (1987a). Regeneration of soybean (*Glycine max* (L.) Merr) from cultured leaf tissue. *Plant Cell Reports* **6**: 83-89.
33. Wright, M. S., Williams, M. H., Pierson, P. E. and Carnes, M. G. (1987b). Initiation and propagation of *Glycine max* (L.) Merrill: Plants from tissue cultured epicotyls. *Plant Cell, Tissue and Organ Culture* **8**: 83-90.
34. Zeven, A.C. and J.M.J. de Wet (1982). Dictionary of cultivated plants and their regions of diversity. Centre for Agricultural Publication and Documentation, Wageningen.

