

EFFECTS OF EXPOSURE TO LOW LEVELS OF WATER-BORNE 17 β -ESTRADIOL ON NEST HOLDING ABILITY AND SPERM QUALITY IN FATHEAD MINNOWS

Heiko L. Schoenfuss

University of Minnesota

Dalma Martinovic

University of Minnesota

Peter W. Sorensen

University of Minnesota

ABSTRACT

This study assessed the effects of exposure to low levels of estradiol (E2), as they have been found in some treated sewage effluents, on nest holding ability and sperm quality, two measures likely related to reproductive success in fathead minnows (*Pimephales promelas*), a species with paternal nest care. Male fathead minnows were exposed for 21 days to either 50 ng/L (E2) in 25 μ L ethanol, an E2 concentration found in the effluent of some British and Canadian sewage treatment plants (STP), or to a 25 μ L ethanol control (aqueous flow-through exposure). After exposure, groups of six males from one treatment were placed into an observation aquarium with females and a limited number of spawning sites and allowed to spawn for four days. Plasma vitellogenin (VTG) concentrations, the ability to acquire and defend a nest site, and sperm quality were compared between E2 exposed and control males. E2 exposure resulted in a significant increase in VTG concentrations in E2 exposed males, but did not affect nest holding ability or sperm quality. These preliminary results indicate little effect of 21-day exposure to levels of E2 that match the highest reported E2 loads in STP effluent, and parallel our findings from a previous study using goldfish. However, caution is warranted in extrapolating these results to populations of wild fish, as other factors such as the overall estrogenic potency of effluent, the duration and timing of exposure to estrogenic compounds, and the competition between fish for spawning opportunities need to be estimated.

INTRODUCTION

The steroid 17 β -estradiol (E2) is an important hormone in endocrine pathways of fishes (Arcand-Hoy & Benson 1998) and occurs frequently in effluent discharge of municipal sewage treatment plants (STP) (Desbrow et al., 1998; Ternes et al., 1999; Rodgers-Gray et al., 2000) where it can cause endocrine

disruption in fishes (Purdom et al., 1994; Harries et al., 1997; Routledge et al., 1998). Numerous studies have assessed the effects of E2 and associated xenoestrogens on wild fish (Folmar et al., 1996; Jobling et al., 1998; Lee et al., 2000; Folmar et al., 2001), fish that were caged (Purdom et al., 1994; Harries et al., 1996; Harries et al., 1997; Routledge et al., 1998; Rodgers-Gray et al., 2001), and on fish in the laboratory (Kramer et al., 1998; Panter et al., 1998; Miles-Richardson et al., 1999; Korte et al., 2000; Bjerselius et al., 2001). All of these studies found induction of the yolk protein vitellogenin (VTG) in exposed male fish, as well as other effects such as gonadal abnormalities (Gimeno et al., 1998; Jobling et al., 1998; Miles-Richardson et al., 1999; Rodgers-Gray et al., 2001), changes in behavior (Bayley et al., 1999; Bjerselius et al., 2001), and sperm production (Gimeno et al., 1998; Bjerselius et al., 2001). However, little is known about the reproductive effects of exposure to E2 at these low concentrations. In a previous study (Schoenfuss et al., in press), we found that 50 ng/L E2 exposure resulted in minor reductions in spawning behavior and sperm production in male goldfish (*Carassius auratus*), which have no parental care. Based on these results, we designed a new experiment to determine whether exposure to similar E2 concentrations would be more detrimental to fish with greater paternal involvement in reproduction. E2 exposure concentrations were chosen to match the highest E2 concentrations reported in sewage effluent in Great Britain (Desbrow et al., 1998; Rodgers-Gray et al., 2000) and Canada (Ternes et al., 1999). In this study we exposed male fathead minnows (*Pimephales promelas*), a species with paternal nest care, to low levels of E2 and assessed the effects of exposure on the fishes ability to hold nest sites and to produce viable sperm.

The fathead minnow is widely used in toxicological studies (Ankley et al., 1998) and reproduces in the

laboratory. Fathead minnows breed year round with the male guarding a cavernous nest site until eggs hatch. The ability to defend a nest is determined by the individual's social status in which dominant males, identifiable by their tubercles, dorsal pad, and bright banding pattern, establish successful breeding territories (Cole & Smith, 1987). Smith (1974) suggested that these morphological characteristics are associated with elevated androgen levels. The male defends the nest site continuously until hatching as eggs are quickly eaten by other fathead minnows in the absence of the nest-guarding male (Sargent, 1989). Estrogen exposure could increase circulating E2 levels in male fish and suppress androgen levels, by altering neuroendocrine feedback loops (Trudeau et al., 1993; Jobling et al., 1996; Ankley et al., 1998).

We expect nest holding ability to be a sensitive indicator of reproductive success as the opportunity to spawn is conditional upon possession of a nest site (Unger, 1983; Huntingford & Torricelli, 1993). We also expect sperm production to serve as a sensitive and relevant measure of the effects of estrogen exposure as sperm production is important in externally fertilizing fish (Petersen & Warner, 1998). A reduction in sperm quantity (which normally increases in association with interactions with ovulating females (Stacey & Sorensen, 1986; Dulka et al., 1987; DeFraipont & Sorensen, 1993)) or quality can lead to reduced fertilization rates (Nakatsuru & Kramer, 1982; Zheng et al., 1997; Suquet et al., 1998; Cosson et al., 2000). By exposing fathead minnows to E2 we addressed two questions. (1) Is sperm quality in fathead minnows compromised by E2 exposure? (2) Are E2 exposed male fathead minnows able to maintain nest sites?

MATERIALS & METHODS

Experimental Animals

Fish for this study were bred and reared at the University of Minnesota. Fish were held in 28°C water (flow-through, 300 ml/min) from an in-house well, at a constant photoperiod (16 h:8 h light:dark, 0800 lights on), and were fed frozen brine shrimp (San Francisco Bay Company, CA) twice daily *ad libitum*.

Experimental Design

An experiment was conducted to assess the effects of E2 exposure on nest holding ability and sperm quality in male fathead minnows. In four trials, males were exposed for 21 days to either E2 or an ethanol control. After exposure, groups of males were placed with

females and allowed to spawn. Male behavior was observed, after which plasma VTG concentrations, and sperm quality were quantified.

Exposure

For each trial, 16 spermiating fathead minnows were netted from a stock aquarium and distributed evenly among four 20 L aquaria. Aquaria were aerated, shielded on three sides, and fish were maintained otherwise as in the stock aquaria. Males were held for 21 days in either (i) 50 ng/L E2 dissolved in 25 µl/L ethanol, or in (ii) a 25 µl/L ethanol control. Experiments with guppies (*Poecilia reticulata*) and goldfish (Bayley et al., 1999; Schoenfuss et al., in press) found no effects of ethanol exposure on spawning behavior and sperm production. Both treatments were administered via continuous flow-through delivery (280 ml/min). A peristaltic pump continuously supplied two aquaria with an E2 stock solution, while the other two aquaria received ethanol at concentrations similar to the one used to prepare the E2 stock solution. E2 was extracted weekly from E2 exposure aquaria and bi-weekly from the control aquaria. Water samples were concentrated using C18 SepPaks (Waters Corporation, MA), and then measured with an E2 ELISA test kit (Cayman Chemicals, MI). Aquarium concentrations of E2 were approximately 50 percent of the 50 ng/L nominal concentration at 24 ± 3.6 ng/L E2 (mean \pm standard error). E2 concentrations in the ethanol control aquaria were below detection limit (9 ng/L).

Spawning Assay

After 21 days, groups of six male fish from the same treatment were moved from the exposure aquaria into 160 L observation aquaria (day one). Each observation aquarium contained six mature female fathead minnows and three nest sites (3" PVC pipe sections) to ensure that only nest sites are limiting reproductive success. In preliminary studies performed with unexposed males in a similar setting, males routinely occupied all nest sites within 24 hours. Males were marked using a combination of the same three latex colors that were injected subcutaneously. A pre-experiment found no effects of latex injection on the reproductive ability of male fathead minnows.

Male fathead minnows were observed twice each morning (days two through five) for five-minute intervals. During the observation period the nest holding male in each of the three nest cavities was identified. Each nest defended by a male fathead minnow was scored as a "nest holding event" for the treatment. Data from all four replicates were pooled

resulting in a total of 48 “nest holding events” per aquarium and treatment (three nests x four observation days x four replicates).

On the fifth day, fish were anesthetized with 0.1% phenoxyethanol (Sigma, St. Louis, MO), placed upside-down into a grooved sponge, and then milt (sperm and seminal fluid) was drawn into a capillary tube by applying light abdominal pressure following established protocols (Stacey & Sorensen, 1986). Care was taken not to contaminate the sample with feces or urine. For storage, 1 µl of milt was transferred into 100 µL of isoosmotic sperm extender following published protocols (Chao et al., 1987). For the sperm analysis, 10 µl of sperm extender solution was suspended in 100 µl distilled water (1:2000 final dilution) and vortexed. The drop in osmotic pressure activated the sperm sample (Billard et al., 1995), which was then analyzed using a Hamilton-Thorn motility analyzer (Danvers, MA). Mean Progressive Velocity (MPrV), a measure of sperm velocity along its path was used to characterize sperm quality (Kime et al., 1996). Each sperm sample was analyzed three times and values were averaged to determine the MPrV for each fish. MPrV was calculated only for those fish with expressible milt.

VTG Assay

After completion of sperm analysis, blood was collected from the caudal vasculature, stored on ice, and centrifuged at 10,000 rpm for one minute. Plasma was transferred into aprotinin coated microcentrifuge tubes and stored at -20°C for later analysis. Plasma VTG concentrations were determined using a fathead minnow-specific ELISA following published protocols (Parks et al., 1999). For the statistical analysis VTG concentrations below detection limit were reported as zero.

Statistical Analysis

Sperm quality and plasma vitellogenin concentrations were analyzed using a nonparametric Mann-Whitney U test as some of the data were not normally distributed. The proportional nest holding data were analyzed using

a Fisher’s Exact test. All calculations were performed with the Prism 2.0 statistical package for the Macintosh (Graphpad Software Inc., CA).

RESULTS

Plasma VTG concentrations in E2 exposed male fathead minnows were significantly higher than in control males ($p < 0.01$, Mann-Whitney U test; Table 1). Despite VTG induction in exposed males, sperm quality did not differ between E2 exposed and control males and the MPrV was virtually identical ($p > 0.05$, Mann Whitney U test, Table 1). The ability to hold nest sites also did not differ between E2 exposed and control fish. The number of nest holding events was similar for E2 exposed and control male fathead minnows ($p > 0.05$, Fisher’s Exact test; Table 1). E2 exposed male fathead minnows were holding nest sites during all 48 observations, while control males held nests during 44 out of 48 observations. As males were color-coded it was also possible to identify changes in the identity of the nest holder from day to day. However, very few incidents in which the nest holding male changed from one day to the next were observed in either treatment.

DISCUSSION

Our preliminary results indicate that three-week exposure to 50 ng/L E2, a concentration exceeding E2 loads in the effluent of at least two of the few STP effluents that have been analyzed for E2 loads, consistently induced VTG production in mature male fathead minnows. The reproductive consequences associated with this physiological effect appear minor under the experimental conditions presented in this study. These results match our earlier findings in goldfish, for which ten week exposure to E2 at similar concentrations results in VTG induction but had relatively minor effects on spawning behavior and sperm production (Schoenfuss et al., in press). Ours is the first study to evaluate the effects of aqueous exposure to E2 at concentrations that simulate E2 loads in some STP effluents on nest holding ability and sperm quality in fathead minnows.

Table 1. Plasma vitellogenin (VTG) concentrations, sperm quality, and nest holding ability in male fathead minnows exposed to E2 or a control.

	<u>Control</u>		<u>E2</u>		
	N	Mean ± Stand. Err.	N	Mean ± Stand. Err.	
VTG (mg/ml)	23	0.01±0.006	24	**29.12 ± 0.62	p<0.01
MPrV (µm/sec)	11	45.45 ± 2.12	12	45.41 ± 0.95	ns
Nest Holding (days)		44		48	ns
(%)		92%		100%	ns

The E2 exposure protocol used in this study is notable because the absolute exposure concentration in this study (24 ng/L) is below the highest E2 loads reported for STP effluent. Although the E2 concentration in this study was higher than effluent E2 concentration reported for U.S. STP effluents, some studies have reported E2 concentrations in STP effluent as high as 88 ng/L in Great Britain (Desbrow et al., 1998, Rodgers-Gray et al., 2000) and 64 ng/L in Canada (Ternes et al., 1999). Thus, our exposure protocol represents a high, but still realistic exposure scenario. Furthermore, E2 is but one of many estrogenic compounds that have been found in STP effluent, and the overall estrogenic potency of the effluent could be much greater than simulated in this experiment.

The plasma VTG concentrations found in E2 exposed male fathead minnows in this study match those reported by Parks et al., (1999), in fathead minnows exposed to E2 via injection. Despite VTG induction, the fish appeared healthy and active and in each replicate spawning behavior could be observed within 24 h in both treatments. Nest holding ability of E2 exposed fathead minnows was also not affected by E2 exposure, with E2 exposed males holding all nest sites in all replicates of the experiment. The ability to defend a nest site is crucial for the reproductive success of male fathead minnows as eggs in any unguarded nest are quickly eaten by other fathead minnows (Sargent, 1989). Our results are similar to those reported in two studies on guppies that did not find reproductive impairment at 30 ng/L E2 exposure (Toft & Baatrup, 2001), and saw no behavioral impairment at 100 ng/L E2 exposure (Bayley et al., 1999). In contrast, a previous study conducted in our laboratory (Schoenfuss et al. in press) found a significant decline in reproductive behavior in male goldfish exposed to 50 ng/L E2 for ten weeks. Bjerselius et al., (2001), also found behavioral impairment at E2 exposure concentrations exceeding 1 µg/L E2, but that E2 exposure level is far above any recorded for STP effluents.

MPrV, a measure of sperm quality did not vary between E2 exposed and control fish. Sperm quality has been suggested previously (Kime et al., 1996; Kime 1998; Kime & Nash, 1999) as a tool to analyze the effects of EDCs on the reproductive fitness of fish. However, despite the strong physiological response (VTG induction) in male fish exposed to E2, sperm quality appeared unaffected. In a previous study using goldfish (Schoenfuss et al., in press), the absolute number of motile sperm per fish, a product of sperm quality and sperm quantity, declined with 50 ng/L E2 exposure after ten weeks. In contrast, Toft and Baatrup

(2001) saw an increase in sperm numbers in guppies exposed to 30 ng/L E2. In this experiment we were unable to measure sperm quantity as the small size of the fathead minnow renders it impossible to strip consistently all milt while avoiding contamination by urine.

Although we did not find any impairment of nest holding ability and sperm quality in male fathead minnows exposed to E2, we remain cautious about extrapolating these laboratory results to populations of wild fish. Indeed, ongoing studies in our laboratory suggest that intra-species interactions might be an important component in evaluating the effects of E2 exposure. Furthermore, STP effluent consists of complex mixtures of chemicals, many of which have been shown to be estrogenic to fish and their combined action may exceed our exposure concentrations. Our understanding of the effects of STP effluent on fish populations is still rudimentary and further studies will be necessary to determine the effects of estrogen and xenoestrogen exposure on all life stages of fish. Answering these questions will allow us to determine the effects of effluent exposure on wild fish populations.

ACKNOWLEDGEMENTS

The authors acknowledge funding from the National Sea Grant College Program. We thank Gary Ankley and the staff at the Duluth EPA laboratory for help with setting up a fathead minnow breeding colony and the VTG assay. We are grateful to Mats Troedsson (School of Veterinary Medicine, University of Minnesota) for making the computer assisted motility analyzer available to us, and Kate Loseth for providing training on its use.

AUTHORS

Heiko L. Schoenfuss, Dalma Martinovic, and Peter W. Sorensen, Department of Fisheries, Wildlife, and Conservation Biology, 200 Hodson Hall, 1980 Folwell Avenue, University of Minnesota, St. Paul, MN 55108

REFERENCES

- Ankley, G. T., E. Mihaich, R. Stahl, D. Tillitt, T. Colborn, S. McMaster, R. Miller, J. Bantle, P. Campbell, N. Denslow, R. Dickerson, L. Folmar, M. Fry, J. Giesy, L. Earl Gray, P. Guiney, T.

- Hutchinson, S. Kennedy, V. Kramer, G. LeBlanc, M. Mayes, A. Nimrod, R. Patino, R. Peterson, R. Purdy, R. Ringer, P. Thomas, L. Touart, G. Van Der Kraak, T. Zacharewski. 1998. *Overview of a Workshop on Screening Methods for Detecting Potential (Anti-) Estrogenic/Androgenic Chemicals in Wildlife*. Environ Toxicol Chem 17, 68-87.
- Arcand-Hoy, L. D., W. H. Benson. 1998. *Fish Reproduction: An Ecologically Relevant Indicator of Endocrine Disruption*. Environ Toxicol Chem 17, 49-57.
- Bayley, M., J. R. Nielsen, E. Baatrup. 1999. *Guppy Sexual Behavior as an Effective Biomarker of Estrogen Mimics*. Ecotoxicol Environ Safety 43, 68-73.
- Billard, R., J. Cosson, G. Perchee, O. Linhart. 1995. *Biology of Sperm and Artificial Reproduction in Carp*. Aquaculture 129, 95-112.
- Bjerselius, R., K. Lundstedt Enkel, H. Olsen, I. Mayer, K. Dimberg. 2001. *Male Goldfish Reproductive Behavior and Physiology Are Severely Affected by Exogenous Exposure to 17 β -Estradiol*. Aquat Toxicol 53, 139-152.
- Chao, N-H., W-C. Chao, K-C. Liu, I-C. Liao. 1987. *The Properties of Tilapia Sperm and its Cryopreservation*. J Fish Biol 30, 107-118.
- Cole, K. S., R. J. F. Smith. 1987. *Male Courting Behaviour in the Fathead Minnow, Pimephales Promelas*. Environ Biol Fish 18, 235-239.
- Cosson, J., O. Linheart, S. D. Mims, W. L. Sheldon, M. Rodina. 2000. *Analysis of Motility Parameters from Paddlefish and Shovelnose Sturgeon Spermatozoa*. J Fish Biol 56, 1348-1367.
- DeFraipont, M., P. W. Sorensen. 1993. *Exposure to the Pheromone 17 α , 20 β -Dihydroxy-4-Pregnen-3-One Enhances the Behavioral Spawning Success, Sperm Production and Sperm Motility of Male Goldfish*. Animal Behav 46, 245-256.
- Desbrow, C., E. J. Routledge, G. C. Brighty, J. P. Sumpter, M. Waldock. 1998. *Identification of Estrogenic Chemicals in STW Effluent. I. Chemical Fractionation and aid In-Vitro Biological Screening*. Environ Sci Technol 32, 1549-1558.
- Dulka, J. G., N. E. Stacey, P. W. Sorensen, G. J. Van Der Kraak. 1987. *A Steroid Sex Pheromone Synchronizes Male-Female Spawning Readiness in Goldfish*. Nature 325, 251-253.
- Folmar, L., N. D. Denslow, K. Kroll, E. F. Orlando, J. Enblom, J. Marcino, C. Metcalfe, L. J. Guillette Jr. 2001. *Altered Serum Sex Steroids and Vitellogenin Induction in Walleye (Stizostedion Vitreum) Collected Near a Metropolitan Sewage Treatment Plant*. Arch Environ Contam Toxicol 40, 392-398.
- Folmar, L. C., N. D. Denslow, V. Rao, M. Chow, A. Crain, J. Enblom, J. Marcino, L. J. Guillette Jr. 1996. *Vitellogenin Induction and Reduced Serum Testosterone Concentrations in Feral Male Carp (Cyprinus Carpio) Captured Near a Major Metropolitan Sewage Treatment Plant*. Environ Health Perspect 104, 1096-1100.
- Gimeno, S., H. Komen, S. Jobling, J. Sumpter, T. Bowmer. 1998. *Demasculinisation of Sexually Mature Male Common Carp, Cyprinus Carpio, Exposed to 4-Tert-Pentylphenol During Spermatogenesis*. Aquat Toxicol 43, 93-109.
- Harries, J. E., D. A. Sheahan, S. Jobling, P. Matthiessen, P. Neall, J. P. Sumpter, T. Tylor, N. Zaman. 1997. *Estrogenic Activity in Five United Kingdom Rivers Detected by Measurement of Vitellogenesis in Caged Male Trout*. Environ Toxicol Chem 16, 534-542.
- Harries, J. E., D. A. Sheahan, S. Jobling, P. Matthiessen, P. Neall, J. P. Sumpter, T. Tylor, N. Zaman. 1996. *A Survey of Estrogenic Activity in United Kingdom Inland Waters*. Environ Toxicol Chem 15, 1993-2002.
- Huntingford, F., P. Torricelli. 1993. *Behavioural Ecology of Fishes*. Erice, Italy.
- Jobling, S., D. Sheahan, J. A. Osborne, P. Matthiessen, J. Sumpter. 1996. *Inhibition of Testicular Growth in Rainbow Trout (Oncorhynchus Mykiss) Exposed to Estrogenic Alkylphenolic Chemicals*. Environ Toxicol Chem 15, 194-202.
- Jobling, S., M. Nolan, C. R. Tyler, G. Brighty, J. P. Sumpter. 1998. *Widespread Sexual Disruption in Wild Fish*. Environ Sci Technol 32, 2498-2506.
- Kime, D. E. 1998. *Endocrine Disruption in Fish*. Boston: Kluwer Academic Publishers.
- Kime D. E., J. P. Nash. 1999. *Gamete Viability as an Indicator of Reproductive Endocrine Disruption in Fish*. Sci Total Environ 233, 123-129.

- Kime, D. E., M. Ebrahimi, K. Nysten, I. Roelants, E. Rurangwa, H. D. M. Moore, F. Olivier. 1996. *Use of Computer Assisted Sperm Analysis (CASA) for Monitoring the Effects of Pollution on Sperm Quality of Fish. Application to the Effects of Heavy Metals*. *Aquat Toxicol* 36, 223-237.
- Korte, J. J., M. D. Kahl, K. M. Jensen, M. S. Pasha, L. G. Parks, G. A. LeBlanc, G. T. Ankley. 2000. *Fathead Minnow Vitellogenin: cDNA Sequence and mRNA and Protein Expression After 17 β -estradiol Treatment*. *Environ Toxicol Chem* 19, in press.
- Kramer, V. J., S. Miles-Richardson, S. L. Pierens, J. P. Giesy. 1998. *Reproductive Impairment and Induction of Alkaline-Labile Phosphate, A Biomarker of Estrogen Exposure in Fathead Minnows (Pimephales Promelas) Exposed to Waterborne 17 β -Estradiol*. *Aquat Toxicol* 40, 335-360.
- Lee, K. E., V. S. Blazer, N. C. Denslow, R. M. Goldstein, P. J. Talmage. 2000. *Use of Biological Characteristics of Common Carp (Cyprinus Carpio) to Indicate Exposure to Hormonally Active Agents in Selected Minnesota Streams, 1999*, pp. 47. Mounds View, MN: U.S. Geological Survey.
- Miles-Richardson, S., V. J. Kramer, S. D. Fitzgerald, J. A. Render, B. Yamini, S. J. Barbee, J. P. Giesy. 1999. *Effects of Waterborne Exposure of 17 β -Estradiol on Secondary Sex Characteristics and Gonads of Fathead Minnows (Pimephales Promelas)*. *Aquat Toxicol* 47, 129-145.
- Nakatsuru, K., D. L. Kramer. 1982. *Is Sperm Cheap? Limited Male Fertility and Female Choice in the Lemon Tetra (Pisces, Characidae)*. *Science* 216, 753-755.
- Panter, G. H., R. S. Thompson, J. P. Sumpter. 1998. *Adverse Reproductive Effects in Male Fathead Minnows (Pimephales Promelas) Exposed to Environmentally Relevant Concentrations of the Natural Oestrogens, Oestradiol and Oestrone*. *Aquat Toxicol* 42, 243-253.
- Parks, L. G., A. O. Cheek, N. D. Denslow, S. A. Heppell, J. A. McLachlan, G. A. LeBlanc, C. V. Sullivan. 1999. *Fathead Minnow Pimephales Promelas Vitellogenin: Purification, Characterization, and Quantitative Immunoassay for the Detection of Estrogenic Compounds*. *Comp Biochem Physiol* 123, 113-125.
- Petersen, C. W., R. R. Warner. 1998. *Sperm Competition in Fishes*. In *Sperm Competition and Sexual Selection*, (ed. T. R. Birkhead, and A. P. Moller), pp. 435-464. New York, Academic Press.
- Purdom, C. E., P. A. Hardiman, V. J. Bye, N. C. Eno, C. R. Tyler, J. P. Sumpter. 1994. *Estrogenic Effects of Effluents From Sewage Treatment Works*. *Chem Ecol* 8, 275-285.
- Rodgers-Gray, T., S. Jobling, C. Kelly, S. Morris, G. Brightly, M. J. Waldock, J. P. Sumpter, C. R. Tyler. 2001. *Exposure of Juvenile Roach (Rutilus Rutilus) to Treated Sewage Effluent Induces Dose-Dependent and Persistent Disruption in Gonadal Duct Development*. *Environ Sci Technol* 35, 462-470.
- Rodgers-Gray, T., S. Jobling, S. Morris, C. Kelly, S. Kirby, A. Janbakhsh, J. E. Harries, M. J. Waldock, J. P. Sumpter, C. R. Tyler. 2000. *Long-Term Temporal Changes in the Estrogenic Composition of Treated Sewage Effluent and Its Biological Effects on Fish*. *Environ Sci Technol* 34, 1521-1528.
- Routledge, E. J., D. Sheahan, C. Desbrow, G. C. Brightly, M. Waldock, J. P. Sumpter. 1998. *Identification of Estrogenic Chemicals in STW Effluent. 2. In Vivo Responses In Trout And Roach*. *Environ Sci Technol* 32, 1559-1565.
- Sargent, R. C. 1989. *Allopaternal Care in the Fathead Minnow, Pimephales Promelas: Stepfathers Discriminate Against Their Adopted Eggs*. *Behav Ecol Sociobiol* 25, 379-385.
- Schoenfuss, H. L., J. T. Levitt, G. Van Der Kraak, P. W. Sorensen. in press. *Ten Week Exposure to Treated Sewage Discharge Has Relatively Minor, Variable Effects on Reproductive Behavior and Sperm Production in Goldfish*. *Environ Toxicol Chem*.
- Stacey, N. E., P. W. Sorensen. 1986. *17 α ,20 β -Dihydroxy-4-Pregnen-3-One: A Steroidal Primer Pheromone Which Increases Milt Volume in Goldfish, Carassius Auratus*. *Can J Zool* 64, 2412-2417.
- Suquet, M., C. Dreanno, G. Dorange, Y. Normant, L. Quemener, J. L. Giagnon, R. Billard. 1998. *The Aging Phenomenon of Turbot Spermatozoa: Effects on Morphology, Motility, and Concentration*.

- Intracellular ATP Content, Fertilization, and Storage Capacities*. J Fish Biol 52, 31-41.
- Smith, R. J. F. 1974. *Effects of 17 α -Methyltestosterone on the Dorsal Pad and Tubercles of Fathead Minnows (*Pimephales Promelas*)*. Can J Zool 52,1031-1038.
- Ternes, T., M. Stumpf, J. Mueller, K. Haberer, R-D. Wilken, M. Servos. 1999. *Behavior and Occurrence of Estrogens in Municipal Sewage Treatment Plants - I. Investigations In Germany, Canada, And Brazil*. Sci Total Environ 225, 81-90.
- Toft, G., E. Baatrup. 2001. *Sexual Characteristics Are Altered by 4-Tert-Octylphenol and 17 β -Estradiol in the Adult Male Guppy*. Ecotoxicol Environ Safety 46, 76-84.
- Trudeau, V. L., M. G. Wade, G. Van Der Kraak, R. E. Peter. 1993. *Effects of 17 β -Estradiol on Pituitary and Testicular Function in Male Goldfish*. Can J Zool 71, 1131-1135.
- Unger, L. M. 1983. *Nest Defense by Deceit In the Fathead Minnow, *Pimephales Promelas**. Behav Ecol Sociobiol 13, 125-130.
- Zheng, W., C. Strobeck, N. Stacey. 1997. *The Steroid Pheromone 4-Pregnen-17 α , 20 β -Diol-3-One Increases Fertility and Paternity in Goldfish*. J Exp Biol 200, 2833-2840.