# A Study of Phytochemical Composition of a Few Tribal Medicinal Plants from Sriharikota

\*R. Bharath Kumar and \*\*B. Suryanarayana

\*Associate Professor, Dept.Biotechnology, Vignan's Engineering College, Vadlamudi, Guntur-522213. email: <a href="mailto:drbharathravuru@gmail.com">drbharathravuru@gmail.com</a>

\*\*Associate Professor (Retd.), #8-307, Karnakamma Street, Venkatagiri Town -524132.

email: dr\_bsnarayana@yahoo.com

**Issued 30 January 2009** 

## **ABSTRACT**

SRIHARIKOTA acquired importance botanically because of rich Island vegetation and due to installation of Rocket Launching Station by Indian Space Research Organisation (ISRO). A few Tribal colonies are domiciled there. Some aged tribal men conversed with herbal medicines and practices. With a fear that this tribal medicinal knowledge of the island may be lost, if urgently not pursued in a recorded manner for the benefit of modern society, a project on Ethnobotany and tribal medicines of Sriharikota Island is taken up three years back in 1996. Regular field tours are conducted covering all the seasons and about 300 medicinal plants are collected along with ample field notes on folklore medicinal uses with the support of local aged tribal men. Out of these, a few plants are selected on the basis of endemism and utility and subjected to phytochemical analysis. Investigation for 11 chemical components is made in 21 samples. All the data is recorded in this paper. The results are mostly in conformity with the medicinal uses and they are discussed.

## Introduction

Yanadies an aboriginal tribe are in Sriharikota Island in Andhra Pradesh even after establishment of Rocket Launching Station. They are said to have been migrated from Malaya Peninsula, Africa or Australia.

Until the establishment of SHAR Centre at Sriharikota, yanadies used to live in forests and near sea coast in traditional way, drifted from the natural way of life due to agro-rural developmental activities a few aged tribal men are able to furnish 'ethno-medicinal data' pertaining to their traditional practices and healings. With the help of local tribal men, 300 ethnomedicinal plants are recorded. Based on their pharmaceutical uses they are classified into 16 – categories [Annexure II].

Out of 300 species surveyed 21 samples falling under 6 –categories are selected for phytochemical screening. Out of them 14 species are endemic/rare. The results are discussed in detail in this article.

## **Topography**

Sriharikota Island geometrically is located at 800 .21' E and 130 .22' to 140 N. It is a spindle shaped land mass sandwitched between Bay of Bengal on the east and Pulicat lake on west. It is 18 km east of Sullurpet, the nearest Railway Station connecting Madras-Kolkata trunk line. Madras is 98 km away from Sriharikota.

#### **Materials and Methods**

Intensive medico-ethnobotanical survey is conducted in Sriharikota Island for 3 years since 1996-99. Field tours at regular intervals were conducted covering all the seasons so as not to miss seasonal elements having pharmaceutical value. Plants are selected for phytochemical screening are popularly used by tribals for their general ailments [Annexure 1].

Phytochemical screening was done by the standard procedures prescribed by Bhattacharya (1956), later modified by Chhabra et.al (1984) and Harborne (1973, 1977).

#### **Preparation of Extracts**

The plant parts (root bark, stem, leaf and whole plant etc.) were washed with water, chopped into small fragments and shade dried. The dried samples were ground to power (each 200 gr.) and stored in polythene containers at room temperature. These samples are used for screening to detect the different classes of chemical constituents. Extracts of sample are prepared by taking of 20 gr. of sample in 200 ml. methanol. Each sample is tested for 11 components Results are given in Annexure I.

16 Field tours of 5-7 days duration of each tour are conducted for three years during 1996-99. with the assistance of local tribal men. 300 species having medicinal uses and ample field notes also are collected regarding flowering, fruiting, flower colour and smell etc; phenological data. Further data pertaining to ethnomedicinal information of each species also recorded. 21 plant samples are choosen for phytochemical screening based on their popular pharmaceutical importance, coupled with their endemic or rare in occurrence. Active constituents present in the plant species in different plant parts (root bark, stem, whole plant, leaves etc.) are responsible for their therapeutic effects. A preliminary phytochemical analysis is conducted to detect the presence of alkaloids, saponins, tannins, carbohydrates, proteins, steroids, trterpinoids, amino acids etc., which act as possible curing agent during folk therapy.

Tests are conducted for 11 components as detailed in the Annexure-I. Their occurrence in noted with symbol (+)in the table given total number of components present in each sample is given in the last column (17) in parenthesis. The table also contains tribal medicinal uses and other folklore medicinal uses, so as to relate medicinal properties with the components of positive occurrence. Further these samples also grouped into 6 broad pharmaceutical categories as given in Annexure-II. Results of the tests for chemical constituents of each of the pharmaceutical group are discussed in detail, here under.

## P.C. No. 5: RHEUMATISM AND ARTHRITIS

Seven test samples have PC 5 property based on their tribal /folklore medicinal use.

Upon chemical analysis five components viz. alkaloids, flavonoids, phenols, amino acids, steroidal nucleus are positive in variable combinations in 7 samples. Saponins occur in 4 samples, tannins occur in 2 samples. Presence of alkaloids, flavonoids, phenols, amino acids, steroidal nucleus in the samples consistently in variable combinations and minimum of two components positive for any 2 samples is indicative of having curative effect for PC 5 either individually or cumulatively

#### PC No. 7: SEXUAL PROBLEMS

4 samples positive for alkaloids (t 6)

Flavonoids (t 7), phenols (t 8), saponins (t 9) and steroidal nucleus (t 15) are positive for 3 samples out of 4 tested samples variably.

Thus alkaloids seems to have curative effect in combination with flavonoids, phenols, saponins and steroidal

nucleus for PC 7.

# PC No. 8: SKIN & DERMATITIS

2 samples when tested, out 11 components resulted positive for four tests i.e., t 7, 8, 9, and 15 viz. flavonoids, phenols, steroids/triterpinoids and steroidal nucleus.

# PC NO. 9: OPTHALMIC & ENT

5 samples are reported for PC 9

All the 5samples are positive for flavonoids (t7) where as 3 samples are positive in variable order for alkaloids (t 6), phenols (t 8), saponins (t 9) and tannins (t 10).

All the 5 components occur in 1 sample i.e., Eugenia bracteata.

Thus flavonoids (t7) seem to be curative component either indivially as in *Capparis brevispina* or cumulatively in other samples.

## PC No. 13: Cooling and Alterative

1 sample – Colubrina asiatica shown positive for t 7, t 9, t 13, and t 15.

## PC No.16: Diabetes

1 sample tested for 11 components. Interestingly it is positive for 9 components. T 12 and t 16 are negative. Out of nine components which of them is specific curative is not certain for PC 16.

#### Annexure-II

## PHARMACEUTICAL CLASSIFICATION

PC	Pharmaceutical Category [P.C.]	No. of
Category		Species
No.		Used
1.	Respiratory and Bronchial problems	
2.	Cordiac	
3.	Urinary and Kidney	
4.	Piles and Constipation	
5.	Arthritis and Rheumatism	7
6.	Anodyne elements	
7.	Sexual problems	4
8.	Skin and Dermatitis	2
9.	Opthalmic and E.N.T	5
10.	Psychoactive and Nervous	
11.	Jaundice and Liver disorders	
12.	Digestive and Stomachic	
13.	Cooling and Alterative	1
14.	Antipyretic, Anthelmintic and	
	Antiperiodic	
15.	Antidotes and Antivenom	
16.	Diabetes	1

## Conclusion

21 Phyto-samples falling under 6 pharmaceutical categories are investigated for 11- chemical components. Results obtained are analysed to note that the presence of specific classes of chemical constituents are in confirmity with the tribal medico-therapy, given in Annexure –II.

Further it is interesting to note that 14 samples out of 21 belong to endemic/ rare species. In view of Endemism/Rarity of the species chosen for phytochemical analysis, these specimens need further investigation to detect specific active principles of curative effect.

#### Acknowledgements

Authors are thankful to the Principal, Visodava Government College, Venkatagiri, for providing facilities and encouragement to the Director, SHAR Centre, Sriharikota, for providing facilities to carryout field work and to Dr. V. Veerraju, Head, C& LD Division, SHAR Centre, Sriharikota, for help during field work. Authors are also thankful to ISRO, Bangalore, for extending financial assistance.

#### **Literature Cited**

Anonymous, 1948-76... *The Wealth of India (Raw Materials)*. Vol. 1-11.CSIR, New Delhi, India.

Bhattacharya, S.C. 1956 a ... Constituents of *Centella asiatica* Linn. Examination of the Ceylonese Variety. *J. Indian. Chem. Soc.* 33: 599-586.

Chhabra, S.C., F.C.Viso and E.N.Mshiu. 1984... Phytochemical screening of Tanzanian Medicinal plants 1. *J. Ethno pharmacology* 11: 157-179.

Chopra, R.N., S.L.Nayar & I.C.Chopra, 1956... Glossary of Indian Medicinal plants.

CSIR, New Delhi..

Harborne, J.B. 1984... *Phytochemical methods : A guide to Modern Techniques of plant Analysis.* 1 - 278 P. Chapman and Hall, London.

Nagaraju, N and Rao, K.N. 1994... Plant Phenolics of some Indegenous medicinal pants of Apocynaceae.

Higher Plants of Indian subcontinent.(Addit. Series of Indian Jour. of Forestry No. VI). 211-215.

Rameshwar Dayal 1997... Chemical Investigation of Medicinal Plants. *Indian Forester*, (1): 249-255.

Saiprasad Goud, P. 1995... Ethno-medico-Botanical studies in Kurnool district of Andhra Pradesh and screening of selected species for Biological activity. Ph.D. Thesis, S.K.University, Anantapur.


Sudarsanam, G. and N.S. Balaji Rao. 1994... Medicinal Plants used by the Yanadi tribe of Nellore district.


Andhra Pradesh. India. Bull. Pure and Applied Sci. 13b: 65-70.


Suryanarayana, B., A.S.Rao, A.M.Rao and V.Veerraju. 1998. Flora of Sriharikota Island – Technical Report. I. S.R.O., Bangalore.


Ullah, N, Ahmed, Z, Anis, F. and Malik, A. 1997...Phytochemical Investigation of *Daphne oleoides*. *Fitoterapia*. Vol. 1.XIX, No.3.280.

Presented at **Indian Science Congress** (ISCA), 3 –7<sup>th</sup> January 2000, Pune University, **Pune**.









# Annexure 1

## PHYTOCHEMICAL SCREENING OF SOME SELECTED TRIBAL MEDICINAL PLANTS

Tribal Medicinal Plant species is about 19 no., with their crude drug samples 21 (leaf, Root, Root bark, stem, stem bark and fruit etc.) were tested against for 11 Phytoactive components (Phytochemical screening), which are widely present in various plant species their respective results are also given here. (\*with their Phytoactive sample nos.)

S.No. Bn., Ln., Family Part Tribal uses in Reported other used island uses

# PC NO.5; ARTHRITIS AND RHEUMATISM:

1. Breynia vitis-idea Lf. Warmed with Leaves smoked Adavimunaga Dalbergia paniculata like tobacco in EUPHORBIACEAE leaves and applied in rheumatic pains. Leaves smoked smoked like tobacco in and tonsils

Phytoactive components are present in the drug sample/s: 1(+), 2(+), 5(+), 6(+), 7(+), 8(+), 9(+), 10(+), 11(+), Total: 9.

2.ClerodendrumLf.Warmed and appliedLeaf juice givenphlomidisfor filarial andin neglectedPeddanelli/ TakkedarheumaticsyphiliticVERBENACEAEswellings.complaints.

Phytoactive components are present in the drug sample/s: 1(+), 2(+), 5(+), 7(+), 8(+), 9(+), 10(+), 11(+) Total:

3. Crotalaria Lf. Leaf paste with Infusion of laburnifolia coconut oil applied the Wh.plant for

FABACEAE. inflammation of

the

mouth.

Phytoactive components are present in the drug sample/s: 2(+), 3(+), 4(+), 6(+), 7(+), 9(+), 10(+), 11(+),

Total: 8.

**4.** Dalbergia Lf. Warmed, applied for —

paniculata rheumatic swellingsPacchari and painful parts.

Phytoactive components are present in the drug sample/s: 1(+), 2(+), 3(+), 6(+), 8(+), 9(+), 10(+), 11(+) Total:

8.

r5. Garcinia spicata Rt.B Paste applied for —

Pidatha / pains and rheumatic

Errijambi swellings.

CLUSIACEAE.

Phytoactive components are present in the drug sample/s: 1(+), 2(+), 3(+), 4(+), 6(+), 7(+), 8(+), 9(+), 10(+),

11(+) **Total: 8.** 

6.*Holoptelia*St.B Warmed and applied Juice of boiled
integrifolia on rheumatic bark applied on
Peddamanu/ swellings. rheumatic swellings.

Tabasu,

ULMACEAE.

Phytoactive components are present in the drug sample/s: 2(+), 5(+), 6(+), 10(+) Total: 4.

7. Pisonea aculeata Rt.B Leaves+Rt.B warPeetrinchi med and applied for used for swellings
NYCTAGINACEAE rheumatic swellings. & rheumatic pains

Phytoactive components are present in the drug sample/s: 1(+), 3(+), 4(+), 5(+), 6(+),

10(+) **Total: 6.** 

PC.No.7; SEXUAL PROBLEMS:

8. *Capparis* Rt.B Used for treatment Cholera, zeylanica for janni (Post natal stomachic and

Aridonda complications of cholagogue. CAPPARACEAE. ladies: fever accom-

panied with chillness).

Phytoactive components are present in the drug sample/s: 1(+), 2(+), 5(+), 10(+)

Total: 4.

9.*Crinum defixum*Porla gadda/
Bul. Scale leaves warmed Emetic, diaphoand applied for retic used for

Chembu gadda swellings of testis. treatment of the

**AMARYLIDACEAE** 

burns.

Phytoactive components are present in the drug sample/s: 1(+), 4(+), 6(+) Total: 4.

10.**Hybanthus** Lf. Powder with turmeric Plant tonic, diuretic; leaves enneaspermus powder taken orally for general debility Ratnapurusha and tender sta-**VIOLACEAE** and vigor. lks demulcent. Phytoactive components are present in the drug sample/s: 1(+), 2(+), 3(+), 4(+), 9(+), 10(+) **Total:6.** 11.Linociera Rt. Paste taken orally in case of sexual diseases zeylanica Punagani (syphilis, gonorrhoea) **OLEACEAE** and menstrual problems. Phytoactive components are present in the drug sample/s: 1(+), 2(+), 3(+), 10(+) Total: 4. PC.No.8; SKIN AND DERMATITIS: 12.Pisonea Lf. Leaf curry orally Juice mixed with aculeata given to children pepper administefor mumps and red for pulmonary Peetrinchi **NYCTAGINACEAE** leprosy. complaints of children. Phytoactive components are present in the drug sample/s: 2(+),3(+),8(+),10(+) Total:4. r13.Ventilago Rt.B. Paste applied for Juice remedy for pains, and madaraspatana skin deseases. Churidi/Surudu malarial fever. RHAMNACEAE. Phytoactive components are present in the drug sample/s: 1(+),2(+),3(+),4(+),6(+),7(+),8(+),9(+),10(+),11(+)**Total:4.** PC.No.9; OPTHALMIC AND E.N.T.: 14. Capparis Rt. Paste used in brevispina tooth ache and Palasuri infected gums. **CAPPARACEAE** Phytoactive components are present in the drug sample/s: 2(+) Total: 1. r<sub>15.</sub>C. rotundifolia Rt. Paste applied Nagetimullu / incase of head Suramullu ache CAPPARACEAE. Phytoactive components are present in the drug sample/s: 1(+), 2(+), 3(+), 4(+), 5(+)Total: 5. r<sub>16</sub>.Cymbidium Lf. Juice warmed and Emetic, purgative administered nutrient and aloifolium Thatisaga incase of earache. demulcent. **ORCHIDACEAE** Phytoactive components are present in the drug sample/s:2(+),4(+),5(+),8(+) Total:4. \*17.Eugenia Lf. Smoke used as bracteata mosquito repellent. Kundanedu /

Kundaneredu

**MYRTACEAE** 

Phytoactive components are present in the drug sample/s: 1(+),2(+)3(+),4(+),5(+),

6(+),7(+),8(+),9(+),10(+),11(+) **Total:11.** 

\*18.*E. bracteata* Rt. Paste mixed with —

Kundanedu / goat milk and Kundaneredu applied for mumps.

**MYRTACEAE** 

Phytoactive components are present in the drug sample/s: 1(+), 2(+), 4(+), 5(+),

6(+),7(+),9(+),10(+),11(+) **Total:10.** 

19. Sarcostemma St. Juice given to chilacidum dren to get relief bitter, cooling.

Pulla teegalu from cold, Latex

ASCLEPIADACEAE dropped in eyes in

case of cataracht for remedy.

Phytoactive components are present in the drug sample/s: 1(+), 2(+), 3(+), 6(+), 8(+),

9(+),10(+)**Total:7.** 

PC.No.13; COOLING AND ALTERATIVE:

\*20.*Colubrina* Lf. Juice taken as tonic. —

asiatica Neetipulavachettu RHAMNACEAE

Phytoactive components are present in the drug sample/s:2(+), 4(+),8(+),10(+) Total:4.

PC.No.16; DIABETES:

\*21. Casearia Rt. Decoction effective Promotes action

esculentaremedy for diabetesof liver. Decoc-Kunda jungerution used inSAMYDACEAE.diabetes and piles.

Phytoactive components are present in the drug sample/s: 1(+),2(+),3(+),4(+),5(+),

6(+), 8(+), 9(+), 10(+)**Total:9.** 

# ABBREVIATIONS USED IN THE TEXT:

1.Alka: Alkaloids2.Flavo.:Flavonoids3.Phen.: Phenols4.Sapo.: Saponins5.Tann.: Tannins

**6.Carbo:** Carbohydrates

**7.Pro.:** Proteins

**8.Stero.** /**Triter.:** Steroids/Triterpinoids

**9.Amino.:** Amino acids

10.Ster.Nuc.: Steroidal nucleus11.Leu.antho.: Leuco anthocyanins

**Total: Total Components** 

PLANT SPECIES : 19. CRUDE DRUG SAMPLES : 21.

ABBREVIATIONS : St.: Stem, St.B: Stem bark, Rt.: Root, Rt.B: Root

bark, Lf.: Leaf, Fr.: Fruit and

PC.No.: Pharmaceutical category number Bn: Botanical name, Ln: Local name and

Fy: Family.\*: Endemic; r: Rare

COLOUR APPEARENCE : R: Red, Y: Yellow, B: Blue, G: Green, P: Pink

(or) Rose, M: Majanta (or) Crimson & V: Violet.

Plant samples extracts are prepared with methanol. (Universal solvent).