

2011

A Characterization of Power Method Transformations through L -Moments

Todd C. Headrick

Southern Illinois University Carbondale, headrick@siu.edu

Follow this and additional works at: http://opensiuc.lib.siu.edu/epse_pubs

Published in *Journal of Probability and Statistics*, Vol. 2011 (2011) at [10.1155/2011/497463](https://doi.org/10.1155/2011/497463)

Recommended Citation

Headrick, Todd C. "A Characterization of Power Method Transformations through L -Moments." (Jan 2011).

This Article is brought to you for free and open access by the Educational Psychology and Special Education at OpenSIUC. It has been accepted for inclusion in Publications by an authorized administrator of OpenSIUC. For more information, please contact opensiuc@lib.siu.edu.

Research Article

A Characterization of Power Method Transformations through L -Moments

Todd C. Headrick

*Section on Statistics and Measurement, Department EPSE, 222-J Wham Building, Mail Code 4618,
Southern Illinois University Carbondale, Carbondale, IL 62901, USA*

Correspondence should be addressed to Todd C. Headrick, headrick@siu.edu

Received 4 July 2010; Revised 27 December 2010; Accepted 20 January 2011

Academic Editor: Rongling Wu

Copyright © 2011 Todd C. Headrick. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Power method polynomial transformations are commonly used for simulating continuous non-normal distributions with specified moments. However, conventional moment-based estimators can (a) be substantially biased, (b) have high variance, or (c) be influenced by outliers. In view of these concerns, a characterization of power method transformations by L -moments is introduced. Specifically, systems of equations are derived for determining coefficients for specified L -moment ratios, which are associated with standard normal and standard logistic-based polynomials of order five and three. Boundaries for L -moment ratios are also derived, and closed-formed formulae are provided for determining if a power method distribution has a valid probability density function. It is demonstrated that L -moment estimators are nearly unbiased and have relatively small variance in the context of the power method. Examples of fitting power method distributions to theoretical and empirical distributions based on the method of L -moments are also provided.

1. Introduction

The power method (see Fleishman [1], Headrick [2], and [3–7]) is a traditional procedure used for simulating continuous nonnormal distributions. Some applications of the power method have included such topics as ANOVA [8–10], asset pricing theories [11], business-cycle features [12], cluster analysis [13], item parameter estimation [14], item response theory [15, 16], factor analysis [17–19], price risk [20], structural equation models [21–24], and toxicology [25].

In the context of univariate data generation and for the purposes considered herein, the power method can be generally summarized by the polynomial transformation as in

Headrick [2, equation (2.7)]

$$p(W) = \sum_{i=1}^m c_i W^{i-1}, \quad (1.1)$$

where W can be either a standard normal (Z) or a standard logistic (L) random variable with probability density function (pdf) and cumulative distribution function (cdf)

$$f_W(w) = \begin{cases} f_{W:=Z}(w := z) = \phi(z) = (2\pi)^{-(1/2)} \exp\left\{-\frac{z^2}{2}\right\}, \\ f_{W:=L}(w := l) = \frac{(\pi/\sqrt{3}) \exp\left\{-(\pi/\sqrt{3})l\right\}}{\left(1 + \exp\left\{-(\pi/\sqrt{3})l\right\}\right)^2}, \end{cases} \quad (1.2)$$

$$F_W(w) = \begin{cases} F_{W:=Z}(w := z) = \Phi(z) = \int_{-\infty}^z \phi(u) du, & -\infty < z < +\infty, \\ F_{W:=L}(w := l) = \frac{1}{1 + \exp\left\{-(\pi/\sqrt{3})l\right\}}, & -\infty < l < +\infty. \end{cases}$$

Setting $m = 4$ (or $m = 6$) in (1.1) gives the Fleishman [1] (or Headrick [3]) class of distributions associated with standard normal-based polynomials. The shape of $p(W)$ in (1.1) is contingent on the coefficients c_i , which are determined by moment-matching techniques. For example, see Headrick [2, equations (2.18)–(2.21); (2.22)–(2.25)] for determining the coefficients associated with standard normal or standard logistic-based polynomials for $m = 4$ in (1.1).

The pdf and cdf associated with $p(W)$ in (1.1) are given in parametric form ($\mathbb{R}^2$) as in Headrick [2, equations (2.12) and (2.13)]

$$f_{p(W)}(p(w)) = f_{p(W)}(p(x, y)) = f_{p(W)}\left(p(w), \frac{f_W(w)}{p'(w)}\right), \quad (1.3)$$

$$F_{p(W)}(p(w)) = F_{p(W)}(p(x, y)) = F_{p(W)}(p(w), F_W(w)), \quad (1.4)$$

where the derivative $p'(w) > 0$ in (1.3), that is, the polynomial p is a strictly increasing monotonic function and where $f_W(w)$ and $F_W(w)$ are the pdf and cdf associated with the random variable W in (1.2). To demonstrate, Figure 1 gives the graphs of a pdf and cdf associated with a standard normal-based power method distribution used in a Monte Carlo study by Berkovits et al. [8] (and, similarly, Enders [21] and Olsson et al. [24]).

The graphs in Figure 1 and numerical solutions for the coefficients c_i , which are based on the shape parameters in Table 1, were obtained using (1.3), (1.4), and the software package developed by Headrick et al. [26]. Note that the parameters in Table 1, γ_3 (skew), γ_4 (kurtosis), γ_5 , and γ_6 , are standardized cumulants and are scaled such that the normal distribution would have values of $\gamma_3 = \gamma_4 = \gamma_5 = \gamma_6 = 0$.

Conventional moment-based estimators, such as $g_3, \dots, g_6$ in Table 1, have unfavorable attributes to the extent that they can be substantially biased, have high variance, or can be influenced by outliers and thus may not be representative of the true population parameters

(e.g., see [27, 28]). Some of these attributes are exemplified in Table 1 as the estimates of $g_3, \dots, g_6$ and their respective bootstrap confidence intervals attenuate below the population parameters of $\gamma_3, \dots, \gamma_6$ with increased bias and variance as the order of the estimate increases. That is, on average, the estimates of $g_3, \dots, g_6$ are only 83.67%, 63.81%, 38.35%, and 21.38% of their associated population parameters, respectively.

The estimates of $g_3, \dots, g_6$ in Table 1 were calculated based on samples of size $n = 200$ and Fisher's k -statistics (see, e.g., [29, pages 299-300]), that is, the formulae currently used by most commercial software packages such as SAS, SPSS (PASW), Minitab, and so forth, for computing values of skew and kurtosis. Thus, it should also be pointed out that these estimates ($g_3, \dots, g_6$) have another undesirable property of being algebraically bounded based on the sample size, that is, $|g_3| \leq n^{1/2}$, $g_4 \leq n$, $|g_5| \leq n^{3/2}$, $g_6 \leq n^2$. As such, if a researcher was using a value of kurtosis in a Monte Carlo study, such as $\gamma_4 = 21$ in Table 1, and drawing samples of size $n = 15$, then the largest sample estimate of kurtosis possible is $g_4 = 15$ or 71.43% of the parameter.

The method of L -moments is an attractive alternative to conventional method of moments in terms of describing theoretical or empirical probability distributions, estimating parameters, and hypothesis testing (see [27, 28, 30]). More specifically, the first four L -moments are analogous to conventional moments as they describe the location, scale, skew, and kurtosis of a data set. However, L -moments have demonstrated to be superior to conventional moments to the extent that they (a) exist whenever the mean of the distribution exists, (b) only require that a distribution have finite variance for their standard errors to be finite, (c) are nearly unbiased for all sample sizes and distributions, (d) do not suffer from the deleterious effects of sampling variability, (e) are more robust in the presence of outliers, and (f) are not algebraically bounded based on sample size (see [27, 28, 30-33]). Further, it has been demonstrated that there are conditions where the method of L -moments can also yield more accurate and efficient parameter estimates than the method of maximum likelihood when sample sizes are small to moderate (see [27, 34-37]). Other advances have also been made. For example, Elamir and Seheult (see [38, 39]) introduced trimmed L -moments and derived expressions for the exact variances and covariances of sample L -moments. Further, Necir and Meraghni [40] demonstrated that L -moments and trimmed L -moments are useful for estimating L -functionals in the context of heavy-tailed distributions.

Estimates of L -moments and L -moment ratios are based on linear combinations of order statistics unlike conventional moments that are based on raising the data to successive powers which in turn gives greater weight to data points located farther away from the mean and thus may result with estimates (g_j) having substantial bias and (or) variance. For example, the L -moment ratio estimates (t_j) in Table 2 are relatively much closer to their respective population parameters (τ_j) with smaller variance than their corresponding conventional moment-based analogs (g_j) in Table 1. Specifically, the ratios (t_j) are, on average, 99.03%, 99.55%, 98.35%, and 99.29% of their respective parameters.

In view of the above, the present aim is to obviate the problems associated with conventional moments in the context of power method transformations of the form in (1.1) by characterizing these transformations through L -moments. The focus is on standard normal and standard logistic-based polynomials. In Section 2, the essential requisite information associated with L -moments for theoretical and empirical distributions is provided as well as the derivations for the systems of equations for computing polynomial coefficients. Closed-formed formulae are provided for determining if any particular polynomial has a valid pdf. Further, the boundary regions for valid pdfs are derived and graphed for polynomials of order three and for the symmetric case associated with polynomials of order five. In Section 3,




Figure 1: Standard normal-based power method pdf and cdf. The coefficients used in (1.3) and (1.4) are $c_1 = -0.252299$, $c_2 = 0.418610$, $c_3 = 0.252299$, and $c_4 = 0.147593$ and are associated with the parameters in Tables 1 and 2.

Table 1: Conventional moment-based parameters (γ) and their estimates (g) for the pdf in Figure 1. Each estimate (Est.), and bootstrap confidence interval (C.I.) was based on resampling 25,000 statistics. Each sample statistic was based on a sample size of $n = 200$.

Parameter	Est. (95% C.I.)	Standard error
skew: $\gamma_3 = 3$	g_3 : 2.51 (2.50, 2.52)	0.007522
kurtosis: $\gamma_4 = 21$	g_4 : 13.4 (13.2, 13.5)	0.069400
$\gamma_5 = 182$	g_5 : 69.8 (68.4, 71.3)	0.723200
$\gamma_6 = 2329$	g_6 : 498 (485, 514)	7.289000

conventional moments and L -moments are compared in terms of estimation and distribution fitting to demonstrate the superior characteristics that L -moments have in the context of the power method. In Section 4, third- and fifth-order standard normal and logistic-based polynomials are compared in terms of their upper and lower L -moment ratio boundary points.

2. Methodology

2.1. Preliminaries

L -moments are defined as linear combinations of probability weighted moments (PWMs). For a continuous theoretical distribution with a cdf denoted as $F(x)$, the PWMs can be generally defined as in Hosking [27]

$$\beta_r = \int x \{F(x)\}^r dF(x), \quad (2.1)$$

where $r = 0, 1, 2, \dots$. The L -moments can be determined by summing the PWMs as

$$\lambda_{r+1} = \sum_{i=0}^r p_{(r,i)} \beta_i, \quad (2.2)$$

Table 2: L -moment ratios (τ) and their estimates (t) for the pdf in Figure 1. Each estimate and bootstrap confidence interval (C.I.) was based on resampling 25,000 statistics. Each sample statistic was based on a sample size of $n = 200$.

Parameter	Est. (95% C.I.)	Standard error
L -skew: $\tau_3 = .313041$	$t_3: 0.310 (.309, .311)$	0.000407
L -kurtosis: $\tau_4 = .333498$	$t_4: 0.332 (.332, .333)$	0.000268
$\tau_5 = .0986303$	$t_5: 0.097 (.097, .098)$	0.000320
$\tau_6 = .157110$	$t_6: 0.156 (.156, .157)$	0.000252

where

$$p_{(r,i)} = \frac{(-1)^{r-i}(r+i)!}{(i!)^2(r-i)!} \quad (2.3)$$

are coefficients from shifted orthogonal Legendre polynomials. Specifically, the first six L -moments based on (2.1)–(2.3) are expressed as

$$\begin{aligned}
 \lambda_1 &= \beta_0, \\
 \lambda_2 &= 2\beta_1 - \beta_0, \\
 \lambda_3 &= 6\beta_2 - 6\beta_1 + \beta_0, \\
 \lambda_4 &= 20\beta_3 - 30\beta_2 + 12\beta_1 - \beta_0, \\
 \lambda_5 &= 70\beta_4 - 140\beta_3 + 90\beta_2 - 20\beta_1 + \beta_0, \\
 \lambda_6 &= 252\beta_5 - 630\beta_4 + 560\beta_3 - 210\beta_2 + 30\beta_1 - \beta_0.
 \end{aligned} \quad (2.4)$$

Analogous to conventional moment theory, the values of λ_1 and λ_2 are parameters associated with the location and scale of the distribution. More specifically, the L -mean (λ_1) is the usual arithmetic mean, and L -scale ($\lambda_2 \geq 0$) is one-half of Gini's coefficient of mean difference (see, e.g., [29, pages 47-48]). Higher-order L -moments are transformed to dimensionless indices referred to as L -moment ratios defined as $\tau_{r+1} = \lambda_{r+1}/\lambda_2$ for $r \geq 2$. In general, L -moment ratios are bounded such that $|\tau_{r+1}| < 1$ as is the index of L -skew (τ_3) where a symmetric distribution implies that $\tau_{2r-1} = 0$. Smaller boundaries can be found for specific cases. For example, in the context of continuous distributions, L -kurtosis (τ_4) and τ_6 have boundaries of (see Jones [41])

$$\begin{aligned}
 \frac{5\tau_3^2 - 1}{4} &< \tau_4 < 1, \\
 \frac{1}{25}(42\tau_4^2 - 14\tau_4 - 3) &< \tau_6 < 1,
 \end{aligned} \quad (2.5)$$

Table 3: L -moment parameters (λ, τ) and percentiles for the Beta $(\alpha = 5, \beta = 4)$ distribution and power method approximations in Figure 3.

Parameters	%	Beta (Exact)	Percentiles	
			Third order	Fifth order
$\lambda_1 = 0.555556$	1	0.1982	0.1679	0.1983
$\lambda_2 = 0.0895836$	5	0.2892	0.2874	0.2892
$\tau_3 = -.0251716$	25	0.4445	0.4507	0.4445
$\tau_4 = .0921653$	75	0.6709	0.6648	0.6709
$\tau_5 = -.0066496$	95	0.8071	0.8097	0.8071
$\tau_6 = .0284846$	99	0.8791	0.9072	0.8789

which indicate that τ_4 and τ_6 have lower bounds of $-1/4$ and $-1/6$, respectively. An example of a set of computed L -moments (λ_1, λ_2) and L -moment ratios $(\tau_3, \dots, \tau_6)$ based on (2.1) through (2.4) is provided in the first column of Table 3 for a Beta distribution.

Empirical L -moments for a set of data of size n are linear combinations of the sample order statistics $X_{1:n} \leq X_{2:n} \leq \dots \leq X_{n:n}$. The unbiased estimates of the PWMs are

$$b_r = \frac{1}{n} \sum_{i=r+1}^n \frac{(i-1)(i-2) \cdots (i-r)}{(n-1)(n-2) \cdots (n-r)} X_{i:n} \quad (2.6)$$

for $r = 0, 1, 2, \dots$ and where b_0 is the sample mean. The sample L -moments are obtained by substituting b_r in place of β_r in (2.4). The notations used for sample L -moments and L -moment ratios are $t_{r+1} = \ell_{r+1}/\ell_2$ for $r \geq 2$.

2.2. L -Moments for Standard Normal Polynomial Transformations

Using (1.1)-(1.2) and (2.1), the PWMs for power method polynomials based on the standard normal distribution are

$$\beta_r = \int_{-\infty}^{\infty} \sum_{i=1}^m c_i z^{i-1} \{\Phi(z)\}^r \phi(z) dz. \quad (2.7)$$

Integrating (2.7) for $\beta_{r=0,\dots,5}$ and subsequently substituting these PWMs into (2.4) (and after several tedious manipulations) and simplifying yields the following system of equations for fifth-order polynomials ($m = 6$):

$$\lambda_1 = c_1 + c_3 + 3c_5, \quad (2.8)$$

$$\lambda_2 = \frac{\Delta}{2} = \frac{(4c_2 + 10c_4 + 43c_6)/(2\sqrt{\pi})}{2}, \quad (2.9)$$

$$\tau_3 = \frac{6c_3 + 26c_5}{\sqrt{3}\Delta\pi}, \quad (2.10)$$

$$\tau_4 = \frac{20\sqrt{2}(\delta_1c_2 + \delta_2c_4 + \delta_3c_6)}{\Delta\pi^{3/2}} - \frac{3}{2}, \quad (2.11)$$

$$\tau_5 = \frac{84\sqrt{5}(\delta_4c_3 + \delta_5c_5) - 5\pi\sqrt{3}(3c_3 + 13c_5)}{3\Delta\pi^2}, \quad (2.12)$$

$$\tau_6 = \frac{15\Delta\pi^{5/2} - 560\pi\sqrt{2}(\delta_1c_2 + \delta_2c_4 + \delta_3c_6) + 1344\sqrt{3}(\delta_6c_2 + \delta_7c_4 + \delta_8c_6)}{8\Delta\pi^{5/2}}, \quad (2.13)$$

where $\Delta = (4c_2 + 10c_4 + 43c_6)/(2\sqrt{\pi})$ is the coefficient of mean difference. The analytical expressions for the constants $\delta_{i=1,\dots,8}$ in (2.11)–(2.13) are given in Appendix A.

The derivations above yield a system of six equations (2.8)–(2.13) expressed in terms of six real variables $c_1, \dots, c_6$. The first two equations of this system can be standardized by setting (2.8) to zero and (2.9) to $1/\sqrt{\pi}$. The next four equations, (2.10)–(2.13), are set to the desired values of $\tau_3, \dots, \tau_6$. Simultaneously solving this system of equations yields the solutions of $c_1, \dots, c_6$. The coefficients are then substituted into (1.1) to generate $p(W)$ which has zero mean ($\lambda_1 = 0$), one-half of the coefficient of mean difference for the unit-normal distribution ($\lambda_2 = 1/\sqrt{\pi}$), and the specified values of $\tau_3, \dots, \tau_6$. If the negatives of τ_3 and τ_5 are desired, then inspection of (2.8)–(2.13) indicates that all that is needed are simultaneous sign reversals between c_1, c_3 , and c_5 . These sign reversals will have no effect on τ_2, τ_4 , or τ_6 . It is worthy to point out that it is not necessary to numerically solve the system of equations in (2.8)–(2.13) as the coefficients $c_1, \dots, c_6$ have unique solutions which can be determined by evaluating equations (A.9)–(A.14) in Appendix A. See Figures 2(a) and 2(b) for examples of standard normal-based fifth-order power method pdfs.

Setting $c_5 = c_6 = 0$ in (2.8)–(2.11) and simplifying yields the system of equations for the smaller class of distributions associated with third-order polynomials ($m = 4$) in (1.1) as

$$\lambda_1 = c_1 + c_3, \quad (2.14)$$

$$\lambda_2 = \frac{\Delta}{2} = \frac{(4c_2 + 10c_4)/(2\sqrt{\pi})}{2}, \quad (2.15)$$

$$\tau_3 = \frac{2\sqrt{3}c_3}{\Delta\pi}, \quad (2.16)$$

$$\tau_4 = \frac{20\sqrt{2}(\delta_1c_2 + \delta_2c_4)}{\Delta\pi^{3/2}} - \frac{3}{2}, \quad (2.17)$$

where the solutions for the coefficients in (2.14)–(2.17) are

$$\begin{aligned} c_1 &= -c_3 = -\tau_3 \sqrt{\frac{\pi}{3}}, \\ c_2 &= \frac{-16\delta_2 + \sqrt{2}(3 + 2\tau_4)\pi}{8(5\delta_1 - 2\delta_2)}, \\ c_4 &= \frac{40\delta_1 - \sqrt{2}(3 + 2\tau_4)\pi}{20(5\delta_1 - 2\delta_2)}. \end{aligned} \quad (2.18)$$

Examples of a fifth-order $(c_1, \dots, c_6)$ and a third-order $(c_1, \dots, c_4)$ power method distribution are given in Figure 3, both of which provide an approximation of the Beta $(\alpha = 5, \beta = 4)$ distribution. The coefficients associated with the two polynomial approximations in Figure 3 were used in (1.4) to compute the percentiles in Table 3 for the power method pdfs. Inspection of the graphs in Figure 3 and the percentiles in Table 3 indicate that the fifth-order polynomial provides a much more accurate approximation of the Beta distribution than the third-order polynomial. The reason for this is because the fifth-order system could produce a valid pdf that is based on an exact match with the Beta distribution's L -moment ratios $\tau_3, \dots, \tau_6$, whereas the third-order system was unable to produce a valid pdf that has an exact match with τ_4, τ_5 , and τ_6 . Thus, it is important to consider the boundary conditions for valid power method pdfs.

2.3. *L-Moment Boundaries for Standard Normal Polynomial Transformations*

The restriction that $p'(w) > 0$ in (1.3) implies that a set of solved coefficients may not necessarily produce a valid pdf. To determine if a third-order polynomial produces a valid pdf, we first set the quadratic equation $p'(w) = 0$ and subsequently solve for w as

$$w = \frac{-c_3 \pm (c_3^2 - 3c_2c_4)^{1/2}}{3c_4}. \quad (2.19)$$

A set of solved coefficients will produce a valid pdf if the discriminant $c_3^2 - 3c_2c_4$ in (2.19) is negative. That is, the complex solutions for w must have nonzero imaginary parts. As such, setting $c_3^2 = 3c_2c_4$ will yield the point where the discriminant vanishes and thus real-valued solutions exist to $p'(w) = 0$.

Standardizing (2.15), by setting the coefficient of mean difference to $1/\sqrt{\pi}$, and solving for c_4 give

$$c_4 = \frac{2}{5} - \frac{2c_2}{5}. \quad (2.20)$$




Figure 2: Standard normal and standard logistic-based power method pdfs.

Substituting the right-hand side of (2.20) into (2.16) and (2.17) and setting $c_3 = (3c_2c_4)^{1/2}$, because we only need to consider positive values of L -skew, gives

$$\tau_3 = 3\sqrt{\frac{2}{5\pi}}\sqrt{c_2(1-c_2)}, \quad (2.21)$$

$$\tau_4 = \frac{2\sqrt{2}(5\delta_1c_2 + 2\delta_2 - 2\delta_2c_2)}{\pi} - \frac{3}{2}. \quad (2.22)$$




Figure 3: Third- and fifth-order standard normal-based polynomial approximations (dashed lines) of a Beta ($\alpha = 5$, $\beta = 4$) distribution. The L -moment ratios for this Beta distribution are given in Table 3. The fifth-order polynomial coefficients are $c_1 = .02697$, $c_2 = 1.071$, $c_3 = -.0297$, $c_4 = -.0312$, $c_5 = .0009$, and $c_6 = .00065$. The third-order polynomial coefficients are $c_1 = .02576$, $c_2 = 0.9993$, $c_3 = -.02576$, and $c_4 = .00027$.




Figure 4: Boundary region and values of L -skew $|\tau_3|$ and L -kurtosis τ_4 for third-order standard normal power method pdfs. For asymmetric distributions, valid pdfs exist in the region inside the graphed boundary. For symmetric distributions, valid pdfs exist for τ_4 between $a_4 < \tau_4 < b_4$. The lower boundary point $(a_{3,4})$ denotes the normal distribution. Note that a_3 and a_4 are lower limits and b_3 and b_4 are upper limits.

Inspection of (2.21) indicates that for real values of τ_3 to exist then we must have $c_2 \in [0, 1]$ and thus from (2.20) $c_4 \in [0, 0.4]$. Using (2.21) and (2.22), the graph of the region for valid third-order power method pdfs is given in Figure 4 along with the minimum and maximum values of τ_3 and τ_4 . In summary, a valid standardized nonnormal third-order pdf will have the properties of (a) $0 < c_2 < 1$, (b) $0 < c_4 < 2/5$, and (c) $c_3^2 - 3c_2c_4 < 0$.




Figure 5: Boundary region and values of L -kurtosis τ_4 and τ_6 for symmetric fifth-order standard normal-based power method pdfs. Valid pdfs exist inside the elliptical graph. The point at the origin represents the regular uniform distribution ($\tau_4 = \tau_6 = 0$). Note that a_4 and a_6 are lower limits and b_4 and b_6 are upper limits.

In terms of fifth-order polynomials, the formulae that solve the (quartic) equation $p'(w) = 0$ must also have nonzero imaginary parts to ensure a valid pdf. Specifically, the closed-formed expressions for evaluating pdfs associated with symmetric distributions are

$$w = \pm \left[\frac{\pm(9c_4^2 - 20c_2c_6)^{1/2} - 3c_4}{10c_6} \right]^{1/2}, \quad (2.23)$$

$$w = \mp \left[\frac{\pm(9c_4^2 - 20c_2c_6)^{1/2} - 3c_4}{10c_6} \right]^{1/2},$$

and for asymmetric distributions the expressions are

$$w = \pm \frac{1}{2}A^{1/2} \pm \frac{1}{2}(B \pm C)^{1/2} - \frac{c_5}{5c_6}, \quad (2.24)$$

$$w = \pm \frac{1}{2}A^{1/2} \mp \frac{1}{2}(B \pm C)^{1/2} - \frac{c_5}{5c_6}.$$

The expressions for A , B , and C in (2.24) are given in Appendix B.

The boundary for the larger class of symmetric fifth-order pdfs (i.e., $\tau_3 = \tau_5 = 0$; $c_1 = c_3 = c_5 = 0$) can be viewed by making use of (2.23). Specifically, the graph of the region for valid symmetric pdfs is given in Figure 5 along with the minimum and maximum values of

τ_4 and τ_6 . The elliptical graph in Figure 5 consists of four separate segments where τ_4 in (2.11) and τ_6 in (2.13) are expressed solely as functions of c_2 . See Appendix C for further details.

2.4. L-Moments and Boundaries for Standard Logistic Polynomial Transformations

The method for deriving the system of equations for power method transformations based on the standard logistic distribution is similar to the derivation of the system associated with (2.8)–(2.13). However, the derivation is more straight forward because the cdf for the logistic distribution is available in closed form. As such, the system of equations for fifth-order polynomials in (1.1) is

$$\lambda_1 = c_1 + c_3 + \frac{21c_5}{5}, \quad (2.25)$$

$$\lambda_2 = \frac{\Delta}{2} = \frac{2\sqrt{3}(c_2 + 3c_4 + 21c_6)/\pi}{2}, \quad (2.26)$$

$$\tau_3 = \frac{12c_3 + 72c_5}{\Delta\pi^2}, \quad (2.27)$$

$$\tau_4 = \frac{2\sqrt{3}(90c_4 + 900c_6 + \pi^2(c_2 + 3c_4 + 21c_6))}{6\Delta\pi^3}, \quad (2.28)$$

$$\tau_5 = \frac{5\pi^2c_3 + 252c_5 + 30\pi^2c_5}{\Delta\pi^4}, \quad (2.29)$$

$$\tau_6 = \frac{2\Delta\pi^5 + 2\sqrt{3}(315\pi^2c_4 + 11340c_6 + 3150\pi^2c_6)}{30\Delta\pi^5}, \quad (2.30)$$

where $\Delta = 2\sqrt{3}(c_2 + 3c_4 + 21c_6)/\pi$ is the coefficient of mean difference and for standardized distributions the solutions for the coefficients $c_1, \dots, c_6$ are given in Appendix D. The system of equations for the smaller family of distributions based on third-order polynomials can be obtained by setting $c_5 = c_6 = 0$ in (2.25)–(2.28), and for standardized distributions, the solutions for the coefficients are

$$\begin{aligned} c_1 &= -c_3 = -\frac{\tau_3\pi}{2\sqrt{3}}, \\ c_2 &= \frac{30 + \pi^2(1 - 6\tau_4)}{30}, \\ c_4 &= \frac{\pi^2(6\tau_4 - 1)}{90}. \end{aligned} \quad (2.31)$$

The derivation of the L -skew (τ_3) and L -kurtosis (τ_4) boundary for third-order polynomials is also analogous to the steps taken in Section 2.3. As such, the graph of the boundary for valid third-order power method pdfs is given in Figure 6 along with the minimum and maximum values of τ_3 and τ_4 . Similarly, a valid standardized nonlogistic third-order pdf will have the properties of (a) $0 < c_2 < 1$, (b) $0 < c_4 < 1/3$, and (c) $c_3^2 - 3c_2c_4 < 0$.




Figure 6: Boundary region and values of L -skew $|\tau_3|$ and L -kurtosis τ_4 for third-order standard logistic power method pdfs. For asymmetric distributions, valid pdfs exist in the region inside the graphed boundary. For symmetric distributions, valid pdfs exist for τ_4 between $a_4 < \tau_4 < b_4$. The lower boundary point (a_3, a_4) represents the standard logistic distribution. Note that a_3 and a_4 are lower limits and b_3 and b_4 are upper limits.

In terms of fifth-order polynomials, the boundary region for symmetric pdfs and minimum and maximum values of τ_4 and τ_6 are given in Figure 7. The elliptical graph in Figure 7 also consists of four segments. See Appendix E for further details. Note that the expressions in (2.23) and (2.24) are also used to determine whether fifth-order pdfs are valid or not. An example of a logistic-based power method pdf is given in Figure 2(c).

3. A Comparison of Conventional Moments and L -Moments

3.1. Estimation

One of the advantages that sample L -moment ratios $(t_{3,\dots,6})$ have over conventional moment-based estimates, such as skew (g_3) and kurtosis (g_4) , is that $t_{3,\dots,6}$ are less biased (e.g., [28]). This advantage can be demonstrated in the context of the power method by considering the simulation results associated with the indices for the percentage of relative bias (RBias%) and standard error (SE) reported in Table 4.

Specifically, a Fortran algorithm was coded to generate twenty-five thousand independent samples of size $n = 50, 100, 1000$, and the estimates of $g_{3,\dots,6}$ and $t_{3,\dots,6}$ were computed for each of the (3×25000) samples based on the parameters and coefficients listed in Table 4. The estimates of $g_{3,\dots,6}$ were computed based on Fisher's k -statistics, and the estimates of $t_{3,\dots,6}$ were based on (2.6). Bootstrapped average estimates, confidence intervals, and SEs were obtained for $g_{3,\dots,6}$ and $t_{3,\dots,6}$ using twenty-five thousand resamples via the commercial software package Spotfire S+ [42]. The percentage of relative bias for each estimate was computed as $\text{RBias}\%, (g_j) = 100 \times (g_j - \gamma_j) / \gamma_j$ and $\text{RBias}\%, (t_j) = 100 \times (t_j - \tau_j) / \tau_j$.




Figure 7: Boundary region and values of L -kurtosis τ_4 and τ_6 for symmetric fifth-order standard logistic-based power method pdfs. Valid pdfs exist inside the elliptical graph. Note that a_4 and a_6 are lower limits and b_4 and b_6 are upper limits.

The results in Table 4 demonstrate the substantial advantage that L -moment ratios have over conventional estimates in terms of both bias and error. These advantages are most pronounced in the context of smaller sample sizes and higher-order moments. For example, given a sample size of $n = 50$, the conventional estimates of g_3 (g_5) and g_4 (g_6) generated in the simulation were, on average, 11.40% (56.50%) and 31.93% (78.36%) less than their respective parameters. On the other hand, the amounts of relative bias associated with the L -moment ratios were essentially negligible. Further, the SEs associated with $t_{3,\dots,6}$ are relatively much smaller than the corresponding SEs for $g_{3,\dots,6}$.

3.2. Distribution Fitting

Presented in Figure 8 are conventional moment and L -moment-based power method pdfs superimposed on a histogram of data from the Project Match Research Group studies (see [43, 44]). The data are associated with participants who were assigned to a Twelve-Step Facilitation (TSF) treatment condition for alcoholism. Specifically, these data are the total number of drinks per ninety-day period which were determined by each participant's reports of daily standard ethanol drinks (one drink was equal to one-half of an ounce). Drinking was assessed approximately every three months, beginning at pretreatment (baseline), immediately following treatment, and at three-month intervals over a twelve-month posttreatment follow-up period (i.e., six, nine, twelve, and fifteen months posttreatment).

The sample estimates ($g_{3,\dots,6}; t_{3,\dots,6}$) associated with Figure 8 were based on a sample size of $n = 311$ participants. The estimates were also used to solve for the two sets of coefficients, which produced the power method pdfs based on (1.3). Note that the two polynomials were linearly transformed using the location and scale estimates $(m, s; \ell_1, \ell_2)$ from the data. Visual inspection of the approximations in Figure 8 and the goodness of fit statistics given in Table 5

Table 4: Overall average estimates (Est.), bootstrap confidence intervals (C.I.), standard errors (SE), and percentage of relative bias (RBias%) for the conventional moment-based statistics (g) and L -moment ratios (t).

(a)					
Parameters ($\chi^2_{df=8}$) and coefficients					
$\gamma_1 = 0$	$c_1 = -0.163968$	$\lambda_1 = 0$	$c_1 = -0.169160$		
$\gamma_2 = 1$	$c_2 = 0.950794$	$\lambda_2 = 1/\sqrt{\pi}$	$c_2 = 0.980897$		
$\gamma_3 = 1$	$c_3 = 0.165391$	$\tau_3 = 0.164666$	$c_3 = 0.170627$		
$\gamma_4 = 1.5$	$c_4 = 0.007345$	$\tau_4 = 0.131237$	$c_4 = 0.007577$		
$\gamma_5 = 3$	$c_5 = -0.000474$	$\tau_5 = 0.051194$	$c_5 = -0.000489$		
$\gamma_6 = 7.5$	$c_6 = 0.000014$	$\tau_6 = 0.048334$	$c_6 = 0.000015$		
(b)					
Est. (95% C.I.)	SE	RBias%	Est. (95% C.I.)	SE	RBias%
$n = 50$					
$g_3: .8860 (.8808, .8917)$	.0028	-11.40	$t_3: .1626 (.1618, .1634)$	.0004	-1.25
$g_4: 1.021 (.9975, 1.043)$	.0116	-31.93	$t_4: .1314 (.1308, .1321)$	.0003	0.12
$g_5: 1.302 (1.218, 1.389)$	.0439	-56.60	$t_5: .0510 (.0505, .0516)$	.0003	-0.38
$g_6: 1.623 (1.313, 1.973)$	.1673	-78.36	$t_6: .0484 (.0480, .0489)$	.0002	0.14
$n = 100$					
$g_3: .9395 (.9350, .9436)$	.0022	-6.05	$t_3: .1635 (.1629, .1640)$	.0003	-0.71
$g_4: 1.225 (1.204, 1.245)$	.0104	-18.33	$t_4: .1314 (.1309, .1318)$	.0002	0.09
$g_5: 1.930 (1.847, 2.032)$	.0466	-35.67	$t_5: .0511 (.0507, .0514)$	.0002	-0.18
$g_6: 3.385 (3.004, 3.862)$	.2160	-54.87	$t_6: .0483 (.0479, .0486)$	.0002	-0.07
$n = 1000$					
$g_3: .9941 (.9926, .9957)$	.0008	-0.59	$t_3: .1645 (.1643, .1647)$	.0001	-0.10
$g_4: 1.471 (1.462, 1.480)$	.0046	-1.93	$t_4: .1313 (.1311, .1314)$	.0001	0.05
$g_5: 2.869 (2.822, 2.921)$	.0252	-4.37	$t_5: .0512 (.0510, .0512)$	.0001	0.01
$g_6: 6.889 (6.586, 7.186)$	.1544	-8.15	$t_6: .0483 (.0482, .0484)$	.0001	-0.01

indicate that the L -moment pdf provides a more accurate fit to the actual data. The reason for this is partially attributed to the fact that the conventional moment-based power method pdf does not have an exact match with g_6 , whereas the L -moment pdf is based on an exact match with all of the sample estimates. Note also that the asymptotic P values in Table 5 are based on a chi-square distribution with degrees of freedom: $df = 13(\text{classes}) - 6(\text{estimates}) - 1(\text{sample size}) = 6$.

4. Discussion and Conclusion

This paper introduced L -moments in the context of standard normal and standard logistic-based power method polynomials. The boundaries were analytically derived for polynomials of order three and for symmetric pdfs associated with polynomials of order five. The lower boundary point associated with the standard normal third-order system is the standard normal distribution (i.e., $\tau_4 \approx 0.1226$ and $\tau_6 \approx 0.0437$). As such, it is worthy to point out that the larger family of pdfs for the standard normal fifth-order system includes the L -moment




Figure 8: Histograms and fifth-order power method approximations for the TSF group. ^aThe value of g_6 had to be increased to -0.28898 to ensure a valid pdf.

Table 5: Chi-square goodness of fit statistics for the conventional (C) moment and L -moment approximations for the TSF group data ($n = 311$) in Figure 8.

%	Expected	Obs (C)	Obs (L)	Total number of drinks (C)	Total number of drinks (L)
10	31.10	38	32	<207.51	<176.05
20	31.10	26	32	207.51–286.14	176.05–287.51
30	31.10	22	25	286.14–380.05	287.51–397.76
40	31.10	31	36	380.05–490.40	397.76–514.61
50	31.10	37	33	490.40–620.50	514.61–644.07
60	31.10	29	25	620.50–777.13	644.07–794.99
70	31.10	33	35	777.13–973.41	794.99–977.85
75	15.55	17	15	973.41–1094.27	977.85–1090.33
80	15.55	18	16	1094.27–1238.41	1090.33–1224.72
85	15.55	15	16	1238.41–1418.18	1224.72–1393.72
90	15.55	16	15	1418.18–1660.33	1393.72–1625.71
95	15.55	15	16	1660.33–2045.66	1625.71–2010.65
100	15.55	14	15	2045.66 or more	2010.65 or more
				C: $\chi^2 = 7.135$	L: $\chi^2 = 3.920$
				$P = .309$	$P = .688$

ratios associated with the regular uniform or Beta ($\alpha = 1, \beta = 1$) distribution (i.e., $\tau_4 = 0$ and $\tau_6 = 0$). Further, the upper boundary of L -kurtosis for the third-order system of $\tau_4 \approx .5728$ extends up to a boundary of $\tau_4 \approx .8233$ for the fifth-order system which is a substantial increase. That is, in terms of conventional moments, this is equivalent to increasing kurtosis from $\gamma_4 \approx 43$ to $\gamma_4 \approx 833$.

In terms of the third-order logistic-based family of pdfs, it is noted that maximum L -skew is $\tau_3 \approx .5513$ whereas the maximum value of L -skew for the standard normal-based system is $\tau_3 \approx .5352$. Again, this is actually a considerable difference—as these values correspond to values of conventional skew of $\gamma_3 \approx 8.4$ and $\gamma_3 \approx 4$, respectively. Similarly, the maximum values of L -kurtosis for the standard logistic-based and normal-based systems are $\tau_4 \approx .6733$ and $\tau_4 \approx .5728$ and are equivalent to values of conventional kurtosis of $\gamma_4 \approx 472$ and $\gamma_4 \approx 43$, respectively.

Finally, we note that Mathematica ([45], Version 7.0) software for computing polynomial coefficients, cumulative probabilities, and fitting (graphing) power method pdfs to data are available on request.

Appendices

A. Analytical Expressions and Polynomial Coefficients (Normal)

The expressions for $\delta_{i=1,\dots,8}$ in (2.11)–(2.13) and the coefficients $c_{i=1,\dots,6}$ in (2.8)–(2.13) are

$$\delta_1 = \frac{3 \tan^{-1}[\sqrt{2}]}{\sqrt{2}} - \frac{3\pi}{4\sqrt{2}} = 0.360451474758732546420622\dots, \quad (\text{A.1})$$

$$\delta_2 = \frac{15 \tan^{-1}[\sqrt{2}]}{2\sqrt{2}} - \frac{15\pi}{8\sqrt{2}} + \frac{1}{4} = 1.151128686896831366051556\dots, \quad (\text{A.2})$$

$$\delta_3 = \frac{129 \tan^{-1}[\sqrt{2}]}{4\sqrt{2}} - \frac{129\pi}{16\sqrt{2}} + \frac{77}{48} = 5.479020020323041540688357\dots, \quad (\text{A.3})$$

$$\delta_4 = \sqrt{15} \tan^{-1}\left[\sqrt{\frac{5}{3}}\right] - \frac{\pi\sqrt{15}}{4} = 0.489313210040355820119964\dots, \quad (\text{A.4})$$

$$\begin{aligned} \delta_5 &= -\sqrt{15} \tan^{-1}\left[\sqrt{\frac{5}{3}}\right] - 8\sqrt{\frac{5}{3}} \tan^{-1}[\sqrt{15}] + \frac{19\pi}{4}\sqrt{\frac{5}{3}} + \frac{1}{4} \\ &= 2.370357243508208553853177\dots, \end{aligned} \quad (\text{A.5})$$

$$\begin{aligned} \delta_6 &= \frac{5\pi \tan^{-1}[\sqrt{2}]}{4\sqrt{3}} + \frac{5}{8}\sqrt{3} \sin^{-1}\left[\frac{1}{3}\right]^2 - \frac{5\pi \sin^{-1}[1/3]}{8\sqrt{3}} - \frac{5\pi^2}{16\sqrt{3}} + \frac{5\sqrt{3}}{2}\varepsilon_1 \\ &= 0.261016893769534900409394\dots, \end{aligned} \quad (\text{A.6})$$

$$\begin{aligned} \delta_7 &= \frac{25\pi \tan^{-1}[\sqrt{2}]}{8\sqrt{3}} + \frac{25}{4}\sqrt{3} \sin^{-1}\left[\frac{1}{\sqrt{3}}\right]^2 - \frac{25\pi \sin^{-1}[1/\sqrt{3}]}{4\sqrt{3}} - \frac{25\pi^2}{64\sqrt{3}} + \frac{5\pi}{8\sqrt{6}} - \frac{5}{4\sqrt{6}} \\ &\quad + \frac{25\sqrt{3}}{4}\varepsilon_2 = 0.960807335426376575971986\dots, \end{aligned} \quad (\text{A.7})$$

$$\delta_8 = \frac{215\pi \tan^{-1}[\sqrt{2}]}{16\sqrt{3}} + \frac{215}{8}\sqrt{3} \sin^{-1}\left[\frac{1}{\sqrt{3}}\right]^2 - \frac{215\pi \sin^{-1}[1/\sqrt{3}]}{8\sqrt{3}} - \frac{215\pi^2}{128\sqrt{3}} + \frac{385\pi}{96\sqrt{6}} - \frac{385}{48\sqrt{6}} + \frac{215\sqrt{3}}{8}\varepsilon_3 = 4.908966006122127514487203\dots, \quad (\text{A.8})$$

$$c_2 = \frac{40320\sqrt{2}(\delta_3\delta_7 - \delta_2\delta_8) - 504(43\delta_7 - 10\delta_8)(3 + 2\tau_4)\pi + 5\sqrt{6}(43\delta_2 - 10\delta_3)\pi^2(27 + 28\tau_4 + 8\tau_6)}{10080\sqrt{2}(43\delta_2\delta_6 - 10\delta_3\delta_6 - 43\delta_1\delta_7 + 4\delta_3\delta_7 + 10\delta_1\delta_8 - 4\delta_2\delta_8)}, \quad (\text{A.9})$$

$$c_3 = \frac{-140\tau_3\delta_5\sqrt{3\pi} + 13\pi^{3/2}\sqrt{5}(2\tau_5 + 5\tau_3)}{140(13\delta_4 - 3\delta_5)}, \quad (\text{A.10})$$

$$c_4 = \frac{-40320\sqrt{2}(\delta_3\delta_6 - \delta_1\delta_8) + 504(43\delta_6 - 4\delta_8)(3 + 2\tau_4)\pi + 5\sqrt{6}(-43\delta_1 + 4\delta_3)\pi^2(27 + 28\tau_4 + 8\tau_6)}{10080\sqrt{2}(43\delta_2\delta_6 - 10\delta_3\delta_6 - 43\delta_1\delta_7 + 4\delta_3\delta_7 + 10\delta_1\delta_8 - 4\delta_2\delta_8)}, \quad (\text{A.11})$$

$$c_5 = \frac{28\tau_3\delta_4\sqrt{3\pi} - (3\pi^{3/2}(2\tau_5 + 5\tau_3))/\sqrt{5}}{364\delta_4 - 84\delta_5}, \quad (\text{A.12})$$

$$c_6 = \frac{20160(\delta_2\delta_6 - \delta_1\delta_7) - 252\sqrt{2}(5\delta_6 - 2\delta_7)(3 + 2\tau_4)\pi + 5\sqrt{3}(5\delta_1 - 2\delta_2)\pi^2(27 + 28\tau_4 + 8\tau_6)}{5040(43\delta_2\delta_6 - 10\delta_3\delta_6 - 43\delta_1\delta_7 + 4\delta_3\delta_7 + 10\delta_1\delta_8 - 4\delta_2\delta_8)}, \quad (\text{A.13})$$

$$c_1 = -c_3 - 3c_5. \quad (\text{A.14})$$

Equations (A.6)–(A.8) were derived by adapting the techniques used by Renner [46] and where ε_1 , ε_2 , and ε_3 are based on the standard recurrence relation for $\int \sec^p u \, du$. The coefficients in (A.9)–(A.14) are for standardized distributions (i.e., $\lambda_1 = 0$, $\lambda_2 = 1/\sqrt{\pi}$).

B. Expressions for (2.24)

The expressions for A , B , and C in (2.24) are

$$\begin{aligned} A &= \frac{4c_5^2}{25c_6^2} + \frac{(E^{1/2} + D)^{1/3}}{15c_6 2^{1/3}} + \frac{2^{1/3}F}{5c_6(E^{1/2} + D)^{1/3}} - \frac{2c_4}{5c_6}, \\ B &= \frac{8c_5^2}{25c_6^2} - \frac{(E^{1/2} + D)^{1/3}}{15c_6 2^{1/3}} - \frac{2^{1/3}F}{5c_6(E^{1/2} + D)^{1/3}} - \frac{4c_4}{5c_6}, \\ C &= A^{-(1/2)} \left(\frac{48c_4c_5}{100c_6^2} - \frac{16c_3}{20c_6} - \frac{64c_5^3}{500c_6^3} \right), \\ D &= 54c_4^3 - 216c_3c_4c_5 + 432c_2c_5^2 + 540c_3^2c_6 - 1080c_2c_4c_6, \end{aligned}$$

$$\begin{aligned}
 E &= D^2 - 4(9c_4^2 - 24c_3c_5 + 60c_2c_6)^3, \\
 F &= 3c_4^2 - 8c_3c_5 + 20c_2c_6.
 \end{aligned}
 \tag{B.1}$$

C. Symmetric Distributions Boundary (Normal)

The elliptical graph in Figure 5 is based on (2.11) and (2.13) and setting $c_4 = (4 - 4c_2 - 43c_6)/10$ from (2.9) and then setting c_6 as follows:

$$\begin{aligned}
 \text{Segment 1: } 0 \leq c_2 \leq 1, \quad c_6 &= -\frac{4(137c_2 - 387)}{16641} + \frac{40\sqrt{5}\sqrt{387c_2 - 262c_2^2}}{16641}, \\
 \text{Segment 2: } 1 \leq c_2 \leq \frac{387}{262}, \quad c_6 &= -\frac{4(137c_2 - 387)}{16641} + \frac{40\sqrt{5}\sqrt{387c_2 - 262c_2^2}}{16641}, \\
 \text{Segment 3: } 0 \leq c_2 \leq 1, \quad c_6 &= -\frac{4(137c_2 - 387)}{16641} - \frac{40\sqrt{5}\sqrt{387c_2 - 262c_2^2}}{16641}, \\
 \text{Segment 4: } 1 \leq c_2 \leq \frac{387}{262}, \quad c_6 &= -\frac{4(137c_2 - 387)}{16641} - \frac{40\sqrt{5}\sqrt{387c_2 - 262c_2^2}}{16641}.
 \end{aligned}
 \tag{C.1}$$

D. Polynomial Coefficients (Logistic)

The coefficients for the standard logistic-based system in (2.25)–(2.30) are

$$\begin{aligned}
 c_2 &= \frac{840 + 28\pi^2(1 - 6\tau_4) + \pi^4(1 - 14\tau_4 + 20\tau_6)}{840}, \\
 c_3 &= \frac{42\tau_3\pi + \pi^3(5\tau_3 - 12\tau_5)}{84\sqrt{3}}, \\
 c_4 &= \frac{\pi^2(42(6\tau_4 - 1) + 5\pi^2(14\tau_4 - 1 - 20\tau_6))}{3780}, \\
 c_5 &= \frac{\pi^3(12\tau_5 - 5\tau_3)}{504\sqrt{3}}, \\
 c_6 &= \frac{\pi^4(1 - 14\tau_4 + 20\tau_6)}{7560}, \\
 c_1 &= -c_3 - \frac{21c_5}{5}.
 \end{aligned}
 \tag{D.1}$$

E. Symmetric Distributions Boundary (Logistic)

The elliptical graph in Figure 7 is based on (2.28) and (2.30) and setting $c_4 = (1 - c_2 - 21c_6)/3$ from (2.26) and then setting c_6 as follows:

$$\begin{aligned}
 \text{Segment 1: } 0 \leq c_2 \leq 1, \quad c_6 &= \frac{21 - 11c_2}{441} + \frac{2\sqrt{5}\sqrt{21c_2 - 16c_2^2}}{441}, \\
 \text{Segment 2: } 1 \leq c_2 \leq \frac{21}{16}, \quad c_6 &= \frac{21 - 11c_2}{441} + \frac{2\sqrt{5}\sqrt{21c_2 - 16c_2^2}}{441}, \\
 \text{Segment 3: } 0 \leq c_2 \leq 1, \quad c_6 &= \frac{21 - 11c_2}{441} - \frac{2\sqrt{5}\sqrt{21c_2 - 16c_2^2}}{441}, \\
 \text{Segment 4: } 1 \leq c_2 \leq \frac{21}{16}, \quad c_6 &= \frac{21 - 11c_2}{441} - \frac{2\sqrt{5}\sqrt{21c_2 - 16c_2^2}}{441}.
 \end{aligned} \tag{E.1}$$

References

- [1] A. I. Fleishman, "A method for simulating non-normal distributions," *Psychometrika*, vol. 43, no. 4, pp. 521–532, 1978.
- [2] T. C. Headrick, *Statistical Simulation: Power Method Polynomials and other Transformations*, Chapman & Hall/CRC, Boca Raton, Fla, USA, 2010.
- [3] T. C. Headrick, "Fast fifth-order polynomial transforms for generating univariate and multivariate nonnormal distributions," *Computational Statistics & Data Analysis*, vol. 40, no. 4, pp. 685–711, 2002.
- [4] T. C. Headrick and S. S. Sawilowsky, "Simulating correlated multivariate nonnormal distributions: extending the Fleishman power method," *Psychometrika*, vol. 64, no. 1, pp. 25–35, 1999.
- [5] K. Høyland, M. Kaut, and S. W. Wallace, "A heuristic for moment-matching scenario generation," *Computational Optimization and Applications*, vol. 24, no. 2-3, pp. 169–185, 2003.
- [6] P. R. Tadikamalla, "On simulating non-normal distributions," *Psychometrika*, vol. 45, no. 2, pp. 273–279, 1980.
- [7] C. D. Vale and V. A. Maurelli, "Simulating multivariate nonnormal distributions," *Psychometrika*, vol. 48, no. 3, pp. 465–471, 1983.
- [8] I. Berkovits, G. R. Hancock, and J. Nevitt, "Bootstrap resampling approaches for repeated measure designs: relative robustness to sphericity and normality violations," *Educational and Psychological Measurement*, vol. 60, no. 6, pp. 877–892, 2000.
- [9] H. J. Keselman, R. R. Wilcox, J. Algina, A. R. Othman, and K. Fradette, "A comparative study of robust tests for spread: asymmetric trimming strategies," *The British Journal of Mathematical and Statistical Psychology*, vol. 61, no. 2, pp. 235–253, 2008.
- [10] L. M. Lix and R. T. Fouladi, "Robust step-down tests for multivariate independent group designs," *British Journal of Mathematical and Statistical Psychology*, vol. 60, no. 2, pp. 245–265, 2007.
- [11] J. Affleck-Graves and B. MacDonald, "Nonnormalities and tests of asset pricing theories," *Journal of Finance*, vol. 44, no. 4, pp. 889–908, 1989.
- [12] G. D. Hess and S. Iwata, "Measuring and comparing business-cycle features," *Journal of Business and Economic Statistics*, vol. 15, no. 4, pp. 432–444, 1997.
- [13] D. Steinley and R. Henson, "OCLUS: an analytic method for generating clusters with known overlap," *Journal of Classification*, vol. 22, no. 2, pp. 221–250, 2005.
- [14] L. Kirisci, T.-c. Hsu, and L. Yu, "Robustness of item parameter estimation programs to assumptions of unidimensionality and normality," *Applied Psychological Measurement*, vol. 25, no. 2, pp. 146–162, 2001.
- [15] M. Harwell, C. A. Stone, T. C. Hsu, and L. Kirisci, "Monte Carlo studies in item response theory," *Applied Psychological Measurement*, vol. 20, no. 2, pp. 101–125, 1996.

- [16] C. A. Stone, "Empirical power and type I error rates for an IRT fit statistic that considers the precision of ability estimates," *Educational and Psychological Measurement*, vol. 63, no. 4, pp. 566–583, 2003.
- [17] J. Benson and J. A. Fleishman, "The robustness of maximum likelihood and distribution-free estimators to non-normality in confirmatory factor analysis," *Quality & Quantity*, vol. 28, no. 2, pp. 117–136, 1994.
- [18] D. B. Flora and P. J. Curran, "An empirical evaluation of alternative methods of estimation for confirmatory factor analysis with ordinal data," *Psychological Methods*, vol. 9, no. 4, pp. 466–491, 2004.
- [19] F. Nasser and J. Wisenbaker, "A Monte Carlo study investigating the impact of item parceling on measures of fit in confirmatory factor analysis," *Educational and Psychological Measurement*, vol. 63, no. 5, pp. 729–757, 2003.
- [20] O. Mahul, "Hedging price risk in the presence of crop yield and revenue insurance," *European Review of Agricultural Economics*, vol. 30, no. 2, pp. 217–239, 2003.
- [21] C. K. Enders, "The impact of nonnormality on full information maximum-likelihood estimation for structural equation models with missing data," *Psychological Methods*, vol. 6, no. 3, pp. 352–370, 2001.
- [22] K.-T. Hau and H. W. Marsh, "The use of item parcels in structural equation modelling: non-normal data and small sample sizes," *The British Journal of Mathematical and Statistical Psychology*, vol. 57, no. 2, pp. 327–351, 2004.
- [23] J. M. Henson, S. P. Reise, and K. H. Kim, "Detecting mixtures from structural model differences using latent variable mixture modeling: a comparison of relative model fit statistics," *Structural Equation Modeling*, vol. 14, no. 2, pp. 202–226, 2007.
- [24] U. H. Olsson, T. Foss, and S. V. Troye, "Does the ADF fit function decrease when the kurtosis increases?" *The British Journal of Mathematical and Statistical Psychology*, vol. 56, no. 2, pp. 289–303, 2003.
- [25] L. Hothorn and W. Lehmacher, "A simple testing procedure "control versus k treatments" for one-sided ordered alternatives, with application in toxicology," *Biometrical Journal*, vol. 33, no. 2, pp. 179–189, 2007.
- [26] T. C. Headrick, Y. Sheng, and F. A. Hodis, "Numerical computing and graphics for the power method transformation using mathematica," *Journal of Statistical Software*, vol. 19, no. 3, pp. 1–17, 2007.
- [27] J. R. M. Hosking, "L-moments: analysis and estimation of distributions using linear combinations of order statistics," *Journal of the Royal Statistical Society. Series B*, vol. 52, no. 1, pp. 105–124, 1990.
- [28] J. R. M. Hosking and J. R. Wallis, *Regional Frequency Analysis: An Approach Based on L-Moments*, Cambridge University Press, Cambridge, UK, 1997.
- [29] M. Kendall and A. Stuart, *The Advanced Theory of Statistics*, Macmillan, New York, NY, USA, 4th edition, 1977.
- [30] J. R. M. Hosking, "Moments or L-moments? An example comparing two measures of distributional shape," *American Statistician*, vol. 46, no. 3, pp. 186–189, 1992.
- [31] J. R. M. Hosking, "Some theory and practical uses of trimmed L-moments," *Journal of Statistical Planning and Inference*, vol. 137, no. 9, pp. 3024–3039, 2007.
- [32] P. Royston, "Which measures of skewness and kurtosis are best?" *Statistics in Medicine*, vol. 11, no. 3, pp. 333–343, 1992.
- [33] R. M. Vogel and N. M. Fennessey, "L moment diagrams should replace product moment diagrams," *Water Resources Research*, vol. 29, no. 6, pp. 1745–1752, 1993.
- [34] P. Delicado and M. N. Gorla, "A small sample comparison of maximum likelihood, moments and L-moments methods for the asymmetric exponential power distribution," *Computational Statistics & Data Analysis*, vol. 52, no. 3, pp. 1661–1673, 2008.
- [35] J. R. M. Hosking, J. R. Wallis, and E. F. Wood, "Estimation of the generalized extreme-value distribution by the method of probability-weighted moments," *Technometrics*, vol. 27, no. 3, pp. 251–261, 1985.
- [36] J. R. M. Hosking and J. R. Wallis, "Parameter and quantile estimation for the generalized Pareto distribution," *Technometrics*, vol. 29, no. 3, pp. 339–349, 1987.
- [37] A. Hussein and J. Liu, "Parametric estimation of mixtures of two uniform distributions," *Journal of Statistical Computation and Simulation*, vol. 79, no. 3–4, pp. 395–410, 2009.
- [38] E. A. H. Elamir and A. H. Seheult, "Trimmed L-moments," *Computational Statistics & Data Analysis*, vol. 43, no. 3, pp. 299–314, 2003.
- [39] E. A. H. Elamir and A. H. Seheult, "Exact variance structure of sample L-moments," *Journal of Statistical Planning and Inference*, vol. 124, no. 2, pp. 337–359, 2004.
- [40] A. Necir and D. Meraghni, "Estimating L-functionals for heavy-tailed distributions and application," *Journal of Probability and Statistics*, vol. 2010, Article ID 707146, 34 pages, 2010.

- [41] M. C. Jones, "On some expressions for variance, covariance, skewness and L -moments," *Journal of Statistical Planning and Inference*, vol. 126, no. 1, pp. 97–106, 2004.
- [42] *TIBCO Spotfire S+ 8.1 for Windows*, TIBCO Software, Palo Alto, Calif, USA, 2008.
- [43] M. E. Mattson, J. P. Allen, W. R. Miller et al., "Project MATCH: Rationale and methods for a multisite clinical trial matching patients to alcoholism treatment," *Alcoholism: Clinical and Experimental Research*, vol. 17, no. 6, pp. 1130–1145, 1993.
- [44] J. P. Allen, M. E. Mattson, W. R. Miller et al., "Matching alcoholism treatments to client heterogeneity: project MATCH posttreatment drinking outcomes," *Journal of Studies on Alcohol*, vol. 58, no. 1, pp. 7–29, 1997.
- [45] S. Wolfram, *The Mathematica Book*, Wolfram Media, Champaign, Ill, USA, 5th edition, 1999.
- [46] R. M. Renner, "Evaluation by power series of the means of normal order statistics of samples of sizes six and seven," *Mathematical Chronicle*, vol. 4, no. 2-3, pp. 141–147, 1976.