

1-2016

Metapopulation Viability of Swamp Rabbits in Southern Illinois: Potential Impacts of Habitat Change

Christopher D. Robinson
Southern Illinois University Carbondale

Joanne C. Crawford
Southern Illinois University Carbondale

Lyann Corcoran
Southern Illinois University Carbondale


Eric M. Schaubert
Southern Illinois University Carbondale, schauber@siu.edu

Clayton K. Nielsen
Southern Illinois University Carbondale

Follow this and additional works at: http://opensiuc.lib.siu.edu/zool_pubs

Recommended Citation

Robinson, Christopher D., Crawford, Joanne C., Corcoran, Lyann, Schaubert, Eric M. and Nielsen, Clayton K. "Metapopulation Viability of Swamp Rabbits in Southern Illinois: Potential Impacts of Habitat Change." *Journal of Mammalogy* 97, No. 1 (Jan 2016): 68-79. doi:10.1093/jmammal/gyv154.



Metapopulation viability of swamp rabbits in southern Illinois: potential impacts of habitat change

| | |
|-------------------------------|--|
| Journal: | <i>Journal of Mammalogy</i> |
| Manuscript ID | JMAMM-2014-039.R3 |
| Manuscript Type: | Feature Article |
| Date Submitted by the Author: | n/a |
| Complete List of Authors: | Robinson, Christopher; Southern Illinois University Crawford, Joanne; Southern Illinois University, Carbondale and Department of Forestry Corcoran, Lyann; Southern Illinois University Schauber, Eric; Southern Illinois University, Cooperative Wildlife Research Lab Nielsen, Clayton; Southern Illinois University |
| Keywords: | bottomland, corridors, dispersal, fragmentation, metapopulation, Illinois, population viability analysis, swam rabbit, Sylvilagus aquaticus, model |
| | |



1
2
3
4
5
6
7 1 Corresponding Author:
8 2 Eric Schauber
9 3 SIU Cooperative Wildlife Research Laboratory
10 4 Mailcode 6504
11 5 Carbondale, IL 62901
12 6 Schauber@siu.edu
13 7 618-453-6940 (voice) 618-453-6944 (fax)
14 8
15 9

16 10 **RH: SWAMP RABBIT METAPOPOPULATION VIABILITY**
17 11

18 12 **Metapopulation viability of swamp rabbits in southern Illinois: potential impacts of**
19 13 **habitat change**
20 14

21 15 Christopher D. Robinson, Joanne C. Crawford, Lyann Corcoran, Eric M. Schauber*, and
22 16 Clayton K. Nielsen
23 17

24 18 *Cooperative Wildlife Research Laboratory, Mailcode 6504, Southern Illinois University,*
25 19 *Carbondale, IL 62901, USA (CDR, JCC, LC, EMS, CKN)*

26 20 *Department of Zoology, Mailcode 6501, Southern Illinois University, Carbondale, IL 62901,*
27 21 *USA (CDR, LC, EMS)*

28 22 *Department of Forestry, Mailcode 4411, Southern Illinois University, Carbondale, IL 62901,*
29 23 *USA (JCC, CKN)*

30 24 *Center for Ecology, Mailcode 317, Southern Illinois University, Carbondale, IL 62901, USA*
31 25 *(JCC, EMS, CKN)*
32 26
33 27
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 28 Swamp rabbits (*Sylvilagus aquaticus*) in southern Illinois exist as a metapopulation
8
9 29 due to fragmentation of the bottomland hardwood forests in which they live. This
10
11 30 fragmentation makes their persistence in Illinois uncertain. We used population viability
12
13 31 analysis (PVA) to estimate the probability of persistence of the swamp rabbit metapopulation
14
15 32 in Illinois, using a habitat suitability map we created and life history parameters drawn from
16
17 33 the literature. We varied the parameters used in our PVA from 50 to 150% of the initial value
18
19 34 to compare their effects on extinction risk and to direct future management and research. We
20
21 35 tested the effects of potential habitat loss and fragmentation by removing patches individually
22
23 36 and in groups from the analysis, and by adding 60, 120, and 180 m to the edge of all patches.
24
25 37 We also tested the potential effect of dispersal corridors by increasing dispersal between
26
27 38 connected patches. Under baseline conditions, the model suggests a 0% chance of quasi-
28
29 39 extinction (90% metapopulation decline) of swamp rabbits within 25 (or even 50) years.
30
31 40 Changes in fecundity values and the effects of catastrophic flooding had the greatest effect on
32
33 41 extinction risk, and changes in no other parameter yielded any appreciable impact. Removing
34
35 42 the largest patches from the population increased the 25-year risk of extinction to 4%,
36
37 43 whereas any other modifications to the habitat did not change the extinction risk. We suggest
38
39 44 that managers focus on sustaining habitat quality, particularly upland habitats adjacent to
40
41 45 occupied bottomland hardwood forests to improve the likelihood of swamp rabbit persistence
42
43 46 in Illinois.

43
44
45 48 Key words: bottomland, corridors, dispersal, fragmentation, Illinois, metapopulation, model,
46
47 49 population viability analysis, swamp rabbit, *Sylvilagus aquaticus*

48
49
50
51 51 *Correspondent: Schauber@siu.edu

1
2
3
4
5
6
7 52 Habitat fragmentation can have immediate and long-term harmful effects ranging
8
9 53 from the genetic level to the community level (Bowers et al. 1996; Dooley and Bowers 1998;
10 54 Haag et al. 2010; Krauss et al. 2010). Habitat fragments are typically surrounded by a hostile
11
12 55 matrix, which can act as a deterrent to dispersal attempts, reduce survival of individuals that
13
14 56 do attempt to disperse, and provide suitable habitat for predators or competitors that may not
15
16 57 have encountered the patch inhabitants otherwise (Rolstad 1991; Wilcove et al. 1997; Åström
17
18 58 and Pärt 2013).

19
20 59 Habitat loss has affected many kinds of wildlife, including those inhabiting
21
22 60 bottomland hardwood forests in the Mississippi River floodplain. Swamp rabbits (*Sylvilagus*
23
24 61 *aquaticus*) are bottomland hardwood forest specialists (Allen 1985) found throughout much
25
26 62 of the Mississippi River floodplain, making them an important indicator species for the
27
28 63 integrity of bottomland hardwood forests in this area. They are classified as endangered in
29
30 64 Indiana (Indiana Department of Natural Resources 2013) and rare in Missouri (Dailey et al.
31
32 65 1993; Scheibe and Henson 2003), and population declines have been noted throughout their
33
34 66 range (Platt and Bunch 2000). Swamp rabbit abundance in Illinois has apparently declined
35
36 67 since the 1970s, and swamp rabbits are now patchily distributed along the major rivers and
37
38 68 some interior river drainages in the southern portion of the state (Kjølhaug et al. 1987;
39
40 69 Barbour et al. 2001). Given this spatial structure, swamp rabbits are thought to exist as a
41
42 70 metapopulation (i.e., a system of local populations connected by dispersing individuals—
43
44 71 Hanski and Gilpin 1991), with small and large patches that may share dispersers scattered
45
46 72 across the landscape (Woolf and Barbour 2002; Roy Nielsen et al. 2008).

47 73 Human activities substantially affect habitat quality for swamp rabbits, which
48
49 74 predominantly prefer early-successional forests with close proximity to wooded wetlands
50
51 75 (Scharine et al. 2009, 2011). Selective logging or burning can replace natural disturbances
52
53 76 that create early-successional habitat (Lorimer 2001), leading to high-quality habitat in the
54
55

1
2
3
4
5
6
7 77 long-term, but clear-cutting large areas of land can have the opposite effect of selective
8
9 78 disturbance, decreasing the amount of high-quality habitat for swamp rabbits. Allen (1985)
10
11 79 suggested that conversion of land to agricultural production is the most significant cause of
12
13 80 swamp rabbit habitat loss, and considerable losses in swamp rabbit habitat have occurred
14
15 81 throughout their range, most notably near the northern edge (Sole 1994; Zollner et al. 2000a;
16
17 82 Fowler and Kissell 2007; Vale 2008).

18
19 83 Given the potential impact of past and future habitat alterations on swamp rabbits,
20
21 84 managers are interested in predicting the fate of the species in Illinois under a range of
22
23 85 possible action scenarios. Population viability analysis (PVA) uses quantitative models to
24
25 86 assess the future status of a population or metapopulation, predict the success of potential
26
27 87 recovery strategies, and identify aspects of a population (e.g., life history stages or
28
29 88 demographic processes) that should receive the highest priority in research and management
30
31 89 (Morris et al. 2002; Possingham et al. 2002; Ralls et al. 2002). Because PVAs are only as
32
33 90 accurate as the parameters and assumptions used to create them, their use in management has
34
35 91 been debated (Brook et al. 2002; Ellner et al. 2002). For instance, estimating parameters
36
37 92 accurately can be troublesome for rare species, on which these analyses are typically
38
39 93 performed (Holmes 2001; Ludwig and Walters 2002; Possingham et al. 2002; Ratcliffe et al.
40
41 94 2005), and most population viability analyses include stochasticity, incorporating
42
43 95 demographic and environmental variances that can be even more problematic to estimate than
44
45 96 average values (Beissinger and Westphal 1998; O'Grady et al. 2004). Although imperfect,
46
47 97 population viability analyses can make useful comparisons between management tools
48
49 98 (Starfield 1997; McCarthy and Broome 2000; Staples et al. 2004; Bakker and Doak 2009).

50
51 99 Population viability analyses have been implemented for swamp rabbits in the past,
52
53 100 using field data collected through a number of studies in Illinois, Indiana, and Missouri.
54
55 101 Woolf and Barbour (2002) and Roy Nielsen et al. (2008) both used spatially explicit stage-

1
2
3
4
5
6
7 102 structured models and predicted slight negative trends in swamp rabbit populations over time.
8
9 103 Woolf and Barbour (2002) predicted an 8.4% chance of the southern Illinois metapopulation
10
11 104 falling below 1,000 individuals over 25 years, and Roy Nielsen et al. (2008) predicted a slight
12
13 105 (< 10%) decline in patch occupancy over 25 years in Indiana. However, both assumed high
14
15 106 (see below) maximum carrying capacities (1.5 rabbits / ha) based on localized trapping
16
17 107 (Kjolhaug 1986). Woolf and Barbour (2002) also assumed that all patches within 200 m of
18
19 108 each other were parts of the same modeled patch, and they used an average dispersal distance
20
21 109 of 3 km; Roy Nielsen et al. (2008) also chose a high average dispersal distance of 1.5 km.
22
23 110 These parameter choices may explain why their models yielded such optimistic results. More
24
25 111 recent genetic evidence indicates that swamp rabbits in Illinois show limited success in
26
27 112 dispersing (Berkman et al. 2015), with strong genetic differentiation among subpopulations
28
29 113 separated by < 5 km. Also, a recent, extensive study of swamp rabbit home range size and
30
31 114 overlap in southern Illinois suggests that typical densities are likely well below 1.5 rabbits /
32
33 115 ha (Crawford 2014). Our 1st objective was to examine how these recent findings affect the
34
35 116 prognosis for swamp rabbit persistence in southern Illinois. We also expanded on earlier
36
37 117 PVAs by modeling specific changes to the habitat, including changes in fragmentation and
38
39 118 the addition of dispersal corridors, with the goal of suggesting future management practices.
40
41 119

42 MATERIALS AND METHODS

43 121 *Mapping suitable habitat and potential corridors.*—Following methods used by
44
45 122 LaRue and Nielsen (2008, 2011), we applied the analytical hierarchy process (AHP—Saaty
46
47 123 1980) to convert expert survey results into a habitat suitability map for the southern 28
48
49 124 counties of Illinois. This region largely consists of agricultural land and upland forests but
50
51 125 has bottomland hardwood forests in the Cache, Kaskaskia, Saline, Mississippi, Ohio, and Big
52
53 126 Muddy watersheds. The AHP hierarchy for identifying potential swamp rabbit habitat
54
55
56
57
58
59
60

1
2
3
4
5
6
7 127 involved: the goal (suitable swamp rabbit habitat), factors, and attributes within the factors.
8
9 128 We solicited expert opinion of the relative importance of habitat factors (landcover type, road
10 129 type, waterbody classification, and percent canopy cover) and relative suitability of attributes
11 130 within each factor (e.g., wetland forest within the "landcover type" factor; Table 1). We asked
12 131 12 researchers and managers familiar with swamp rabbit ecology to rate pairs of habitat
13 132 factors in terms of relative importance and pairs of attributes in terms of relative habitat
14 133 suitability for swamp rabbits, using a continuous rating scale from 1/9 to 9. For instance,
15 134 when comparing the suitability of agriculture and upland forest attributes of the "land cover"
16 135 feature, a rating of 1/6 would indicate that the expert considers agriculture to be one-sixth as
17 136 suitable as upland forest, whereas a rating of 1 would indicate equal suitability. Six surveys
18 137 were returned. Pairwise rating scores were made comparable by:

19
20
21
22
23
24
25
26
27
28 138
$$a_{ij}^* = a_{ij} / \sum_{k=1}^n a_{kj}$$

29
30
31

32 139 where a_{ij} is the raw score (relative importance or suitability) for attribute or factor i relative to
33 140 attribute or factor j , a_{ij}^* is the normalized score, and n is the number of attributes or factors
34 141 being compared (Kovacs et al. 2004). Then, weight (w_i) of each attribute or factor was
35 142 calculated as follows:

36
37
38
39
40 143
$$w_i = \sum_{j=1}^n a_{ij}^*$$

41
42
43

44 144 Finally, these weights were averaged across the 6 experts who returned surveys.

45
46
47 145 We applied these weightings to assess the suitability of each 30 × 30-m pixel based on
48 146 its attribute value for each habitat factor: land cover type (United States Geological Survey
49 147 2006, 2008), forest canopy cover (United States Geological Survey 2001), water bodies
50 148 (United States Department of Commerce 2011), and roads (United States Department of
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 149 Commerce 2011). We modified some of the data layers for our analysis. Roads, streams, and
8
9 150 water-bodies used in the habitat model were given a 0.25-km buffer. The road dataset
10
11 151 contained 5 classes, but we combined “Primary highway with limited access” and “Primary
12
13 152 road without limited access” into 1 class (Highways). Stream and water-body classes were
14
15 153 grouped based on perennial/intermittent status and divided into streams, water bodies, or
16
17 154 shorelines. The 2001 National Land Cover Dataset contained 30 classes, but we grouped
18
19 155 similar classes into 8 categories thought to be important to swamp rabbit biology (Chapman
20
21 156 and Feldhamer 1981). We also converted the canopy cover data from a continuous variable
22
23 157 into 4 categories (Table 1). These geospatial data layers were then reclassified by multiplying
24
25 158 attribute weights by the relevant factor weights. We summed weights over all attributes and
26
27 159 factors for each pixel using the Raster Calculator in ArcGIS 10.1 (ESRI 2012) to generate a
28
29 160 habitat suitability value for that pixel. We divided the observed range of habitat suitability
30
31 161 values into 5 categories based on observed breaks in the data.

32 162 We considered the highest 2 categories (highest 34% of values) to be suitable habitat
33
34 163 for use in our model. We chose the 34% cutoff values based on an apparent break in the
35
36 164 distribution of suitability values and because swamp rabbits in southern Illinois studied by
37
38 165 Scharine et al. (2009) and Crawford (2014) tended to be located in these categories. Allen
39
40 166 (1985) suggested that areas of contiguous habitat >100 ha are required to support a swamp
41
42 167 rabbit population, but Scharine et al. (2009) located swamp rabbits in habitat patches < 25 ha.
43
44 168 Due to the presence of swamp rabbits in patches far smaller than 100 ha (Kjolhaug 1986;
45
46 169 Porath 1997; Scheibe and Henson 2003; Scharine et al. 2009), we included suitable habitat
47
48 170 areas > 50 ha, as well as patches 25-50 ha that were within 2 km of a patch > 50 ha. We
49
50 171 anticipated that swamp rabbits would rarely disperse as far as 2 km, but included such
51
52 172 isolated patches to better evaluate how decreased fragmentation could influence the
53
54 173 likelihood of persistence (discussed below). From these areas, we used those patches with

Comment [EMS1]: Fixed

Comment [CLB2]: Please cite software here as (Author Year) and in lit cit as Author. Year. Title. Publisher. Please add this citation to the lit cit. For example: (ESRI 2014) and in lit cit: ESRI. 2014. ArcGIS. Ver. 10.2.1. Environmental System Research Institute, Inc. Redlands, California.

1
2
3
4
5
6
7 174 confirmed swamp rabbit presence in the past 30 years (Kjolhaug 1986; Porath 1997; Woolf
8
9 175 and Barbour 2002; Scharine et al. 2009; Crawford 2014) to run simulations.

10 176 Berkman et al. (2015) found limited genetic connectivity of swamp rabbit populations
11
12 177 in the Cache River watershed of Illinois and suggested that forested corridors may improve
13
14 178 metapopulation viability by increasing dispersal between populations. We used ArcGIS to
15
16 179 conduct least-cost path analysis, to identify the most permeable portions of the Cache River
17
18 180 watershed for potential dispersal (Singleton et al. 2002; Adriaensen et al. 2003; LaRue and
19
20 181 Nielsen 2008). This analysis is based on simulating movement over a resistance map, in
21
22 182 which each pixel is assigned a resistance to movement based on its habitat characteristics and
23
24 183 cost is the total resistance encountered over the length of a path (Singleton et al. 2002;
25
26 184 Wikramanayake et al. 2004; LaRue and Nielsen 2008). We employed the AHP to assign each
27
28 185 pixel a resistance to dispersal using the same methodology, habitat factors and attributes, and
29
30 186 experts as for the map of habitat suitability. We used the resulting resistance map to generate
31
32 187 least-cost paths 1 pixel wide between occupied habitat patches.

33
34 188 *Population viability analysis.*—Like Woolf and Barbour (2002), we used a Lefkovich
35
36 189 matrix model (Caswell 2001) to simulate rabbit population dynamics in each patch. The
37
38 190 model included a juvenile and adult stage with 1-year time steps and assumed a pre-breeding
39
40 191 census, such that the juvenile class comprised individuals nearly 1 year old; therefore,
41
42 192 fecundity parameters incorporated survival of individuals through their 1st year. This model
43
44 193 also assumed a female-only population, and allowed patch carrying capacity to differ based
45
46 194 on patch size and suitability.

47
48 195 We used the program RAMAS GIS, version 5.0 (Akçakaya 2005) to run all
49
50 196 simulations for this study. We ran all simulations with 2,500 repetitions for 25 simulated
51
52 197 years, and our primary outputs were the distribution of population sizes each year, as well as
53
54 198 the probability (i.e., proportion of repetitions) of the total metapopulation abundance (i.e.,

1
2
3
4
5
6
7 199 number of female rabbits alive) dropping below our quasi-extinction threshold (Ginzburg et
8
9 200 al. 1982), which we set at 10% of the total regional carrying capacity, within 25 years. We
10
11 201 then compared these outputs between simulations run with varying parameter values
12
13 202 (intermediate value versus 50 to 150% of intermediate value) and management scenarios (i.e.,
14
15 203 changes to the habitat suitability map based on possible interventions). Because the
16
17 204 estimated quasi-extinction probability under intermediate parameter values was essentially
18
19 205 0% (see “Results”), we also examined the effects of simulated habitat management
20
21 206 interventions using a fecundity value 40% lower than the intermediate value.

22 207 The maximum female carrying capacity ($K_{\max,i}$) of each patch (i) equaled the patch
23
24 208 area divided by the mean size of swamp rabbit home ranges, as we assumed a uniform
25
26 209 distribution of individuals. We used an intermediate home range size of 1.93 ha (2.54 ha with
27
28 210 a 24% overlap, based on core areas (50% isopleths from fixed kernel utilization distributions)
29
30 211 of $n = 60$ swamp rabbits—Crawford 2014). Depending on the relative amounts of patch i
31
32 212 made up of suitable and highly suitable habitat, the intermediate carrying capacity (and initial
33
34 213 abundance) of each patch (K_i) ranged from $0.8K_{\max,i}$ (all suitable) to $K_{\max,i}$ (all highly
35
36 214 suitable). As per Woolf and Barbour (2002), we chose to model density dependence as a
37
38 215 ceiling, in which the population in each patch (N_i) increases based on the density-independent
39
40 216 matrix model until they reach or exceed K_i , at which point additional individuals above K_i are
41
42 217 removed from the population (Akçakaya 2005). In intermediate-parameter runs, we kept K_i
43
44 218 constant through time.

45 219 In RAMAS GIS, dispersal rates (m_{ij}) are calculated from each patch to all other
46
47 220 patches based on a migration-distance function:

$$m_{ij} = a \cdot \exp\left(\frac{-D_{ij}^c}{b}\right) \text{ if } D_{ij} < D_{\max}$$

$$m_{ij} = 0 \text{ if } D_{ij} > D_{\max},$$

Equation 1

1
2
3
4
5
6
7 221 where m_{ij} is the proportion of animals in patch i that successfully disperse to patch j in each
8
9 222 year, and D_{ij} is the shortest straight-line distance (km) between the edges of the two patches.
10
11 223 D_{max} represents the maximum dispersal distance, and a , b , and c define the shape of the
12
13 224 migration-distance curve: a is the maximum dispersal rate (i.e., maximum fraction of
14
15 225 individuals from a patch that successfully disperse to any single other patch), b is the average
16
17 226 dispersal distance (km) when $c = 1$, and c determines the shape of the curve.

18 227 Dispersal data for swamp rabbits are limited, but Forys (1995) found that the Lower
19
20 228 Keys marsh rabbit (*Sylvilagus palustris hefneri*) had an average dispersal distance of 300 m.
21
22 229 However, marsh rabbits evolved in a naturally patchy habitat (Forys and Humphrey 1996)
23
24 230 and cottontail rabbits (*Sylvilagus* spp.) do not typically disperse long distances (Shields 1960;
25
26 231 Chapman and Trethewey 1972; Fenderson et al. 2014), which suggests 300 m is likely an
27
28 232 overestimate for swamp rabbit dispersal. Genetic evidence also indicates limited successful
29
30 233 dispersal of swamp rabbits within one watershed in Illinois (Berkman et al. 2015), so we used
31
32 234 an average dispersal distance $b=200$ m. Since Akçakaya and Raphael (1998) suggested that
33
34 235 only a small portion of individuals in a patch will actually disperse to a single neighboring
35
36 236 patch, we set a equal to 0.1. We set the intermediate value of parameter $c = 1$ (exponential
37
38 237 dispersal kernel). Despite their rarity, long-distance dispersers are generally more important
39
40 238 than short-distance dispersers to the persistence and genetic mixing of a metapopulation
41
42 239 (Johst et al. 2002; Trakhtenbrot et al. 2005). We therefore set $D_{max} = 4$ km, although dispersal
43
44 240 in the model beyond 1.5 km was vanishingly rare based on the intermediate values of a , b ,
45
46 241 and c .

47 242 We used an intermediate annual survival rate of 0.3 based on survival analysis of 79
48
49 243 radiocollared swamp rabbits in southern Illinois (Crawford 2014). Swamp rabbits are legal
50
51 244 game species in Illinois, and this estimate included mortality due to hunting. Due to a lack of
52
53 245 data, we assumed that survival rates of juveniles (1 year old) and adults (>1 year old) were

1
2
3
4
5
6
7 246 equal. Using values from Holler et al. (1963) and Sorensen et al. (1968), both from Missouri,
8
9 247 we estimated average numbers of litters per female per year and offspring per litter per
10
11 248 female from these studies as 2.8 and 3.2, respectively. These estimates yielded an estimate of
12
13 249 8.96 offspring per year per female, which is similar to estimates from other studies (Hunt
14
15 250 1959; Toll et al. 1960; Hill 1967). We used an all-female model, so we halved this value to
16
17 251 4.48 female offspring per year per female. Since we assumed equal survival rates regardless
18
19 252 of age, the fecundity matrix element equaled 4.48 multiplied by the 0.3 survival rate, or 1.344
20
21 253 female recruits per female per year for both stages. In a simple 2-stage Lefkovich matrix,
22
23 254 these intermediate survival and fecundity values produce a baseline, deterministic estimate of
24
25 255 the finite rate of increase: $\lambda = 1.644$.

26 256 Flooding can increase swamp rabbit mortality by predation, starvation, hunting, and
27
28 257 drowning, as well as decrease their reproduction due to embryo resorption (Conaway et al.
29
30 258 1960; Platt and Bunch 2000; Zollner et al. 2000b). The quantitative effect of catastrophic
31
32 259 flooding on rabbit populations is poorly understood. Hamilton et al. (2010) estimated that
33
34 260 severe flooding reduced monthly survival of riparian brush rabbits (*Sylvilagus bachmani*
35
36 261 *riparius*) by about 33%, from 0.90-0.96 to 0.61-0.64. Previous floods had greatly reduced the
37
38 262 number of riparian brush rabbits trapped, but rigorous estimates of effects on abundance and
39
40 263 survival were not available. Woolf and Barbour (2002) assumed a 60% decline in population
41
42 264 abundance during a catastrophic flood, so we used this as our intermediate value of the effect
43
44 265 of catastrophes. As catastrophes are rare by definition, we assumed a 10% annual occurrence
45
46 266 rate for our intermediate value, and catastrophes occurred regionally, impacting all patches
47
48 267 simultaneously. The value of 10% roughly corresponds to the frequency of major floods
49
50 268 based on river stage data from the United States Geological Survey (2011) and the US Army
51
52 269 Corps of Engineers (2011). These intermediate values for the effect and occurrence rate of
53
54 270 catastrophic floods reduced deterministic λ to 1.545.

1
2
3
4
5
6
7 271 RAMAS GIS allows for both demographic and environmental stochasticity in
8
9 272 modeling population growth. For demographic stochasticity, RAMAS selects the number of
10
11 273 survivors per patch per year from a binomial distribution and the number of offspring per
12
13 274 patch per year from a Poisson distribution. For environmental stochasticity, RAMAS samples
14
15 275 each vital rate from a lognormal distribution (Akçakaya 2005); means and standard
16
17 276 deviations for these distributions are set by the user. We estimated standard deviations by
18
19 277 applying the coefficients of variation for fecundity and survival estimates of Lower Keys
20
21 278 marsh rabbits to our values (Forys 1995; LaFever et al. 2008). The resulting standard
22
23 279 deviations were 0.166, 0.076, and 0.051 for fecundity, juvenile survival, and adult survival,
24
25 280 respectively.

26 281 Environmental conditions are typically more similar in nearby patches than in distant
27
28 282 patches. We modeled spatial autocorrelation in survival and fecundity values between patches
29
30 283 due to environmental similarities using the correlation-distance function:

$$\rho_{ij} = x \times \exp(-D_{ij}^z/y)$$

31
32 284
33
34 285 where ρ_{ij} is the coefficient of correlation between patches i and j , and D_{ij} is the distance (km)
35
36 286 between the centers of the two patches. The parameters x , y , and z define the shape of the
37
38 287 correlation-distance curve: x is the maximum correlation (as $D_{ij} \rightarrow 0$), y is the rate at which
39
40 288 correlation declines with increasing distance, and z determines the shape of the curve. Woolf
41
42 289 and Barbour (2002) found no significant difference in persistence probability caused by
43
44 290 varying environmental correlation values so we used their intermediate values: $x = 1$, $z = 1$,
45
46 291 and $y = 40$.

47 292 We performed a sensitivity analysis to test the effect of changes in parameters that
48
49 293 could be measured or estimated inaccurately, or affected by management efforts. We
50
51 294 individually varied each parameter in 5% increments from 50 to 150% of the intermediate
52
53 295 value (Table 2), and we ran all simulations with 2,500 replications each for 25 simulated
54

1
2
3
4
5
6
7 296 years, using intermediate values for all other parameters. When varying mean survival and
8
9 297 fecundity values, standard deviations (environmental stochasticity) changed accordingly,
10
11 298 based on the coefficients of variation from Forys (1995) and LaFever et al. (2008). Although
12
13 299 fecundity values incorporated survival to age 1, we kept the fecundity value constant when
14
15 300 measuring sensitivity to survival rates. We also explored the effect of a trend in carrying
16
17 301 capacity, such that the carrying capacity of each patch (i) changed linearly from the initial
18
19 302 carrying capacity ($K_{0,i}$) to the new carrying capacity at year 25 ($0.5K_{0,i} \leq K_{25,i} \leq 1.5 K_{0,i}$).

20 303 To identify which habitat patches may be most important to swamp rabbit viability in
21
22 304 Illinois, we removed one patch at a time with replacement from the habitat map for each
23
24 305 simulation, and also ran two simulations removing the largest and smallest 25% of patches by
25
26 306 population size. To estimate the effect of habitat fragmentation, we reduced fragmentation by
27
28 307 adding 60, 120, and 180 m to the perimeters of all patches, such that the initial test was used
29
30 308 as a “high fragmentation” comparison. We separated the effect of habitat fragmentation *per*
31
32 309 *se* from that of habitat amount by setting the initial abundance and carrying capacity of each
33
34 310 expanded patch equal to that of the original patch or the sum of all original patches (initial
35
36 311 test) the expanded patch incorporated (i.e., total carrying capacity of the landscape was not
37
38 312 changed). We found intermediate model results were so optimistic that a positive change
39
40 313 caused by the addition of dispersal corridors would not be detected, so we set fecundity at
41
42 314 60% of the intermediate test and compared results to the corresponding test from the
43
44 315 sensitivity analysis.

45 316 We used the model to assess the benefit of improving dispersal along corridors
46
47 317 identified by our least-cost path analysis. Because corridors would likely be used by the small
48
49 318 number of individuals near the start of the corridor in each patch, we tested the effect of these
50
51 319 dispersal corridors by adding 5% to the dispersal rate between patches connected by corridors
52
53 320 (i.e. adding 0.05 to m_{ij} calculated via Equation 1), or in the case of one corridor that

1
2
3
4
5
6
7 321 connected seven patches, by adding 0.0083 to the migration rate for each connected pair of
8
9 322 patches. Because we were testing the use of corridors to improve dispersal between patches,
10
11 323 we did not artificially join connected patches as a single patch in our model. As with the
12
13 324 habitat fragmentation tests, we set fecundity at 60%, using the corresponding tests from the
14
15 325 sensitivity analysis as a comparison.
16
17 326

18 327 **RESULTS**

19
20 328 *Mapping suitable habitat and potential corridors.*—Experts deemed that land cover
21
22 329 was the most important factor for identifying suitable swamp rabbit habitat as well as
23
24 330 resistance to dispersal (Fig. 1). Among land cover types (attributes), wetland forest was
25
26 331 deemed most suitable for swamp rabbits and open water and developed/barren lands were
27
28 332 most resistant to dispersal (Table 1). Canopy cover and water bodies were both deemed
29
30 333 intermediate in importance (Fig. 1), with high canopy cover and perennial stream/ditch
31
32 334 providing highest suitability and lowest resistance (Table 1). Roads were considered to be
33
34 335 relatively unimportant (Fig. 1), and unpaved roads were considered most suitable and least
35
36 336 resistant to dispersal (Table 1).

37
38 337 The resulting map of habitat in southern Illinois (Fig. 2a) consisted of 62 patches of
39
40 338 suitable and highly suitable swamp rabbit habitat totaling just under 12,000 ha, with mean
41
42 339 and median patch sizes of 193 ha and 86 ha, respectively (range = 25-3,818 ha). Of these
43
44 340 patches, 19 patches were < 50 ha, 17 patches were 50-100 ha, and 26 patches were > 100 ha,
45
46 341 and the initial metapopulation abundance was 5,577 individuals, resulting in a quasi-
47
48 342 extinction threshold of 558 individuals. Least-cost path analysis identified 31 potential
49
50 343 dispersal corridors, ranging from 0.7 km to 19.1 km long, linking suitable habitat patches in
51
52 344 the Cache River watershed (Fig. 2b).
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 345 *Population viability analysis.*—The initial population viability analysis with
8
9 346 intermediate parameter values predicted relatively little change in the swamp rabbit
10
11 347 metapopulation of southern Illinois over the next 25 years, with a median final
12
13 348 metapopulation abundance of 5,570 rabbits (0.13% decline; Fig. 3). The model estimated a
14
15 349 0% chance of the swamp rabbit metapopulation declining below 10% of the initial abundance
16
17 350 (quasi-extinction) within 25 years (95% *CI*: 0-1.77%). Extending intermediate-parameter
18
19 351 simulations to 50 years resulted in minuscule changes to median final abundance (5,564
20
21 352 rabbits) and quasi-extinction risk (0%).

22 353 Population growth and viability were most sensitive to changes in fecundity: the
23
24 354 range of fecundity values we considered produced deterministic λ values ranging from 0.97
25
26 355 to 2.32, and a 50% reduction in fecundity caused the risk of quasi-extinction to exceed 75%
27
28 356 (Fig. 4). Population growth and viability were moderately sensitive to changes in the effect
29
30 357 of catastrophic flooding, resulting in a quasi-extinction probability as high as 40% (Fig. 4)
31
32 358 despite deterministic λ only varying between 1.50 and 1.60). No other parameters, when
33
34 359 changed, yielded an appreciable impact on the risk of quasi-extinction for the metapopulation
35
36 360 (i.e. quasi-extinction risk exceeding the 95% *CI* from the intermediate test).

37 361 The removal of any single patch had no impact on the quasi-extinction risk of the
38
39 362 overall metapopulation when compared to the original model (0%), nor did removal of the
40
41 363 smallest 25% of patches, but removal of the largest 25% of patches (72% of the
42
43 364 metapopulation abundance) increased the quasi-extinction risk to 4%. Simulations with
44
45 365 fecundity reduced by 40% yielded a probability of quasi-extinction of 20% (95% *CI*: 18-
46
47 366 22%), which we use as a baseline for comparing with simulated habitat or corridor
48
49 367 manipulations. Adding 60, 120, and 180 m to the edge of all patches increased the total patch
50
51 368 area to approximately 17,800 ha, 21,500 ha, and 24,600 ha, respectively, and adjusting for
52
53 369 this increase in habitat area to isolate the effects of habitat fragmentation per se did not

1
2
3
4
5
6
7 370 substantially affect the quasi-extinction risk for any of the three scenarios (21%, 20%, and
8
9 371 20%, respectively). Adding corridors also yielded no appreciable change in the predicted
10
11 372 probability of quasi-extinction (21%).
12
13 373

14 374 **DISCUSSION**

15
16 375 The swamp rabbit has high habitat specificity and the availability and abundance of
17
18 376 swamp rabbit habitat have declined across its range, making an assessment of its viability in
19
20 377 Illinois important to its conservation. Our findings suggest that swamp rabbit populations
21
22 378 show no risk of extinction in Illinois in the next 50 years. The predicted extinction risk was
23
24 379 most sensitive to changes in fecundity and the effect of catastrophes, and was relatively
25
26 380 insensitive to changes in other parameter values. Our findings also suggest that efforts to
27
28 381 increase connectivity, e.g., by adding dispersal corridors, will likely not benefit the overall
29
30 382 metapopulation as much as efforts to increase the overall amount and quality of habitat.

31
32 383 Our model suggests a 0% chance of quasi-extinction within 50 years, with a median
33
34 384 percent decline of < 1%. Similarly to Woolf and Barbour (2002), our model was sensitive to
35
36 385 few of its parameters. The model was most sensitive to changes in fecundity, and the
37
38 386 fecundity value we used was estimated based on two studies of captive individuals from
39
40 387 Missouri conducted over 40 years ago. The rabbits in these studies may have reproduced less
41
42 388 than they would have in the wild due to high stress resulting from limited space in the
43
44 389 enclosure, as crowding can lead to higher than normal rates of litter resorption (Conaway et
45
46 390 al. 1960; Holler et al. 1963; Sorensen et al. 1968), which suggests that our fecundity value
47
48 391 may be an underestimate yet still resulted in a 0% chance of extinction. Other studies have
49
50 392 reported similar reproductive rates in captive and free-living *Sylvilagus*, however (e.g.,
51
52 393 Kirkpatrick and Baldwin 1974). Kjolhaug (1986) estimated an 18% annual survival rate for
53
54 394 swamp rabbits in Illinois, which would not increase the likelihood of extinction based on our
55

1
2
3
4
5
6
7 395 sensitivity analysis. However, when this new survival value was used to calculate fecundity,
8
9 396 the risk of extinction increased to about 20%, suggesting that accurate estimates of both
10
11 397 demographic rates are very important to achieve reliable estimates of absolute extinction risk.

12
13 398 Catastrophes also had a large impact on the risk of quasi-extinction. We based the
14
15 399 occurrence rate of catastrophes on flood gauge measurements, but the actual occurrence rate
16
17 400 of catastrophic floods that affect swamp rabbits in southern Illinois is unknown, and climate
18
19 401 change models predict an increase in the occurrence of flooding events around the
20
21 402 Mississippi River (Pinter et al. 2006) and at the global scale (Bouwer 2011; Wilby and
22
23 403 Keenan 2012). Although the occurrence rate of catastrophes appeared to have no impact on
24
25 404 extinction risk, the frequency of catastrophes would amplify the impact the effects
26
27 405 catastrophes have on the population. The model was more sensitive to the effect that
28
29 406 catastrophic floods have on swamp rabbit populations than occurrence rate, but the value for
30
31 407 the effect of catastrophic floods was assumed with little empirical data to support it,
32
33 408 suggesting more research is needed to determine their impacts. Severe flooding in 1976 and
34
35 409 again in 1997 appeared to nearly eradicate the only known (at the time) population of riparian
36
37 410 brush rabbits (United States Fish and Wildlife Service 1998), and Hamilton et al. (2010)
38
39 411 estimated that a severe (but apparently brief) flood reduced survival rate of riparian brush
40
41 412 rabbits by approximately 30%. However, the impact of flooding is likely to vary strongly
42
43 413 with local topography as well as flood duration. Crawford (2014) reported that swamp
44
45 414 rabbits in some but not all sites within the Cache River National Wildlife Refuge, Illinois,
46
47 415 were able to find refuge from a severe and persistent flood event. Combined, these
48
49 416 observations reinforce the need to ensure that upland refugia remaining near fragmented
50
51 417 populations vulnerable to flooding are given conservation priority.

52
53 418 The model appeared to be relatively insensitive to changes in home range area (which
54
55 419 determined carrying capacity) and the trend in carrying capacity, but our simulations showed

1
2
3
4
5
6
7 420 that the total metapopulation abundance tended to stay at approximately the total carrying
8
9 421 capacity of the landscape. Thus, home range area and temporal changes in suitability are
10
11 422 likely to influence the overall abundance of the species in Illinois. Removal of any single
12
13 423 patch had no appreciable impact on the extinction risk for the metapopulation as a whole,
14
15 424 suggesting that individual patches are not crucial to metapopulation persistence, concurring
16
17 425 with the results of our dispersal and connectivity tests. However, removal of the largest 25%
18
19 426 of patches did yield a negative effect, pointing out that further habitat loss and degradation
20
21 427 can reverse the current prediction of low extinction risk. Conversely, our results confirm that
22
23 428 increasing the amount and suitability (and therefore local carrying capacity) of habitat can
24
25 429 further increase the total swamp rabbit metapopulation size (i.e., number of rabbits).

26 430 Swamp rabbit persistence in the model was relatively insensitive to changes in
27
28 431 average dispersal distance or maximum dispersal rates, as well as to the shape of the dispersal
29
30 432 kernel (parameter c). Decreased fragmentation per se and addition of dispersal corridors also
31
32 433 had little effect on model results, especially when compared to the effect of habitat loss, again
33
34 434 suggesting that habitat quality and quantity is more important to manage than habitat
35
36 435 connectivity. Although this seems contrary to the logic behind a metapopulation, the
37
38 436 intermediate parameter values resulted in very rare dispersal between patches at the average
39
40 437 nearest neighbor distance of 2.09 km. Rare dispersal coupled with the strong baseline
41
42 438 population growth rates meant that in situ population dynamics were much more important
43
44 439 for persistence than metapopulation dynamics. Akçakaya et al. (2003) obtained similar
45
46 440 results for metapopulation viability of California least terns (*Sterna antillarum browni*):
47
48 441 where strong in situ population growth resulted in essentially zero near-term risk of
49
50 442 extinction under intermediate parameter values, and greater sensitivity to demographic rates
51
52 443 than dispersal parameters. In contrast, Medici and Desbiez (2012) found that dispersal was
53
54 444 crucial for persistence of fragmented populations of lowland tapirs (*Tapirus terrestris*),

1
2
3
4
5
6
7 445 whose maximum population growth rates were much lower than we estimate for swamp
8 rabbits. Frequent dispersal also enables persistence of American pikas (*Ochotona princeps*)
9
10 447 in a metapopulation occupying a complex of small mine tailing patches (Moilanen et al.
11
12 448 1998; Smith and Nagy 2015), a system that differs from ours in that dispersal is frequent and
13
14 449 extinction rates are very high because most patches typically only support a few individuals.
15
16 450 We would expect connectivity to play a larger role for swamp rabbits over longer timescales
17
18 451 (with increased probability of extinction), but the amount and distribution of habitat as well
19
20 452 as flooding regimes are likely to change dramatically over longer time horizons.

21
22 453 Our findings are based on several assumptions that require further validation. The
23
24 454 habitat suitability map we used to build our models assumed a clear dichotomy between
25
26 455 suitable and unusable habitat in southern Illinois, although swamp rabbits have been found in
27
28 456 lower quality or smaller areas than the cut-off we chose (Rubert 2007; Scharine et al. 2009,
29
30 457 2011). Empirical validation or participation by a greater number of experts would improve
31
32 458 confidence in habitat suitability values. Also, we treated patches separated by > 30 m as 2
33
34 459 separate patches, even though the patches may have been separated by a river or a lower
35
36 460 quality wooded area. Small areas such as these that did not register as suitable habitat may
37
38 461 not act as a barrier for a swamp rabbit in reality, suggesting our map may represent an
39
40 462 artificially fragmented landscape. We also assumed that all patches identified were fully
41
42 463 occupied at the start, although this is likely a false assumption (Kjolhaug 1986; Barbour et al.
43
44 464 2001). However, our results indicate that predicted persistence was much more sensitive to
45
46 465 demographic parameters than to the amount or connectivity of suitable habitat. Even
47
48 466 removing the patches containing most of the simulated rabbits in the model only increased
49
50 467 the predicted probability of extinction to 4%. Our findings point to refining estimates of
51
52 468 those demographic rates rather than estimates of habitat suitability as the most important
53
54 469 avenue to increasing confidence in PVA results.

1
2
3
4
5
6
7 470 There has been much debate about whether the primary approach of species
8
9 471 conservation should be to increase connectivity or to conserve existing habitat (Simberloff
10
11 472 and Cox 1987; Simberloff et al. 1992; Beier and Noss 1998). Enhancing connectivity can
12
13 473 increase genetic variability, decrease local extinctions, and increase abundance in patches
14
15 474 with smaller populations (Fahrig and Merriam 1985; Dunning et al. 1995; Haddad and Baum
16
17 475 1999). However, our models suggest that efforts to improve dispersal amongst swamp rabbit
18
19 476 populations in southern Illinois will have less impact on their overall persistence in the state
20
21 477 than improvements in patch habitat quality. This conclusion is in concordance with the
22
23 478 outcome of agent-based simulation models of Ye et al. (2013), who found that long-term
24
25 479 abundance of habitat specialists in heterogeneous environments depended mainly on the size
26
27 480 and quality of suitable habitat patches whereas that of generalists was more strongly
28
29 481 influenced by patch isolation. Thus, while habitat quality is obviously important for
30
31 482 demographic parameters and population persistence of all species, it may be particularly
32
33 483 crucial (relative to connectivity) for habitat specialists with low vagility.

34 484 This study suggests some potential changes to management practices that would help
35
36 485 swamp rabbit population in Illinois and will likely have similar positive effects in other
37
38 486 fragmented areas in the northern portion of the swamp rabbit range (e.g. Indiana—Roy
39
40 487 Nielsen et al. 2008), as well as for other species occupying similar habitats. Wildlife
41
42 488 managers working to conserve individual species are often torn between improving and
43
44 489 expanding existing habitat (Hobbs 1992; Beier and Noss 1998; Hootor et al. 2000). Woolf
45
46 490 and Barbour (2002) and Scharine et al. (2009) suggested that maintaining quality of existing
47
48 491 patches, particularly upland areas adjacent to the bottomland hardwood forests currently
49
50 492 occupied, should be a major goal in swamp rabbit conservation, and our results agree. High-
51
52 493 quality upland habitat (e.g., with thicker understory growth for food and protection) provides
53
54 494 a refuge from flooding without high predation (Kjolhaug et al. 1987), which will decrease the

1
2
3
4
5
6
7 495 effect of floods on survival and population abundance, and increase reproduction during
8
9 496 flooding periods. We also identified further research to improve our knowledge of swamp
10
11 497 rabbit persistence in southern Illinois and create more accurate models of swamp rabbit
12
13 498 population dynamics in the future. Most important are better estimates of fecundity of swamp
14
15 499 rabbits in Illinois and the effect of floods on swamp rabbit populations. These 2 parameters
16
17 500 had the greatest effect on extinction risk in the model and are the least studied in swamp
18
19 501 rabbits range wide.

502 **ACKNOWLEDGMENTS**

20
21
22 503 We thank M. Wefer and J. Cole and the Illinois Department of Natural Resources for
23
24 504 funding this project via Federal Aid in Wildlife Restoration Project W-106-R. We also thank
25
26 505 the Cooperative Wildlife Research Laboratory, Department of Zoology, and the Graduate
27
28 506 School at Southern Illinois University for support. We also wish to thank M. LaRue and G.
29
30 507 Feldhamer for their assistance with this project, and 2 anonymous reviewers whose
31
32 508 comments and suggestions substantially improved this manuscript.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Literature Cited

- 509
510 Adriaensen, F., J. P. Chardon, G. De Blust, E. Swinnen, S. Villalba, H. Gulinck, AND E.
511 Matthysen. 2003. The application of 'least-cost' modeling as a functional landscape
512 model. *Landscape and Urban Planning* 64:233-247.
- 513 Akçakaya, H. R. 2005. RAMAS GIS: Linking spatial data with population viability analysis
514 (version 5.0). Applied Biomathematics, Setauket, New York.
- 515 Akçakaya, H. R., J. L. Atwood, D. Breininger, C. T. Collins, AND B. Duncan. 2003.
516 Metapopulation viability of the California least tern. *Journal of Wildlife Management*
517 67:829-842.
- 518 Akçakaya, H. R., AND M. G. Raphael. 1998. Assessing human impact despite uncertainty:
519 viability of the northern spotted owl metapopulation in the northwestern USA.
520 *Biodiversity and Conservation* 7:875-894.
- 521 Allen, A. W. 1985. Habitat suitability index model: swamp rabbit. United States Fish and
522 Wildlife Service. Habitat Evaluations Procedure Group. Fort Collins, Colorado, USA.
- 523 Åström, J., AND T. Pärt. 2013. Negative and matrix-dependent effects of dispersal corridors
524 in an experimental metacommunity. *Ecology* 94:72-82.
- 525 Bakker, V. J., AND D. F. Doak. 2009. Population viability management: ecological standards
526 to guide adaptive management for rare species. *Frontiers in Ecology and the*
527 *Environment* 7:158-165.
- 528 Barbour, M. S., A. Woolf, AND J. W. Porath. 2001. Recent trends and future outlook for the
529 swamp rabbit (*Sylvilagus aquaticus*) in Illinois. *Transactions of the Illinois State*
530 *Academy of Science* 94:151-160.
- 531 Beier, P., AND R. F. Noss. 1998. Do habitat corridors provide connectivity? *Conservation*
532 *Biology* 12:1241-1242.

- 1
2
3
4
5
6
7 533 Beissinger, S. R., AND M. I. Westphal. 1998. On the use of demographic models of
8
9 534 population viability in endangered species management. *Journal of Wildlife*
10
11 535 *Management* 62:821-841.
- 12
13 536 Berkman, L. K., C. K. Nielsen, C. L. Roy, AND E. J. Heist. 2015. Comparative genetic
14
15 537 structure of sympatric leporids in southern Illinois. *Journal of Mammalogy* 96:552-
16
17 538 563.
- 18
19 539 Bouwer, L. M. 2011. Have disaster losses increased due to anthropogenic climate change?
20
21 540 *Bulletin of the American Meteorological Society* 92:39-46.
- 22
23 541 Bowers, M. A., S. F. Matter, J. L. Dooley, Jr., J. L. Dauten, AND J. A. Simkins. 1996.
24
25 542 Controlled experiments of habitat fragmentation: a simple computer simulation and a
26
27 543 test using small mammals. *Oecologia* 108:182-191.
- 28
29 544 Brook, B. W., M. A. Burgman, H. R. Akçakaya, J. J. O'Grady, AND R. Frankham. 2002.
30
31 545 Critiques of PVA ask the wrong questions: throwing the heuristic baby out with the
32
33 546 numerical bath water. *Conservation Biology* 16:262-263.
- 34
35 547 Caswell, H. 2001. *Matrix population models*. Sinauer Associates, Inc., Sunderland,
36
37 548 Massachusetts.
- 38
39 549 Chapman, J. A., AND G. A. Feldhamer. 1981. *Sylvilagus aquaticus*. *Mammalian Species*
40
41 550 151:1-4.
- 42
43 551 Chapman, J. A., AND D. E. C. Trethewey. 1972. Movements within a population of
44
45 552 introduced eastern cottontail rabbits. *Journal of Wildlife Management* 36:155-158.
- 46
47 553 Conaway, C. H., T. S. Baskett, AND J. E. Toll. 1960. Embryo resorption in the swamp rabbit.
48
49 554 *Journal of Wildlife Management* 24:197-202.
- 50
51 555 Crawford, J. C. 2014. Ecology of the swamp rabbit and eastern cottontail in bottomland
52
53 556 hardwood forests in southern Illinois. Ph.D. Dissertation, Southern Illinois
54
55 557 University, Carbondale.

- 1
2
3
4
5
6
7 558 Dailey, T. V., T. M. Vangilder, AND L. W. Burger, Jr. 1993. Swamp rabbit distribution in
8
9 559 Missouri. Proceedings of the Annual Conference for the Association of Southeastern
10
11 560 Fish and Wildlife Agencies 47:251-256.
- 12 561 Dooley, J. L., Jr., AND M. A. Bowers. 1998. Demographic responses to habitat fragmentation:
13
14 562 experimental tests at the landscape and patch scale. Ecology 79:696-980.
- 15
16 563 Dunning, J. B., R. Borgella, K. Clements, AND G. K. Meffe. 1995. Patch isolation, corridor
17
18 564 effects, and colonization by a resident sparrow in a managed pine woodland.
19
20 565 Conservation Biology 9:542-550.
- 21
22 566 Ellner, S. P., J. Fieberg, D. Ludwig, AND C. Wilcox. 2002. Precision of population viability
23
24 567 analysis. Conservation Biology 16:258-261.
- 25
26 568 ESRI. 2012. ArcGIS. Ver. 10.1. Environmental System Research Institute, Inc. Redlands,
27
28 569 California.
- 29
30 570 Fahrig, L., AND G. Merriam. 2003. Habitat patch connectivity and population survival.
31
32 571 Ecology 66:1762-1768.
- 33
34 572 Fenderson, L. E., A. I. Kovach, J. A. Litvaitis, K. M. O'Brien, K. M. Boland, AND W. J.
35
36 573 Jakubas. 2014. A multiscale analysis of gene flow for the New England cottontail, an
37
38 574 imperiled habitat specialist in a fragmented landscape. Ecology and Evolution
39
40 575 4:1853-1875.
- 41 576 Forsys, E. A. 1995. Metapopulations of marsh rabbits: a population viability analysis of the
42
43 577 Lower Keys marsh rabbit (*Sylvilagus palustris hefneri*). Ph. D. dissertation,
44
45 578 University of Florida, Gainesville, USA.
- 46
47 579 Forsys, E. A., AND S. R. Humphrey. 1996. Home range and movements of the Lower Keys
48
49 580 marsh rabbit in a highly fragmented habitat. Journal of Mammalogy 77:1042-1048.
- 50
51 581 Fowler, A., AND R. E. Kissell, Jr. 2007. Winter relative abundance and habitat associations of
52
53 582 swamp rabbits in eastern Arkansas. Southeastern Naturalist 6:247-258.

- 1
2
3
4
5
6
7 583 Ginzburg, L. R., L. B. Slobodkin, K. Johnson, and A. G. Bindman. 1982. Quasiextinction
8 probabilities as a measure of impact on population growth. *Risk Analysis* 21:171-
9 584 181.
10
11
12 586 Haag, T., A. S. Santos, D. A. Sana, R. G. Morato, L. Cullen, Jr., P. G. Crawshaw, Jr., C. De
13
14 587 Angelo, D. S. Di Bitetti, F. M. Salzano, AND E. Eizirik. 2010. The effect of habitat
15
16 588 fragmentation on the genetic structure of a top predator: loss of diversity and high
17
18 589 differentiation among remnant populations of Atlantic Forest jaguars (*Panthera*
19
20 590 *onca*). *Molecular Ecology* 19:4906-4921.
21
22 591 Haddad, N. M. AND K. A. Baum. 1999. An experimental test of corridor effects on butterfly
23
24 592 densities. *Ecological Applications* 9:623-633.
25
26 593 Hamilton, L. P., P. A. Kelly, D. F. Williams, D. A. Kelt, AND H. U. Wittmer. 2010. Factors
27
28 594 associated with survival of reintroduced riparian brush rabbits in California.
29
30 595 *Biological Conservation* 143:999-1007.
31
32 596 Hanski, I., and M. Gilpin. 1991. Metapopulation dynamics: a brief history and conceptual
33
34 597 domain. *Biological Journal of the Linnaean Society* 42:3-16.
35
36 598 Hill, E. P., III. 1967. Notes on the life history of the swamp rabbit in Alabama. *Proceedings*
37
38 599 of the Annual Conference of Southeastern Association of Game and Fish
39
40 600 Commissioners 21:117-123.
41
42 601 Hobbs, R. J. 1992. The role of corridors in conservation: solution or bandwagon? *Trends in*
43
44 602 *Ecology and Evolution* 7:389-392.
45
46 603 Hooctor, T. S., M. H. Carr, AND P. D. Zwick. 2000. Identifying a linked reserve system using
47
48 604 a regional landscape approach: The Florida Ecological Network. *Conservation*
49
50 605 *Biology* 14:984-1000.
51
52 606 Holler, N. S., T. S. Baskett, AND J. P. Rogers. 1963. Reproduction in confined swamp rabbits.
53
54 607 *Journal of Wildlife Management* 27:179-183.
55
56
57
58
59
60

- 1
2
3
4
5
6
7 608 Holmes, E. E. 2001. Estimating risks in declining populations with poor data. Proceedings of
8
9 609 the National Academy of Sciences of the United States of America 98:5072-5077.
10
11 610 Hunt, T. P. 1959. Breeding habits of the swamp rabbit with notes on its life history. Journal
12
13 611 of Mammalogy 40:82-91.
14
15 612 Indiana Department of Natural Resources. 2013. Division of Fish and Wildlife. Nongame &
16
17 613 Endangered Wildlife. <http://www.in.gov/dnr/fishwild/2356.htm>. Accessed 5 March
18
19 614 2013.
20
21 615 Johst, K., R. Brandl, AND S. Eber. 2002. Metapopulation persistence in dynamic landscapes:
22
23 616 the role of dispersal distance. *Oikos* 98:263-270.
24
25 617 Kirkpatrick, R. L., and D. M. Baldwin. 1974. Population density and reproduction in penned
26
27 618 cottontail rabbits. *Journal of Wildlife Management* 38:482-487.
28
29 619 Kjolhaug, M. S. 1986. Status, distribution, and factors determining habitat quality of the
30
31 620 swamp rabbit in Illinois. M. S. thesis, Southern Illinois University Carbondale, USA.
32
33 621 Kjolhaug, M. S., A. Woolf, AND W. D. Klimstra. 1987. Current status and distribution of the
34
35 622 swamp rabbit in Illinois. *Transactions of the Illinois Academy of Science* 80:299-308.
36
37 623 Kovacs, J. M., J. Malczewski, AND F. F. Verdugo. 2004. Examining local ecological
38
39 624 knowledge of hurricane impacts in a mangrove forest using an analytical hierarchy
40
41 625 process (AHP) approach. *Journal of Coastal Research* 20:792-800.
42
43 626 Krauss, J., R. Bommarco, M. Guardiola, R. K. Heikkinen, A. Helm, M. Kuussaari, R.
44
45 627 Lindborg, E. Öckinger, M. Pärtel, J. Pino, J. Pöyry, K. M. Raatikainen, A. Sang, C.
46
47 628 Stefanescu, T. Teder, M. Zobel, AND I. Steffan-Dewenter. 2010. Habitat
48
49 629 fragmentation causes immediate and time-delayed biodiversity loss at different
50
51 630 trophic levels. *Ecology Letters* 13:597-605.
52
53 631 LaFever, D. H., P. M. Schmidt, N. D. Perry, C. A. Faulhaber, R. R. Lopez, N. J. Silvy, AND
54
55 632 E. A. Forsys. 2008. Use of a population viability analysis to evaluate human-induced

- 1
2
3
4
5
6
7 633 impacts and mitigation for the endangered Lower Keys marsh rabbit. *Human-Wildlife*
8 634 *Conflicts* 2:260-269.
- 9
10 635 LaRue, M. A., AND C. K. Nielsen. 2008. Modelling potential dispersal corridors for cougars
11
12 636 in midwestern North America using least-cost path methods. *Ecological Modelling*
13 637 212:372-381.
- 14
15
16 638 LaRue, M. A., AND C. K. Nielsen. 2011. Modelling potential habitat for cougars in
17
18 639 midwestern North America. *Ecological Modelling* 222:897-900.
- 19
20 640 Lorimer, C. G. 2001. Historical and ecological roles of disturbance in eastern North
21
22 641 American forests: 9,000 years of change. *Wildlife Society Bulletin* 29:425-439.
- 23
24 642 Ludwig, D., AND C. J. Walters. 2002. Fitting population viability analysis into adaptive
25
26 643 management. Pp. 511-520 in *Population viability analysis* (S. R. Beissinger and D. R.
27 644 McCullough, eds.). University of Chicago Press, Chicago, Illinois.
- 28
29
30 645 McCarthy, M. A., AND L. S. Broome. 2000. A method for validating stochastic models of
31
32 646 population viability: a case study of the mountain pygmy-possum (*Burramys parvus*).
33 647 *Journal of Animal Ecology* 69:599-607.
- 34
35
36 648 Medici, E. P., AND A. L. J. Desbiez. 2012. Population viability analysis: using a modeling
37
38 649 tool to assess the viability of tapir populations in fragmented landscapes. *Integrative*
39 650 *Zoology* 7:356-372.
- 40
41 651 Moilanen, A., A. T. Smith, AND I. Hanski. 1998. Long-term dynamics in a metapopulation
42
43 652 of the American pika. *American Naturalist* 152:530-542.
- 44
45 653 Morris, W. F., P. L. Bloch, B. R. Hudgens, L. C. Moyle, AND J. R. Stinchcombe. 2002.
46
47 654 Population viability analysis in endangered species recovery plans: past use and future
48
49 655 improvements. *Ecological Applications* 12:708-712.
- 50
51 656 O'Grady, J. J., D. H. Reed, B. W. Brook, AND R. Frankham. 2004. What are the best
52
53 657 correlates of predicted extinction risk? *Biological Conservation* 118:513-520.
- 54
55
56
57
58
59
60

- 1
2
3
4
5
6
7 658 Pinter, N., B. S. Ickes, J. H. Wlosinski, AND R. R. van der Ploeg. 2006. Trends in flood
8 stages: contrasting results from the Mississippi and Rhine River systems. Journal of
9 659 Hydrology 331:554-566.
10 660
11
12 661 Platt, S. G., AND M. Bunch. 2000. Distribution and status of the swamp rabbit in South
13 Carolina. Proceedings of the Annual Conference for the Association of Southeastern
14 662 Fish and Wildlife Agencies 54:408-414.
15 663
16
17 664 Porath, W. J. 1997. Swamp rabbit (*Sylvilagus aquaticus*) status, distribution, and habitat
18 characteristics in southern Illinois. M. S. thesis, Southern Illinois University,
19 Carbondale, USA.
20 665
21
22 666
23
24 667 Possingham, H. P., D. B. Lindenmayer, AND G. N. Tuck. 2002. Decision theory for
25 population viability analysis. Pp. 470-489 in Population viability analysis (S. R.
26 668 Beissinger and D. R. McCullough, eds.). University of Chicago Press, Chicago,
27 669 Illinois.
28
29 670
30
31 671 Ralls, K., S. R. Beissinger, AND J. F. Cochrane. 2002. Guidelines for using population
32 viability analysis in endangered-species management. Pp. 521-550 in Population
33 672 viability analysis (S. R. Beissinger and D. R. McCullough, eds.). University of
34 673 Chicago Press, Chicago, Illinois.
35
36 674
37
38 675 Ratcliffe, N., S. Schmitt, AND M. Whiffin. 2005. Sink or swim? Viability of a black-tailed
39 godwit population in relation to flooding. Journal of Applied Ecology 42:834-843.
40 676
41
42 677 Rolstad, J. 1991. Consequences of forest fragmentation for the dynamics of bird populations:
43 conceptual issues and the evidence. Biological Journal of the Linnean Society
44 678 42:149-163.
45 679
46
47 680 Roy Nielsen, C. L., S. M. Wakamiya, AND C. K. Nielsen. 2008. Viability and patch
48 occupancy of a swamp rabbit metapopulation at the northern edge of its distribution.
49 681 Biological Conservation 141:1043-1054.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7 683 Rubert, L. 2007. Identifying habitat suitability and potential linkages for swamp rabbit
8 conservation in southern Illinois. M. S. thesis, Southern Illinois University
9 Carbondale, USA.
10 685
11
12 686 Saaty, T. 1980. The analytical hierarchy process: planning, setting priorities, resource
13 allocation. McGraw-Hill, New York, New York.
14 687
15
16 688 Scharine, P. D., C. K. Nielsen, E. M. Schaubert, AND L. Rubert. 2009. Swamp rabbits in
17 floodplain ecosystems: influence of landscape- and stand-level habitat on relative
18 689 abundance. *Wetlands* 29:615-623.
19 690
20
21 691 Scharine, P. D., C. K. Nielsen, E. M. Schaubert, L. Rubert, AND J. C. Crawford. 2011.
22 Occupancy, detection, and habitat associations of sympatric lagomorphs in early-
23 692 successional bottomland forests. *Journal of Mammalogy* 92:880-890.
24 693
25
26 694 Scheibe, J. S., and R. Henson. 2003. The distribution of swamp rabbits in southeast
27 Missouri. *Southeastern Naturalist* 2:327-334.
28 695
29
30 696 Shields, P. W. 1960. Movement patterns of brush rabbits in northwestern California. *Journal*
31 of *Wildlife Management* 24:381-386.
32 697
33
34 698 Simberloff, D. AND J. Cox. 1987. Consequences and costs of conservation corridors.
35 *Conservation Biology* 6:63-70.
36 699
37
38 700 Simberloff, D., J. A. Farr, J. Cox, AND D. W. Mehlman. 1992. Movement corridors:
39 conservation bargains or poor investments? *Conservation Biology* 6:493-504.
40 701
41
42 702 Singleton, P. H., W. L. Gaines, AND J. F. Lehmkühl. 2002. Landscape permeability for large
43 carnivores in Washington: a geographic information system weighted distance and
44 703 least-cost corridor assessment. U. S. Department of Agriculture, Pacific Northwest
45 704 Research Station, Portland, Oregon, USA.
46 705
47
48 706 Smith, A. T., AND J. D. Nagy. 2015. Population resilience in an American pika (*Ochotona*
49 *princeps*) metapopulation. *Journal of Mammalogy* 96:394-404.
50 707
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7 708 Sole, J. D. 1994. Assessing swamp rabbit distribution in Kentucky. Proceedings of the
8
9 709 Annual Conference for the Association of Southeastern Fish and Wildlife Agencies
10
11 710 48:145-151.
- 12
13 711 Sorensen, M. F., J. P. Rogers, AND T. S. Baskett. 1968. Reproduction and development in
14
15 712 confined swamp rabbits. Journal of Wildlife Management 32:520-531.
- 16
17 713 Staples, D. F., M. L. Taper, AND B. Dennis. 2004. Estimating population trend and process
18
19 714 variation for PVA in the presence of sampling error. Ecology 85:923-929.
- 20
21 715 Starfield, A. M. 1997. A pragmatic approach to modeling for wildlife management. Journal
22
23 716 of Wildlife Management. 61:261-270.
- 24
25 717 Toll, J. E., T. S. Baskett, AND C. H. Conaway. 1960. Home range, reproduction, and foods of
26
27 718 the swamp rabbit in Missouri. American Midland Naturalist 63:398-412.
- 28
29 719 Trakhtenbrot, A., R. Nathan, G. Perry, AND D. M. Richardson. 2005. The importance of long-
30
31 720 distance dispersal in biodiversity conservation. Diversity and Distributions 11:173-
32
33 721 181.
- 34
35 722 United States Army Corps of Engineers. 2011. Louisville District Corps of Engineer. Water
36
37 723 levels of rivers and lakes.
38
39 724 <http://www2.mvr.usace.army.mil/WaterControl/new/layout.cfm>. Accessed 14 Dec
40
41 725 2011.
- 42
43 726 United States Department of Commerce. 2011. TIGER/Line data.
44
45 727 <http://www.census.gov/cgi-bin/geo/shapefiles2011/main>. Accessed 14 Dec 2011.
- 46
47 728 United States Fish and Wildlife. Recovery plan for upland species of the San Joaquin Valley,
48
49 729 California. 319 pp. http://ecos.fws.gov/docs/recovery_plans/1998/980930a.pdf.
50
51 730 Accessed 13 May 2015.

- 1
2
3
4
5
6
7 731 United States Geological Survey. 2001. Multi-Resolution Land Characteristics Consortium.
8
9 732 National Landcover Database. <http://www.mrlc.gov/nlcd2001.php>. Accessed 29 Nov
10
11 733 2011.
12
13 734 United States Geological Survey. 2006. Multi-Resolution Land Characteristics Consortium.
14
15 735 National Landcover Database. <http://www.mrlc.gov/nlcd2006.php>. Accessed 29 Nov
16
17 736 2011.
18
19 737 United States Geological Survey. 2008. Project LANDFIRE. LANDFIRE data distribution
20
21 738 site. <http://landfire.cr.usgs.gov/viewer/>. Accessed 30 Jan 2012.
22
23 739 United States Geological Survey. 2011. National Weather Information System. USGS
24
25 740 surface-water daily data for Illinois. <http://waterdata.usgs.gov/il/nwis/dv?>. Accessed
26
27 741 29 Nov 2011.
28
29 742 Vale, K. B. 2008. Spatial ecology of male swamp rabbits (*Sylvilagus aquaticus*) in
30
31 743 southeastern Arkansas. Thesis, University of Arkansas at Monticello, USA.
32
33 744 Wikramanayake, E., M. McKnight, E. Dinerstein, A. Joshi, B. Gurung, AND D. Smith. 2004.
34
35 745 Designing a conservation landscape for tigers in human-dominated environments.
36
37 746 Conservation Biology 18:839-44.
38
39 747 Wilby, R. L., AND R. Keenan. 2012. Adapting to flood risk under climate change. Progress in
40
41 748 Physical Geography 36:348-378.
42
43 749 Wilcove, D. S., C. H. McLellan, AND A. P. Dobson. 1997. Habitat fragmentation in the
44
45 750 temperate zone. Pp. 342-355 in The human impact reader (A. Goudie, ed.). Blackwell
46
47 751 Publishers, Malden, Massachusetts.
48
49 752 Woolf, A., AND M. S. Barbour. 2002. Population dynamics and status of the swamp rabbit in
50
51 753 Illinois. Final Report, Illinois Federal Aid Project W-106-R-12. Cooperative Wildlife
52
53 754 Research Laboratory, Southern Illinois University, Carbondale, Illinois.
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7 755 Ye, X. P., A. K. Skidmore, AND T. J. Wang. 2013. Within-patch habitat quality determines
8 the resilience of specialist species in fragmented landscapes. *Landscape Ecology*
9 756 28:135-147.
10 757
11
12 758 Zollner, P. A., W. P. Smith, AND L. A. Brennan. 2000a. Home range use by swamp rabbits
13 (*Sylvilagus aquaticus*) in a frequently inundated bottomland forest. *American Midland*
14 759 *Naturalist* 143:64-69.
15 760
16
17 761 Zollner, P. A., W. P. Smith, AND L. A. Brennan. 2000b. Microhabitat characteristics of sites
18 762 used by swamp rabbits. *Wildlife Society Bulletin* 28:1003-1011.
19
20 763 ***Submitted 8 September 2014. Accepted XXXXXXXX.***
21
22 764 ***Associate Editor was Harald Beck.***
23
24
25
26 765
27
28 766
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

767 **FIGURE LEGENDS**

768 Fig. 1. Mean importance weights with standard deviations calculated using the analytical
769 hierarchy process (Saaty 1980), representing the relative importance of each factor in the
770 habitat suitability model (solid bar) and the dispersal cost model (hatched bar) for swamp
771 rabbits (*Sylvilagus aquaticus*) in southern Illinois. A mean weight of 1 (dotted line) indicates
772 equal perceived importance relative to other factors.

773
774 Fig. 2. Maps of a) habitat patches ($n = 62$) identified as suitable or highly suitable and with
775 confirmed swamp rabbit (*Sylvilagus aquaticus*) presence in southern Illinois (inset) in 1983-
776 1985, 1995-1997, 2006-2007, or 2009-2011, and b) dispersal corridors (in black) connecting
777 suitable habitat patches with confirmed swamp rabbit presence (in gray) in the Cache River
778 watershed, southern Illinois.

779
780 Fig. 3. Minimum (circles), 5th percentile (squares), and median (50th percentile; diamonds)
781 simulated metapopulation abundance as percentage of initial metapopulation size for swamp
782 rabbits (*Sylvilagus aquaticus*) in southern Illinois over 50 years, based on intermediate
783 parameter values.

784
785 Fig. 4. Probability of quasi-extinction for swamp rabbits (*Sylvilagus aquaticus*) in southern
786 Illinois within 25 years following detrimental changes in fecundity (decreased fecundity;
787 diamonds) and effect of catastrophes (increased impact on populations; squares). Dashed line
788 represents the upper 95% confidence interval around the probability of quasi-extinction (0%)
789 from the initial test using all intermediate values.

790

791 Table 1. Importance of habitat characteristics for swamp rabbits (*Sylvilagus*
 792 *aquaticus*), calculated from expert surveys using the analytical hierarchy process (Saaty
 793 1980). Values are mean weights ($\pm SD$), representing the relative suitability of each attribute
 794 within the factors used in mapping habitat suitability and dispersal resistance for swamp
 795 rabbits in southern Illinois. Bold indicates the attribute with highest suitability or dispersal
 796 resistance within each factor.

| Factor | Attribute | Suitability | Dispersal resistance |
|--------------|-----------------------------|-----------------------------------|-----------------------------------|
| Land cover | Open water/barren/developed | 0.16 \pm 0.02 | 2.68 \pm 0.49 |
| | Agriculture | 0.24 \pm 0.06 | 1.75 \pm 0.58 |
| | Upland forest | 0.39 \pm 0.11 | 0.88 \pm 0.07 |
| | Upland shrub/scrub | 0.54 \pm 0.08 | 0.83 \pm 0.22 |
| | Upland herbaceous | 0.50 \pm 0.18 | 1.15 \pm 0.47 |
| | Wetland forest | 2.26 \pm 0.63 | 0.23 \pm 0.07 |
| | Wetland shrub/scrub | 2.11 \pm 0.57 | 0.20 \pm 0.05 |
| | Wetland herbaceous | 1.80 \pm 0.49 | 0.29 \pm 0.10 |
| Canopy cover | 0-25% | 0.24 \pm 0.06 | 1.93 \pm 0.81 |
| | 26-50% | 0.91 \pm 0.50 | 0.89 \pm 0.35 |
| | 51-75% | 1.36 \pm 0.36 | 0.64 \pm 0.60 |

| | | | | |
|--------------------------|--|---------------------------|--------------------|--------------------|
| | | 76-100% | 1.49 ± 0.75 | 0.54 ± 0.37 |
| Streams and water bodies | | Intermittent shoreline | 0.58 ± 0.48 | 1.16 ± 0.64 |
| | | Perennial shoreline | 1.14 ± 0.56 | 0.89 ± 0.57 |
| | | Intermittent stream/ditch | 0.64 ± 0.31 | 1.01 ± 0.54 |
| | | Perennial stream/ditch | 1.58 ± 0.44 | 0.81 ± 0.50 |
| | | Intermittent lake/pond | 0.76 ± 0.50 | 1.25 ± 0.65 |
| | | Perennial lake/pond | 1.31 ± 0.88 | 0.87 ± 0.94 |
| Roads | | Highways | 0.22 ± 0.05 | 1.81 ± 0.79 |
| | | Secondary | 0.45 ± 0.07 | 0.96 ± 0.28 |
| | | Local/rural | 0.82 ± 0.18 | 0.64 ± 0.20 |
| | | Unpaved | 2.50 ± 0.20 | 0.59 ± 0.92 |

797

798

799 Table 2. Parameter values used in the initial population viability analysis
 800 (intermediate values) and sensitivity tests for swamp rabbits (*Sylvilagus aquaticus*) in
 801 southern Illinois. Effect of catastrophe values are the percent of patch abundance lost in a
 802 flood, and trend in carrying capacity values are year-25 carrying capacity (K_{25}) as a
 803 percentage of K_0 .


| Parameter | Minimum | Intermediate | Maximum |
|---|---------|--------------|---------|
| Home Range (ha) | 0.965 | 1.93 | 2.895 |
| Survival | 0.15 | 0.3 | 0.45 |
| Fecundity | 0.672 | 1.344 | 2.016 |
| Occurrence Rate of Catastrophes | 5% | 10% | 15% |
| Effect of Catastrophes ^a | 30% | 60% | 90% |
| Average Dispersal Distance (m) | 100 | 200 | 300 |
| Maximum Dispersal Rate | 5% | 10% | 15% |
| Dispersal Parameter c | 0.5 | 1 | 1.5 |
| Trend in Carrying Capacity ^b | 50% | 100% | 150% |

805
 806 ^aPercentage of patch population lost during flood.

807 ^bValues indicate carrying capacity in year 25 as a percentage of carrying capacity in year 0.


808 100% indicates no trend.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



278x215mm (300 x 300 DPI)

Only



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

