

Pharmacological Activities of *Mentha piperita*- Mini Review

S. Meenatchisundaram^{1*}, G. Parameswari¹, Diana Sunny¹, M. Brinda², T. Subbraj¹, T. Suganya¹ and A. Michael¹

¹Department of Microbiology, PSG College of Arts and Science, Coimbatore, India

²Department of Microbiology, GRD College of Science, Coimbatore, India

*Corresponding author: Dr.S.Meenatchisundaram, E-Mail: drmscbe@gmail.com

Issued 30 January 2009

Peppermint (*Mentha × piperita*) is a hybrid mint, a cross between watermint (*Mentha aquatica*) and spearmint (*Mentha spicata*). The plant is indigenous in Europe and now widespread in cultivation throughout all regions of the world. It is found wild occasionally with its parent species

Pharmacological Activities of *Mentha piperita*

Antimicrobial activity

Extracts of Folium Menthae Piperitae have antibacterial and antiviral activity in vitro. Addition of ground leaves to the agar medium inhibited the growth of *Salmonella typhimurium*, *Staphylococcus aureus* and *Vibrio parahaemolyticus* at concentrations of 0.1–2.0% (w/v). Aqueous and ethanol extracts of the leaves reduced the number of plaques of the rinderpest virus at concentrations of 4–8mg/ml¹. Aqueous extracts of the leaves demonstrated activity against the following viruses in egg and cell culture: Newcastle disease, herpes simplex, vaccinia, Semliki Forest and West Nile².

Smooth muscle contraction

A 31% ethanol extract of the leaves inhibited both acetylcholine- and histamine-induced smooth muscle contractions in guinea-pig ileum in vitro at a concentration of 10 ml/l. The results were similar to those obtained with 0.13 mg atropine^{3,4}. An aqueous flavonoid fraction isolated from a leaf extract inhibited barium chloride-induced muscle contractions of guinea-pig ileum in vitro at a concentration corresponding to 0.5 g leaves/ml⁵.

Choleretic activity

Injection of a leaf infusion (0.5 ml) or a flavonoid fraction (equivalent to 3.3 g leaves/kg body weight) increased the amount of bile acids in cannulated rats and dogs (dose 0.4 mg/kg body weight)^{5,6}. A mixture of flavonoids, isolated from the leaves, had choleric activity in dogs (2mg/kg body weight)⁷. Flavomentin, a flavonoid isolated from the leaves, stimulated bile secretion and the synthesis of bile acids in dogs (2mg/kg body weight)⁸. Intra-gastric administration of a 30% ethanol extract of the leaves to rats (1ml/kg body weight) increased bile flow by 43%. The extract did not induce sedation in mice at doses up to 10 ml/kg body weight⁴.

Anti-oedema activity

Topical application of a methanol leaf extract to mice (2.0mg/ear) inhibited ear oedema induced by 12-*O*-tetradecanoylphorbol-13-acetate⁹.

Analgesic activity

Intra-gastric administration of a 30% ethanol extract of the leaves inhibited phenylbenzoquinone-induced writhing in mice (ED₅₀ 2.1 ml/kg body weight)⁴.

Toxicology

Intra-gastric administration of a leaf extract (50 g leaves infused with 500 ml hot water for 10 minutes, then spray-dried) to 12 mice (4 g/kg body weight as a single dose) did not result in central nervous system depression, toxic effects or mortality¹⁰.

References

1. Alwan AH. Antiviral activity of some Iraqi indigenous plants. *International Journal of Crude Drug Research*, 1988, 2:107–111.
2. Herrmann EC Jr, Kucera LS. Antiviral substances in plants of the mint family (Labiatae). III. Peppermint (*Mentha piperita*) and other mint plants. *Proceedings of the Society for Experimental Biology and Medicine*, 1967:874–878.
3. Forster HB. Antispasmodic effects of some medicinal plants. *Planta Medica*, 1980, 40:309–319.
4. Leslie GB. A pharmacometric evaluation of nine Bio-Strath herbal remedies. *Medita*, 1978, 8:3–19.
5. Lallement-Guilbert N, Bézanger-Beauquesne L. Recherches sur les flavonoides

quelques Labiées médicinales (romarin, menthe poivrée, sauge officinale). *Plantes médicinales et Phytothérapie*, 1970, 4:92–107.

6. Steinmetzer K. Experimentelle Untersuchungen über Cholagoga. *Wiener Klinische Wochenschrift*, 1926, 39:1418–1422, 1455–1457.
7. Pasechnik IK. Study of choleric properties specific to flavonoids from *Mentha piperita* leaves. *Farmakologiya Toksikologiya*, 1966, 21:735–737.
8. Pasechnik IK, Gella EV. Choleric preparation from peppermint. *Farmatsevtichnyi Zhurnal (Kiev)*, 1966, 21:49–53.
9. Yasukawa K. Inhibitory effect of edible plant extracts on 12-*O*-tetradecanoylphorbol-13-acetate-induced ear edema in mice. *Phytotherapy Research*, 1993, 7:185–189.
10. Della Loggia R. Evaluation of some pharmacological activities of a peppermint extract. *Fitoterapia*, 1990, 61:215–221.