

Glycyrrhizin- A Review


Dr Amrit Pal Singh, MD (Alternative Medicine), Medical Executive, India –Swift Ltd, Super Speciality Division, Chandigarh.

Address for correspondence:

Dr Amrit Pal Singh,
House No: 2101 Phase-7,
MOHALI -160062.
Email amritpal2101@yahoo.com

Chemistry

Glycyrrhizin is a triterpenoid saponin found in *Glycyrrhiza glabra* (licorice). Chemically, glycyrrhizin is a sulphated polysaccharide. It is considered to be the active constituent of the drug and the standardization of licorice is based on glycyrrhizin content. The standardized extracts of licorice sold in the market contain 20% of glycyrrhizin. Glycyrrhizin is converted into glycyrrhetic acid by an enzyme, glycyrrhonase.



Structure of Glycyrrhizin.

Pharmacological activities

A. Hepatoprotective activity:

Glycyrrhizin has demonstrated hepatoprotective activity in animal models against carbon tetrachloride induced toxicity. It reduces alanine transaminase and aspartate transaminase values in serum. The exact mode of action is not clear but it has been proposed that glycyrrhizin has inhibitory effect on immune mediated cytotoxicity against hepatocytes and on nuclear factor (NF)-kappa B, which activates genes encoding inflammatory cytokines in the liver. In addition glycyrrhizin has shown activity in hepatitis and cirrhosis.

B. Anti-viral activity

It has been proven to have antiviral activity against DNA and RNA virus (VZV, HIV, Influenza A and B, types 1 and 2 herpes simplex and hepatitis B and C) in vitro and in vivo. Toxicity has not been found in experimental trials on different cell lines. The Glycyrrhizin acid blocks the first steps on viral replication as well as the viron exit from the capsid.

1. Glycyrrhizin and herpes simplex

In an open, controlled, clinical and multicentric study, glycyrrhizin was evaluated for efficacy, safety and tolerability. 52 adults patients of both sexes (males 62.7%, females 37.3%) with age ranging from 18 to 65 years (mean 38.8) with active lesions of genital herpes simplex were included in this preliminary study. Patients the cases were enrolled under clinical, cytologic and immunologic criteria. A local application of the spray solution on the affected area was done twice a day for 5 days. Clinical efficacy was evaluated with no parametric methods. Total efficacy was observed in 95%, it was excellent in 65%, good in 25% and mild in 9%. The natural history was the same, but not so good were the control of pain and pruritus. The immuno-serological evolution was correlated with the clinical history. Local tolerability was of 94.1% and non-important adverse side events were observed.

2. Glycyrrhizin and Anti HIV activity

Anti HIV activity of glycyrrhizin was first explained IN November 1986 in Aids treatment news. After this the other compounds belonging to the category of glycyrrhizin were investigated for Anti HIV activity. A study from Japan explained the use of glycyrrhizin in 10 patients, who were given glycyrrhizin ranging from 150 to 225mg. At the end it was concluded that glycyrrhizin was more effective than placebo in preventing the occurrence of symptoms.

3. Anti-inflammatory effects

The anti-inflammatory activity of glycyrrhizin has been explained by number of researchers but very limited data is available. The activity has been compared with glucocorticoid hormone. The mode of action is slightly different from the hormone. It has been proposed that glycyrrhizin promotes regeneration of inflammatory tissue, whereas glucocorticoid inhibits the regeneration.

References:

1. William Procter Jr, On glycyrrhizin. America Journal of Pharmacy. Vol. XLIII, 1871.
2. P. Bean. Silymarin and glycyrrhizin are best-known phytomedicine for hepatitis C. Am Clin Lab 2002 May; 21(4);

19-21.

3. Numazaki, K., et al. Effect of glycyrrhizin in children with liver dysfunction associated with cytomegalovirus infection. *Tohoku J Exp Med.* 1994 Feb; 172(2); 147-53.
4. Vick FR, Hidalgo LH, Zenon MC, Martinez S. Local application of glycyrrhizin acid in genital herpes. *Rev Hosp M Gea Glz* 2000; 3(4); 141-144.
5. Yamamura Y, Kotaki H, Tanaka N, Aikawa T, Sawada Y, Iga T. The pharmacokinetics of glycyrrhizin and its restorative effect on hepatic function in patients with chronic hepatitis and in chronically carbon-tetrachloride-intoxicated rats. *Biopharm Drug Dispos* 1997; 18:717-72.
6. Ito, M., et al. Mechanism of inhibitory effect of glycyrrhizin on replication of human immunodeficiency virus. *Antiviral Res.* 1988, Dec 11; 10(6): 289-98.